

















Wild Plants of Sycamore Valley Regional Open Space Preserve

Common Name Version

A Photographic Guide

Sorted by Form, Color and Family with Habitat Descriptions and Identification Notes



New Revised and Expanded Edition - Includes the latest scientific names, habitat descriptions and identification notes

Decimal Inches



English Inches

Notes:

A Photographic Guide to the Wild Plants of

Sycamore Valley Regional Open Space Preserve

More than 2,000 species of native and naturalized plants grow wild in the San Francisco Bay Area. Most are very difficult to identify without the help of good illustrations. This is designed to be a simple, color photo guide to help you identify some of these plants. This guide is published electronically in Adobe Acrobat® format so that it can easily be updated as additional photographs become available. You have permission to freely download, distribute and print this guide for individual use. Photographs are © 2014 Wilde Legard, all rights reserved.

In this guide, the included plants are sorted first by form (Ferns & Fern-like, Grasses & Grass-like, Herbaceous, Woody), then by most common flower color, and finally by similar looking flowers (grouped by genus within each family). Each photograph has the following information, separated by '-':

COMMON NAME According to The Jepson Manual: Vascular Plants of California, Second Edition (JM2) and other references (not standardized).

(Scientific Name) According to JM2 and eFlora (ucjeps.berkeley.edu/IJM.html).

Origin & Longevity Native, Naturalized, or Waif (not reproducing without human care). Annual, Biennial, Perennial, or a combination.

Family Name Common family name according to *JM2*,

(Bloom date range) Period during the year when the plant blooms, according to JM2 and other sources. '-' if plant does not bloom (ie. Ferns).

Habitat Habitat description according to *JM2* and other sources.

ID Characteristics Plant description with identification characteristics and other notes, based on multiple sources including: Annotated Checklist of the East Bay Flora, Second

Edition (2013), JM2, Plants of the San Francisco Bay Region (Revised Edition), and Weeds of California and Other Western States.

Additional notes Occasionally, an additional note may appear (ie. NOXIOUS weed, INVASIVE weed, Fed & Calif. ENDANGERED, etc.).

Revision: 3/2/2014 Grass-like - Green/Brown

Wild Plants of Sycamore Valley Regional Open Space Preserve - Sorted by Form, Color and Family



BROAD-LEAVED CATTAIL (Typha latifolia) Native Perennial - Cattail Family - (Jun–Jul) - Unpolluted to nutrient-rich freshwater (brackish) marshes - Plant 4.9-9.8' tall. Widest leaves 0.4-1.1" wide. Flower cluster with no gap between flower types.



SLENDER WILD OAT (Avena barbata)
Naturalized Annual - Grass Family - (Mar—Jun) Disturbed sites - Plants gen 24-32" Spikelets
0.8-1.2" long. Awns 0.8-1.8" long. Lemma tip
bristles >= 0.1" long. Seeds EDIBLE whole or
around for flour. INVASIVE weed.



WILD OAT (Avena fatua) Naturalized Annual - Grass Family - (Apr–Jun) - Disturbed sites - Plants 1-5' tall. Spikelets 0.7-1.3" long. Low awn 1-1.6" long. Lemma tip bristles < 0.04" long. Seeds EDIBLE whole or ground for flour. INVASIVE weed.



Page 1

CALIFORNIA BROME (Bromus carinatus var. carinatus) Native Perennial - Grass Family - (Apr-Aug) - Coastal prairies, openings in chaparral, plains, open oak and pine woodland -Plant 20-40" tall. Flower cluster 6-16" long. Spikelet 0.8-1.6" long. Lemma 0.5-0.8" long, hairy, awn 0.3-0.6" long.



RIPGUT GRASS (Bromus diandrus) Naturalized Annual - Grass Family - (Apr-Jul) - Open, gen disturbed areas - Plant 6-40" tall. Spikelet 1-2.8" long. Lemma body 0.8-1.2" long, awn > 1.2" long. Barbed seeds can stick in flesh of animals. INVASIVE weed.



SOFT CHESS (Bromus hordeaceus) Naturalized Annual - Grass Family - (Apr–Jul) - Fields, disturbed areas - Plant 4-26" tall. Leaf hairy. Flower cluster 1-5" long, dense, some stalks > spikelet. Spikelet 0.5-0.9". Lemma 0.26-0.4", awn 0.16-0.4". INVASIVE weed.



FOXTAIL CHESS (Bromus madritensis subsp. madritensis) Naturalized Annual - Grass Family -(Apr-Jan) - Disturbed areas, roadsides - Plants 4-20" tall. Stem and sheathes smooth. Flower cluster branches visible, lower spikelets erect, >



RED BROME (Bromus madritensis subsp. rubens) Naturalized Annual - Grass Family -(Mar-Jun) - Disturbed areas, roadsides - Plant (Wal-Sun) - Distribute a reas, readstates - Flath 4-20". Flower cluster condensed, branches obscure, < spikelets. Stem & sheathes hairy. INVASIVE weed.



BEARDLESS WILD RYE (Elymus triticoides) Native Perennial - Grass Family - (Jun–Jul) - Dry to moist, often saline, meadows - Plant 18-50" tall, from rhizomes. Flower cluster 2-8" long. Spikelets generally 2/node. Lemmas 3-7, 0.2-0.5" long, awn to 0.12" long.



BROME FESCUE (Festuca bromoides)
Naturalized Annual - Grass Family - (May–Jun) Uncommon. Dry, disturbed places, coastal-sage
scrub, chaparral - Stem < 20". Inflor 0.6-6" tall,
dense, lower branches erect. Spikelet 0.2-0.4" long, awn 0.1-0.3" long.



RYE GRASS (Festuca perennis) Naturalized Perennial - Grass Family - (May-Sep) - Dry to moist disturbed sites, abandoned fields - Stem 20-40" tall. Spikelet 0.2-0.9" long, > glume, awned or awnless. Sterile shoots at base. INVASIVE weed.




MEDITERRANEAN BARLEY (Hordeum marinum subsp. gussoneanum) Naturalized Annual - Grass Family - (Apr-Jun) - Dry to moist, disturbed sites - Stem 4-20". Leaf blades to 32" long, auricles < 0.08". Central lemma awn 0.25-0.7" long. INVASIVE weed.


HARE BARLEY (Hordeum murinum subsp. leporinum) Naturalized Annual - Grass Family - (Feb-May) - Moist, gen disturbed sites. Common - Stem 12-43" tall. Central spikelet stalk ~0.06" long. Central floret << lateral florets. INVASIVE weed.



WALL BARLEY (Hordeum murinum subsp. murinum) Naturalized Annual - Grass Family - (Feb-May) - Moist, gen disturbed sites - Stem 1-2'. Leaf auricles notable. Central spikelet stalk 0-0.02". Central floret gen = lateral florets. INVASIVE weed.



HARDING GRASS (Phalaris aquatica) Naturalized Perennial - Grass Family - (Apr–Aug) - Disturbed areas, roadsides - Plant 16-79" tall, tufted. Flower cluster 0.6-6" long, unbranched. Lower florets 1 or or unequal. Glume wing untoothed. INVASIVE weed.



RABBITFOOT GRASS (Polypogon monspeliensis) Naturalized Annual - Grass Family - (Apr-Aug) - Moist places, along streams - Stem 8-39". Leaf 0.4-8" long, 0.16-0.24" wide. Flower cluster 0.4-6.7" long, plume-like. Glume awn > 0.14". Lemma awn gen < 0.1". INVASIVE.



FOOTHILL NEEDLE GRASS (Stipa lepida) Native Perennial - Grass Family - (Mar–Jun) - Dry slopes, chaparral, grassland, savanna, coastal scrub - Stem 14-39" tall. Leaf blade 4.7-9.1" long, narrow. Glumes 0.2-0.6" long. Awn 0.8-2" long.



COMMON TULE (Schoenoplectus acutus var. occidentalis) Native Perennial - Sedge Family - (Summer) - Common. Marshes, shores, fens, shallow lakes, often emergent - Stem 3.3-13' tall, fat, round x-section, blue-green.



ITHURIEL'S SPEAR (Triteleia laxa) Native Perennial - Brodiaea Family - (Apr–Jun) - Common. Open forest, conifer or foothill woodland, grassland on clay soil - Flower stem 4-28". Leaves 8-16", 0.16-1" wide. Flowers blue, blue-purple or white, 0.7-1.9" long.



MEDITERRANEAN LINSEED (Bellardia trixago) Naturalized Annual - Broom-rape Family - (Apr-Jun) - Disturbed grassland. - Plant sticky-hairy. Stem 6-32" tall. Leaves lance-shaped, toothed. Flowers 0.8-1" long, 2-lipped: upper lip pink, lower lip white. INVASIVE.



PURPLE OWL'S-CLOVER (Castilleja exserta subsp. exserta) Native Annual - Broom-rape Family - (Mar–May) - Open fields, grassland - Plant sticky, short-hairy. Flower cluster tipped white, pale yellow or rose. Flower upper lip hooked, fuzzy.



NARROW-LEAF MILKWEED (Asclepias fascicularis) Native Perennial - Dogbane Family - (May-Oct) - Dry ground, valleys, foothills - Plant gen hairless. Stem leaves narrow, 3-5 in whorls. Flowers green-white to purple-tinged, ~0.25" long.



SMALL CLARKIA (Clarkia affinis) Native Annual -Evening Primrose Family - (May-Jun) - Openings in woodland, chaparral - Stem < 32". Buds erect, ovary 8-grooved, sepals united on side. Petals 0.2-0.6" long, pale pink to dark wine-red, often purple-flecked.



PANICLED / WEEDY WILLOWHERB (Epilobium brachycarpum) Native Annual - Evening Primrose Family - (Jun-Sep) - Common. Dry open or disturbed woodland, grassland, roadsides - Plant 8-79" tall, diffuse. Petals 2-15 mm long, white to rose-purple.



LONG-BEAKED FILAREE (Erodium botrys) Naturalized Annual - Geranium Family -(Mar–Jul) - Dry, open or disturbed sites - Stem 4-35" tall, coarse-hairy. Leaves lobed. Petals pink, 0.3-0.5" long. Sepals w/reddish tip. Fruit beak 2-4.7" long.



REDSTEM FILAREE (Erodium cicutarium)
Naturalized Annual - Geranium Family (Feb-Sep) - Open, disturbed sites, grassland,
scrub - Stem 4-20". Leaves compound, leaflets
dissected. Sepal tip bristly. Petal pink to purple.
Fruit beak 0.8-2". INVASIVE.



GREENSTEM FILAREE (Erodium moschatum) Naturalized Annual - Geranium Family -(Feb-Sep) - Open, disturbed sites - Stem 4-24", short-hairy. Leaves compound, leaflets toothed. Sepal tip glabrous. Petals 0.4-0.6" long, pink. Fruit beak 0.8-1.6" long.



CUT-LEAVED GERANIUM (Geranium dissectum) Naturalized Annual - Geranium Family - (Mar–Jul) - Open, disturbed sites - Stem 3-28" tall, rough-hairy. Leaf blades deeply divided. Petals violet-red, 0.1-0.25" long w/notched tips. Flower stalk sticky. INVASIVE.



HAIRY DOVE'S FOOT GERANIUM (Geranium molle) Naturalized Annual - Geranium Family - (Feb-Aug) - Open to shaded sites, disturbed ground - Stem 4-17" tall. Petals red-purple, 0.1-0.4" long, notched. Sepals awnless. Fruit smooth, wrinkled.



WESTERN BLUE-EYED-GRASS (Sisyrinchium bellum) Native Perennial - Iris Family - (Mar–May) - Common. Open, gen moist, grassy areas, woodland - Stem < 25" tall. Leaves iris-like. Petals 6, blue-purple with dark veins, 0.4-0.7" long.



BULL MALLOW (Malva nicaeensis) Naturalized Annual - Mallow Family - (Mar–Jun) - Disturbed places - Stem 0.7-2'. Leaf blade 1.2-4.7" wide, 5-7 shallow lobes. Bractlets egg-shaped,0.16-0.2" long. Petals pink to blue-violet, 0.2-0.5" long.



MINIATURE LUPINE (Lupinus bicolor) Native Annual - Pea Family - (Mar–Jun) - Abundant. Open or disturbed areas - Plant 4-16", hairy. Flower 0.16-0.4" long, banner longer than wide, upper keel ciliate near tip, pedicel < 0.12". Fruit 0.12-0.24" wide.



PINPOINT CLOVER (Trifolium gracilentum) Native Annual - Pea Family - (Mar–Jun) - Open, disturbed places, occas serpentine - Leaflet tips not deep-notched. Reflexed pink-purple flowers w/point. Flower bracts completely smooth.



ROSE CLOVER (Trifolium hirtum) Naturalized Annual - Pea Family - (Apr–May) - Disturbed areas, roadsides - Head with 1-2 bract-like leaves immed below. Flowers pink, 0.4-0.6" long, densely-bristly in fruit. INVASIVE weed.



SPRING VETCH (Vicia sativa subsp. sativa) Naturalized Annual - Pea Family - (Mar–Jun) - Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands - Flowers 1-2 at leaf bases, pink-purple to white, 0.7-1.2" long. Leaflets 0.16-0.4" wide.



SPARSELY HAIRY VETCH (Vicia villosa subsp. varia) Naturalized Annual-Biennial - Pea Family - (Mar-Jun) - Grassland, roadside, disturbed areas - Stems w/few hairs. Flowers 10-20, blue-purple to white, 0.4-0.55" long. Lower bract lobes 0.04-0.1" long.



ITALIAN THISTLE (Carduus pycnocephalus subsp. pycnocephalus) Naturalized Annual -Sunflower Family - (Mar–Jul) - Roadsides, pastures, disturbed areas - Stems 8-79", narrow spiny-wings. Lower leaves 4-10 lobed, heads 2-5, flower bracts woolly. NOXIOUS.



BULL THISTLE (Cirsium vulgare) Naturalized Biennial - Sunflower Family - (May-Oct) - Common. Disturbed areas - Plant 12-79" tall. Top of leaves bristly; leaf base forming decurrent wings on stem. Flowers purple, gen 1-2" wide. NOXIOUS weed.



ARTICHOKE THISTLE (Cynara cardunculus subsp. flavescens) Naturalized Perennial - Sunflower Family - (Apr–Jul) - Disturbed places - Plant 1.6-8' tall. Artichoke head 1.6-6" diam, spine-tipped. Flowers blue or purple, 1.2-2" long. NOXIOUS weed.



MILK THISTLE (Silybum marianum) Naturalized Annual-Biennial - Sunflower Family - (Feb–Jun) - Roadsides, pastures, disturbed areas - Stem 1-10', stout. Leaves spiny, shiny green w/white veins and spots. Flowers red-purple. INVASIVE weed.



PURPLE SALSIFY (Tragopogon porrifolius) Naturalized Biennial-Perennial - Sunflower Family - (Mar-Nov) - Common. Disturbed places - Plant 16-39" tall, milky sap. Leaves 0.8-1.6" long, very narrow, waxy blue. Flowers purple.



ROBUST VERVAIN (Verbena lasiostachys var. scabrida) Native Perennial - Vervain Family - (May-Sep) - Open, dry to wet places - Plant 14-32" tall. Leaves green, narrow, 3-lobed, stiff-haired above. Flowers blue to purple on long



CURLY DOCK (Rumex crispus) Naturalized Perennial - Buckwheat Family - (All year) - Abundant. Disturbed places - Stem 16-39" tall. Flower cluster dense, valves 0.2-0.24", winged around tubercles, smooth edged. 1 tubercle enlarged. INVASIVE.



FIDDLE DOCK (Rumex pulcher) Naturalized Perennial - Buckwheat Family - (May–Sep) - Disturbed places, meadows, moist or dry habitats - Stem 8-24" long, branches widely spreading. Leaf blades 1.6-4" long, 1.2-2" wide. Valves w/2-5 teeth.



MUGWORT (Artemisia douglasiana) Native Perennial - Sunflower Family - (May–Nov) - Common. Open to shady areas, often in drainages - Plant 20-60" tall. Leaves gen 0.4-4" long, densely hairy below, some 3-5 lobed. Flower bracts hairy.



SPINY COCKLEBUR (Xanthium spinosum) Native Annual - Sunflower Family - (Jul-Oct) - Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses - Plant 4-24". Stem node spines 0.6-1.2" long, golden, gen 3-lobed. Bur 0.4-0.5" long.



COMMON FIDDLENECK (Amsinckia intermedia) Native Annual - Borage Family - (Mar–Jun) - Abundant. Open, generally disturbed places - Plant 0.5-3' tall. Flowers orange with 5 red spots, 0.3-0.4" long, 0.2-0.4" wide, tube straight.



SHEEP SORREL (Rumex acetosella) Naturalized Perennial - Buckwheat Family - (Apr-Jul) - ± Disturbed, often acidic places - Stem < 16". Leaf gen basal, blade 0.8-2.4" long, lower arrowhead-shaped w/2 lobes. Flowers yellowish, turn reddish. INVASIVE.



SCARLET PIMPERNEL (Anagallis arvensis) Naturalized Annual - Myrsine Family - (Mar–May) - Common. Disturbed places, ocean beaches - Plants 2-16" tall. Flowers 0.2-0.3" long, salmon or occasionally blue. Leaves TOXIC, can cause dermittis.



CALIFORNIA POPPY (Eschscholzia californica) Native Perennial - Poppy Family - (Feb-Sep) - Grassy, open areas - Plant 2-24" tall. Flower w/spreading rim <= 0.2". Petals 0.8-2.4" long, early spring = large and orange, fall = smaller and volley.



POISON HEMLOCK (Conium maculatum) Naturalized Biennial - Carrot Family - (Apr-Jul) - Common. Moist, esp disturbed places - Plants 2-10' tall. Stems purple-spotted. Leaves fern-like, gen 2x divided. Flowers white. TOXIC. INVASIVE weed



TALL SOCK-DESTROYER (Torilis arvensis) Naturalized Annual - Carrot Family - (Apr–Jul) - Disturbed places - Plant erect, 12-40". Flower clusters open, > leaf. Fruits 0.1-0.2" long, covered with uncurved bristles. Flowers white or pinkish. INVASIVE weed.

Herbaceous - White

Wild Plants of Sycamore Valley Regional Open Space Preserve - Sorted by Form, Color and Family



SHORT SOCK-DESTROYER (Torilis nodosa) Naturalized Annual - Carrot Family - (Apr-Jun) - Disturbed places - Plant spreading, 4-20" tall. Flowers white to red, clusters dense, < leaf. Fruit ~ 0.1" long, uncurved bristles on outer surface, bumps inside.



WAVYLEAF SOAP PLANT (Chlorogalum pomeridianum var. pomeridianum) Native Perennial - Century Plant Family - (May-Aug) - Common. Open grassland, chaparral, woodland - Plant 1-8' tall, flowers at night. Bulb juices lather in water.



CALIFORNIA MAN-ROOT (Marah fabacea) Native Perennial - Gourd Family - (Feb-Apr) - Streamsides, washes, shrubby open areas - Vine 6-20' long. Leaves moderately lobed. Flowers cream or white, < 0.3" wide. Fruit spiny all over.



GOOSE GRASS (Galium aparine) Native Annual - Madder Family - (Apr-Jun) - Grassy, ± shady places - Stem 12-35" long, weak, brittle. Leaves 0.5-1.6" long, in whorls of 6-8. Flowers white, 4-lobed. Fruits covered w/short, hooked hairs.



SHORTSTEM MORNING-GLORY (Calystegia subacaulis subsp. subacaulis) Native Perennial - Morning-glory Family - (Apr–Jun) - Dry, open scrub or woodland - Stem gen ~0.8" long. Flowers 1.3-2.4" long, white or cream. Bractlets concealing flower bracts.



BINDWEED (Convolvulus arvensis) Naturalized Perennial - Morning-glory Family - (Mar–Oct) - Roadsides, open areas in many pl communities - Trailing. Leaf 0.8-1.2" long, w/round tip, pointed lobes. Flowers white to pink, 0.8-1" long. NOXIOUS.



SHEPHERD'S PURSE (Capsella bursa-pastoris) Naturalized Annual - Mustard Family - (Jan-Oct) - Disturbed areas - Stem 4-20" long. Basal leaves 1.2-2.4" long, dandelion-like. Petals white, 0.08-0.16" long. Fruits 0.16-0.35" long, flat, heart-shaped.



THREADLEAF PEPPERGRASS (Lepidium nitidum) Native Annual - Mustard Family - (Feb-Mar) - Alkaline soils, pastures, dry vernal pools, fields, beaches - Fruit stalk flat. Fruit smooth, 0.14-0.24" long, cupped, winged.

hite

Herbaceous - White



Revision: 3/2/2014

WATER CRESS (Nasturtium officinale) Native Perennial - Mustard Family - (Mar–Nov) - Streams, springs, marshes, lake margins, swamps - Stem 4-43" long. Leaflets 3-9, 0.3-1" wide. Petals white, 0.12-0.18" long. EDIBLE leaves.



FENNEL (Foeniculum vulgare) Naturalized Perennial - Carrot Family - (May–Sep) - Roadsides, disturbed sites - Plants 3-6.5' tall, anise-scented. Stems waxy-blue, canelike. Flowers yellow. Leaf segments thread-like, edible when young. INVASIVE weed.

Wild Plants of Sycamore Valley Regional Open Space Preserve - Sorted by Form, Color and Family



STICKY MOUSE-EAR CHICKWEED (Cerastium glomeratum) Naturalized Annual - Pink Family - (Spring) - Dry hillsides, grassland, chaparral, disturbed areas - Flowers 0.1-02" long, white, sticky-hairy. Flower bract hairs extend beyond tip; bracts herbaceous.



COMMON CHICKWEED (Stellaria media) Naturalized Annual - Pink Family - (Feb-Sep) - Oak woodland, meadows, disturbed areas - Plant 2.8-20" tall. Stem w/line of hairs on 1 side. Sepals 0.12-0.18" long. Petals white, 2-lobed, 0.7-0.9x sepals.



YELLOW JOHNNY-TUCK (Triphysaria eriantha subsp. eriantha) Native Annual - Broom-rape Family - (Mar–May) - Grassland, foothills - Plant 4-14" tall, purple. Leaves 0.4-2" long, 3-7 lobed. Flowers yellow with dark purple beak, 0.4-1" long.



GOLDEN MONKEYFLOWER (Mimulus guttatus) Native Perennial - Lopseed Family - (Mar–Aug) - Common. Wet places, gen terrestrial, occ emergent or floating in mats - Plant 0.8-60". Flower yellow, gen > 0.8" long; mature bracts flattened sideways, inflated in fruit.



BLACK MUSTARD (Brassica nigra) Naturalized Annual - Mustard Family - (Apr–Sep) - Common. Disturbed areas, fields - Plant 1-6' tall. Petals bright yellow, 0.3-0.4" long. Fruit upright, pressed against stem. INVASIVE weed.



SHORTPOD MUSTARD (Hirschfeldia incana) Naturalized Biennial-Perennial - Mustard Family - (Apr-Oct) - Disturbed areas - Stem 16-60". Longest leaves 1.6-4" long, dense-hairy. Petals pale yellow, ~0.2" long. Fruit appressed. Fall-blooming. INVASIVE.



CHARLOCK (Sinapis arvensis) Naturalized Annual - Mustard Family - (Mar-Sep) - Disturbed areas - Stem 8-39", no rosette. Leaves coarse. Petals 0.35-0.47" long, yellow. Fruit 0.8-1.8", ascending, beak 0.24-0.47" long w/fat base. 3-7 valve veins. INVASIVE weed.



HEDGE MUSTARD (Sisymbrium officinale) Naturalized Annual - Mustard Family - (Apr-Sep) - Disturbed areas, fields, pastures - Stem 10-22" tall, thin, wiry. Petals 0.1-0.16" long, pale yellow. Fruit 0.4-0.55" long, awl-shaped, appressed.



CALIFORNIA BURCLOVER (Medicago polymorpha) Naturalized Annual - Pea Family - (Mar–Jul) - Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas - Stem 4-20". Flower yellow, 0.14-0.24" long. Spiny spiral fruits. INVASIVE.



SOURCLOVER (Melilotus indicus) Naturalized Annual-Biennial - Pea Family - (Apr-Oct) - Open, disturbed areas - Stem 4-24" long. Leaflets 3, 0.4-1" long. Flowers yellow, ~0.1" long.



HOP CLOVER (Trifolium campestre) Naturalized Annual - Pea Family - (Apr-May) - Disturbed areas, roadsides - Flower heads 0.3-0.5" wide, bright yellow, brown w/age, quickly reflexing. Flowers with lengthwise ridges.



LITTLE HOP CLOVER (Trifolium dubium) Naturalized Annual - Pea Family - (Spring) -Agricultural, disturbed areas, lawns - Heads 0.16-0.3" wide. Flowers bright yellow, age brown, quickly reflex, smooth.



YELLOW STAR-THISTLE (Centaurea solstitialis) Naturalized Annual - Sunflower Family -(May-Oct) - Invasive, roadsides, disturbed grassland or woodland - Plant 4-39" tall. Leaves woolly, extend down the stem. Flower bract spines 0.4-1" long. NOXIOUS weed.



BRISTLY OX-TONGUE (Helminthotheca echioides) Naturalized Annual-Biennial - Sunflower Family - (All year) - Common. Disturbed areas - Stem 1-6.5' tall, bristly, Leaf 2-8" long, covered with prickly bumps. Flower heads 0.8-1.6" wide. INVASIVE weed.





SMOOTH CAT'S-EAR (Hypochaeris glabra) Naturalized Annual - Sunflower Family -(Mar–Jun) - Common. Disturbed areas, grassland, open woodland - Plant 4-24" tall, smooth. Rays 0.2-0.3" long, barely > head bracts. Only inner fruit beaked. INVASIVE.



ROUGH CAT'S-EAR (Hypochaeris radicata) Naturalized Perennial - Sunflower Family -(Apr–Jul) - Disturbed areas, grassland, open woodland - Plant 4-32" tall, rough-hairy. Rays 0.4-0.6" long, well exceeding head bracts. Fruits all beaked. INVASIVE weed.



PRICKLY LETTUCE (Lactuca serriola) Naturalized Annual - Sunflower Family -(May-Oct) - Abundant. Disturbed places - Stem 1.6-10' tall, prickly-bristly. Leaves deeply-lobed, midvein and edges prickly-bristly. Flowers pale yellow, 0.16-0.2" wide.



COMMON GROUNDSEL (Senecio vulgaris) Naturalized Annual - Sunflower Family -(Feb-Jul) - Common. Disturbed areas - Plant 4-24" tall, not hairy, w/1-10+ stems. Flower head bracts with black-tips. Milky sap.



PRICKLY SOW THISTLE (Sonchus asper subsp. asper) Naturalized Annual - Sunflower Family - (All year) - Common. Slightly moist disturbed sites, along streams - Plant 4-48". Leaf teeth prickly. Basal lobes of upper leaves rounded, curving downward.



COMMON SOW THISTLE (Sonchus oleraceus) Naturalized Annual - Sunflower Family - (All year) - Abundant. Disturbed places - Plant 4-55" tall. Leaf teeth soft to touch. Basal lobes of upper leaves sharply pointed, straight to curving upward.



SNOWBERRY (Symphoricarpos albus var. laevigatus) Native Perennial - Honeysuckle Family - (May-Jul) - Shady woodland, streambanks, N. slopes - Shrub 20-71" tall. Flowers > 8/cluster, pink, 0.16-0.24" long, bell-shaped, swollen on 1 side.



BAY AREA SILVER LUPINE (Lupinus albifrons var. collinus) Native Perennial - Pea Family - (Mar-Jun) - Cliffs, forest openings - Subshrub 8-16" tall, woody only at base, silvery. Flowers 0.35-0.6" long, violet to lavender.



COAST LIVE OAK (Quercus agrifolia var. agrifolia) Native Perennial - Oak Family - (Mar-Apr) - Valleys, slopes, mixed-evergreen forest, woodland - Tree 30-80'. Leaves convex, hair-tuft below in vein axils. Acorns on 1st year twigs, shell glabrous inside.



VALLEY OAK (Quercus lobata) Native Perennial - Oak Family - (Mar–Apr) - Slopes, valleys, savanna - Tree < 115' tall, deciduous. Leaves 2-5" long, not leathery, deeply lobed, lobes without bristles. Acorns 1.2-2" long, slender, cup 0.4-1.2" deep.



BIG-LEAF MAPLE (Acer macrophyllum) Native Perennial - Soapberry Family - (Mar–Jun) -Common. Streambanks, canyons - Tree < 100'. Leaves 5-lobed, 3-6" long, 4-10" wide. Group of 20-90 hanging flowers appear after leaves. Winged seeds.



WESTERN POISON OAK (Toxicodendron diversilobum) Native Perennial - Sumac Family -(Apr–Jun) - Canyons, slopes, chaparral, coastal scrub, oak woodland - Shrub or vine. Leaflets 3, red in fall, mid leaflet stalked. Flowers green. Fruits white. TOXIC.



BLUE ELDERBERRY (Sambucus nigra subsp. caerulea) Native Perennial - Muskroot Family - (Mar–Sep) - Common. Streambanks, open places in forest - Shrub 7-26' tall. Flower cluster flat-topped, 1.6-13" diam, petals spreading. Fruits waxy blue-black.



HIMALAYAN BLACKBERRY (Rubus armeniacus) Naturalized Perennial - Rose Family - (Mar–Jun) - Common. Disturbed areas, roadsides - Shrub w/thorny 5-angled stem. Leaflets 3-5, white-hairy beneath. Blackberry-type fruit. INVASIVE.



CALIFORNIA BUCKEYE (Aesculus californica) Native Perennial - Soapberry Family - (May–Jun) - Dry slopes, canyons, borders of streams - Large shrub or tree 13-39' tall. 5-7 leaflets. Flowers white to pale rose. Large seeds toxic but edible after leaching out saponins.



RED WILLOW (Salix laevigata) Native Perennial Willow Family - (Dec-Jun) - Common. Riverbanks, seepage areas, lakeshores, canyons
- Tree bark fissured. Leaf lanceolate, glaucous below, gen w/stalk glands. Stamens 5. Fruit glabrous.

Notes:



Decimal Inches



English Inches

Notes: