

# East Bay Regional Park District

## Checklist of Wild Plants

### Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in the East Bay Regional Park District. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
<b>Scientific Name</b>	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> ( <a href="http://ucjeps.berkeley.edu/IJM.html">ucjeps.berkeley.edu/IJM.html</a> )
<b>(JM93 if different)</b>	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
<b>Common Name</b>	According to <i>JM2</i> and other references (not standardized)
<b>Family</b>	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
<b>Status</b>	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): <b>Federal/California (Fed./Calif.)   California Native Plant Society (CNPS)   East Bay chapter of the CNPS (EBCNPS)</b>  <b>Fed./Calif.:</b> FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare  <b>CNPS</b> (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California  <b>EBCNPS</b> (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
<b>Inv</b>	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
<b>OL</b>	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
<b>Habitat (Bloom)</b>	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, [wlegard@ebparks.org](mailto:wlegard@ebparks.org)). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

# East Bay Regional Park District

## Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name					<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum aleuticum</i>	Five-finger Fern	Pterid-	-   -   A2	- NP	Shady, moist banks, streamsides, serpentine -	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	-   -   -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Aspidotis californica</i>	California Lace Fern	Pterid-	-   -   A2	- NP	Rock outcrops, crevices -	
<input type="radio"/> <i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	Western Lady Fern	Woods-	-   -   -	- NP	Woodland, along streams, seepage area -	
<input type="radio"/> <i>Azolla filiculoides</i>	Mosquito Fern	Azoll-	-   -   -	- NP	Common. Ponds, slow streams -	
<input type="radio"/> <i>Azolla microphylla</i> (JM93: <i>A. mexicana</i> )	Mexican Mosquito Fern	Azoll-	-   -   -	- NP	Ponds, slow streams -	
<input type="radio"/> <i>Cheilanthes covillei</i>	Coville's Lip Fern	Pterid-	-   -   A2	- NP	Crevices, bases of rocks, sun or shade -	
<input type="radio"/> <i>Cheilanthes intertexta</i>	Coastal Lip Fern	Pterid-	-   -   A2	- NP	Crevices, bases of rocks -	
<input type="radio"/> <i>Cystopteris fragilis</i>	Brittle Fern	Woods-	-   -   C	- NP	Shady, moist rock crevices, meadows, streamsides -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	-   -   -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum arvense</i>	Common Horsetail	Equiset-	-   -   -	- NP	Streambanks, wet meadows, springs, other wet, shaded places -	
<input type="radio"/> <i>Equisetum hyemale</i> subsp. <i>affine</i>	Common Scouring Rush	Equiset-	-   -   -	- NP	Streams, moist, sandy, gravelly areas -	
<input type="radio"/> <i>Equisetum laevigatum</i>	Smooth Scouring Rush	Equiset-	-   -   B	- NP	Moist, sandy or gravelly areas -	
<input type="radio"/> <i>Equisetum</i> sp.	Scouring Rush	Equiset-	-   -   -	- NP	--	
<input type="radio"/> <i>Equisetum telmateia</i> subsp. <i>braunii</i>	Giant Horsetail	Equiset-	-   -   -	- NP	Streambanks, roadside ditches, seepage areas -	
<input type="radio"/> <i>Isoetes howellii</i>	Howell Quillwort	Isoet-	-   -   A2	- NP	Vernal pools, lake margins (Late spring-summer)	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	-   -   -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Pterid-	-   -   C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Pterid-	-   -   -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	-   -   -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Polypodium scolieri</i>	Leather-leaf Fern	Polypodi-	-   -   A1	- NP	Coast, generally in heavy fog-drip or salt-spray zones, on plants (especially Douglas fir, Eucalyptus), dunes, rocky cliffs, bluffs, mtn ridges, granitic or volcanic rocks, mossy logs, or in soil -	
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Dryopterid-	-   -   -	- NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -	
<input type="radio"/> <i>Pteridium aquilinum</i> var. <i>pubescens</i>	Bracken Fern	Dennstaedti-	-   -   -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Selaginella</i>	Sorted Alphabetically by Scientific Name					<i>Selaginella</i>
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="checkbox"/> <i>Selaginella bigelovii</i>	Spike-moss	Selaginell-	-   -   A2	-	NP	Open sites, mineral soil on rock outcrops, amid shrubs -
<input type="checkbox"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	-   -   -	-	NP	Near streams, springs, seeps -

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like	
<i>Agrostis</i>	Sorted Alphabetically by Scientific Name				<i>Agrostis</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis avenacea</i>	Pacific Bent Grass	Po-	-   -   -	L	ZP	Open, often disturbed place (Jun-Jul)
<input type="radio"/> <i>Agrostis capillaris</i>	Colonial Bent Grass	Po-	-   -   -	-	ZP	Roadsides, open, disturbed places (Jul-Sep)
<input type="radio"/> <i>Agrostis exarata</i>	Spike Bent Grass	Po-	-   -   -	-	NP	Common. Moist or disturbed areas, open woodland, conifer forest (Jun-Aug)
<input type="radio"/> <i>Agrostis hallii</i>	Hall's Bent Grass	Po-	-   -   A2	-	NP	Open oak woodland, conifer forest (May-Jul)
<input type="radio"/> <i>Agrostis microphylla</i>	Small-leaf Bent Grass	Po-	-   -   -	-	NA	Thin, rocky soils, cliffs, vernal pools, occasionally on serpentine (May-Jul)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	-   -   -	-	NP	Common. Open meadows, woodland, forest, subalpine (Jun-Aug)
<input type="radio"/> <i>Agrostis sp.</i>	Bent Grass	Po-	-   -   -	-	- -	
<input type="radio"/> <i>Agrostis stolonifera</i>	Creeping Bent Grass	Po-	-   -   -	L	ZP	Ditches, lake margins, marshes (Jun-Sep)
<input type="radio"/> <i>Aira caryophyllea</i>	Silver Hair Grass	Po-	-   -   -	-	ZA	Sandy soils, open or disturbed sites (Apr-Jun)
<input type="radio"/> <i>Alopecurus saccatus</i>	Pacific Foxtail	Po-	-   -   B	-	NA	Vernal pools, moist, open meadows (Mar-May)
<input type="radio"/> <i>Arundo donax</i>	Giant Reed	Po-	-   -   -	H	ZP	Moist places, seeps, ditchbanks (Mar-Sep)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	-   -   -	M	ZA	Disturbed sites (Mar-Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	-   -   -	M	ZA	Disturbed sites (Apr-Jun)
<input type="radio"/> <i>Bolboschoenus maritimus subsp. paludosus</i> (JM93: <i>Scirpus maritimus</i> )	Marine Bulrush	Cyper-	-   -   C	-	NP	Common in CA. Brackish to saline coastal, inland marshes, shores (Summer)
<input type="radio"/> <i>Bolboschoenus robustus</i> (JM93: <i>Scirpus</i> )	Seacoast Bulrush	Cyper-	-   -   -	-	NP	Brackish to saline coastal marshes (Summer)
<input type="radio"/> <i>Brachypodium distachyon</i>	Purple False-brome	Po-	-   -   -	M	ZA	Disturbed areas, dry slopes (Apr-Jul)
<input type="radio"/> <i>Briza maxima</i>	Rattlesnake Grass	Po-	-   -   -	L	ZA	Shaded sites, roadsides, pastures, weedy on coastal dunes (Apr-Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	-   -   -	-	ZA	Shaded or moist, open sites (Apr-Jul)
<input type="radio"/> <i>Bromus arenarius</i>	Australian Chess	Po-	-   -   -	-	ZA	Open, disturbed places (Apr-Jul)
<input type="radio"/> <i>Bromus carinatus var. carinatus</i>	California Brome	Po-	-   -   -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)
<input type="radio"/> <i>Bromus catharticus var. catharticus</i> (JM93: no var.)	Rescue Grass	Po-	-   -   -	-	ZAP	Open, disturbed places (Apr-Nov)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	-   -   -	M	ZA	Open, gen disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	-   -   -	L	ZA	Fields, disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus japonicus</i>	Japanese Brome	Po-	-   -   -	L	ZA	Open, disturbed areas (May-Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	-   -   -	-	NP	Shrubland, conifer forest, shaded streambanks, roadsides (May-Jul)
<input type="radio"/> <i>Bromus madritensis subsp. madritensis</i>	Foxtail Chess	Po-	-   -   -	-	ZA	Disturbed areas, roadsides (Apr-Jan)
<input type="radio"/> <i>Bromus madritensis subsp. rubens</i>	Red Brome	Po-	-   -   -	H	ZA	Disturbed areas, roadsides (Mar-Jun)

Grasses & Grass-like <i>Bromus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Bromus</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Bromus sterilis</i>	Poverty Brome	Po-	-   -   -	-	ZA	Open, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Bromus tectorum</i>	Cheat Grass	Po-	-   -   -	H	ZA	Open, disturbed areas (May–Aug)
<input type="radio"/> <i>Calamagrostis rubescens</i>	Pine Reed Grass	Po-	-   -   B	-	NP	Wooded slopes, montane forest, chaparral, meadows (Jun–Sep)
<input type="radio"/> <i>Carex aquatilis</i> var. <i>dives</i>	Sitka Sedge	Cyper-	-   -   A2	-	NP	Wet meadows, shores, fresh water (Jun–Sep)
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	-   -   B	-	NP	Seasonally wet places (May–Aug)
<input type="radio"/> <i>Carex brevicaulis</i>	Short-stemmed Sedge	Cyper-	-   -   A1	-	NP	Coastal dunes, headlands, rocky or sandy soil (Apr–May)
<input type="radio"/> <i>Carex globosa</i>	Round-fruited Sedge	Cyper-	-   -   A1	-	NP	Well-drained soil of wooded areas, edges (Apr–Jun)
<input type="radio"/> <i>Carex leporina</i> (JM93: <i>C. ovalis</i> )	Tracy Sedge	Cyper-	-   -   A1	-	NP	Seasonally wet soil (May–Aug)
<input type="radio"/> <i>Carex leptopoda</i> (JM93: <i>C. deweyana</i> subsp. <i>leptopoda</i> )	Slender-footed Sedge	Cyper-	-   -   A1	-	NP	Moist soil, wooded areas (May–Aug)
<input type="radio"/> <i>Carex nebrascensis</i>	Nebraska Sedge	Cyper-	-   -   B	-	NP	Meadows, swamps (Jun–Sep)
<input type="radio"/> <i>Carex nudata</i>	Torrent Sedge	Cyper-	-   -   A2	-	NP	Rocky or sandy streambeds below high-water mark (Apr–Jul)
<input type="radio"/> <i>Carex obnupta</i>	Slough Sedge	Cyper-	-   -   A2	-	NP	Moist openings, shores, redwood forest (Jun–Sep)
<input type="radio"/> <i>Carex pellita</i> (JM93: <i>C. lanuginosa</i> )	Woolly Sedge	Cyper-	-   -   A1	-	NP	Generally marshy places, creekbanks (May–Sep)
<input type="radio"/> <i>Carex praegracilis</i>	Freeway Sedge	Cyper-	-   -   C	-	NP	Common. Often alkaline, ± moist places (Apr–Aug)
<input type="radio"/> <i>Carex serratodens</i>	Bifid Sedge	Cyper-	-   -   B	-	NP	Moist places (Apr–Jul)
<input type="radio"/> <i>Carex</i> sp.	Sedge	Cyper-	-   -   -	-	-	-
<input type="radio"/> <i>Carex subbracteata</i>	Small-bracted Sedge	Cyper-	-   -   B	-	NP	At least seasonally moist soil, grassland to open fores (Apr–Jun)
<input type="radio"/> <i>Carex tumulicola</i>	Foothill Sedge	Cyper-	-   -   C	-	NP	Meadows, open woodland (Apr–Jul)
<input type="radio"/> <i>Cortaderia jubata</i>	Hairy Pampas Grass	Po-	-   -   -	H	ZP	Disturbed sites, many habitats, esp coastal (Sep–Feb)
<input type="radio"/> <i>Cortaderia selleana</i>	Smooth Pampas Grass	Po-	-   -   -	H	ZP	Disturbed sites (Sep–Mar)
<input type="radio"/> <i>Crypsis schoenoides</i>	Swamp Prickle Grass	Po-	-   -   -	-	ZA	Wet places (Jun–Oct)
<input type="radio"/> <i>Crypsis vaginiflora</i>	Prickle Grass	Po-	-   -   -	-	ZA	Wet soils, lake margins, vernal pools (Jun–Sep)
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Po-	-   -   -	M	ZP	Disturbed sites (Jun–Aug)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	-   -   -	M	ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Cyperus acuminatus</i>	Tapertip Flatsedge	Cyper-	-   -   -	-	NA	Edges of temporary pools, ponds, streams (Jun–Oct)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	-   -   -	-	NP	Vernal pools, streambanks (May–Nov)
<input type="radio"/> <i>Cyperus erythrorhizos</i>	Redroot Nutsedge	Cyper-	-   -   B	-	NA	Ditches, riverbanks, shores (Jul–Oct)

Grasses & Grass-like <i>Cyperus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Cyperus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Cyperus odoratus</i>	Coarse Nutsedge	Cyper-	-   -   A1	- NP	Wet disturbed soils (Jul–Oct)
<input type="radio"/> <i>Cyperus sp.</i>	Nutsedge	Cyper-	-   -   -	- NAP - -	
<input type="radio"/> <i>Cyperus strigosus</i>	False Nutsedge	Cyper-	-   -   A1	- NP	Moist soils, pond margins, roadsides (Jul–Oct)
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	-   -   -	L ZP	Disturbed, often moist sites (May–Aug)
<input type="radio"/> <i>Danthonia californica</i> (JM93: var. <i>californica</i> )	California Oat Grass	Po-	-   -   C	- NP	Gen moist meadows, open woodland (Apr–Aug)
<input type="radio"/> <i>Deschampsia cespitosa subsp. cespitosa</i>	Tufted Hair Grass	Po-	-   -   C	- NP	Meadows, streambanks, coastal marshes, forest (Jul–Aug)
<input type="radio"/> <i>Deschampsia cespitosa subsp. holciformis</i>	Pacific Hair Grass	Po-	-   -   A2	- NP	Coastal marshes, meadows (May–Jul)
<input type="radio"/> <i>Deschampsia danthonioides</i>	Annual Hair Grass	Po-	-   -   C	- NA	Moist to drying, open sites, meadows, streambanks, vernal pools, occ alkali soil (Mar–Aug)
<input type="radio"/> <i>Deschampsia elongata</i>	Slender Hair Grass	Po-	-   -   C	- NP	Wet sites, meadows, lakeshores, shaded slopes (May–Sep)
<input type="radio"/> <i>Digitaria ischaemum</i>	Smooth Crab Grass	Po-	-   -   -	- ZA	Disturbed areas (Sep–Nov)
<input type="radio"/> <i>Digitaria sanguinalis</i>	Hairy Crab Grass	Po-	-   -   -	- ZA	Disturbed areas (Jun–Sep)
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Po-	-   -   -	- NP	Salt marshes, coastal dunes, moist, alkaline areas (Apr–Sep)
<input type="radio"/> <i>Echinochloa crus-galli</i>	Barnyard Grass	Po-	-   -   -	- ZA	Gen wet, disturbed sites, fields, roadsides (Jun–Oct)
<input type="radio"/> <i>Ehrharta erecta</i>	Panic Veldt Grass	Po-	-   -   -	M ZP	Disturbed, often shady, moist sites (Mar–Jun)
<input type="radio"/> <i>Eleocharis acicularis var. acicularis</i>	Needle Spikerush	Cyper-	-   -   C	- NP	Common. Fresh wet soil to deeply submersed (Late spring–summer)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	-   -   -	- NP	Common. Fresh to brackish wetland (Spring–summer)
<input type="radio"/> <i>Eleocharis parishii</i>	Parish Spikerush	Cyper-	-   -   A2	- NP	Locally common. Fresh streams, vernal pools, foothill and riparian woodland, pine fores ( Late spring–fall)
<input type="radio"/> <i>Eleocharis parvula</i>	Small Spikerush	Cyper-	-   4.3   *A1	- NP	Brackish wet soil, coastal (Late winter–fall)
<input type="radio"/> <i>Eleocharis sp.</i>	Spikerush	Cyper-	-   -   -	- NP - -	
<input type="radio"/> <i>Elymus caput-medusae</i> (JM93: <i>Taeniatherum</i> )	Medusa Head	Po-	-   -   -	H ZA	Disturbed areas (Apr–Jul)
<input type="radio"/> <i>Elymus condensatus</i> (JM93: <i>Leymus</i> )	Giant Wild-rye	Po-	-   -   -	- NP	Dry slopes, open woodland (Jun–Aug)
<input type="radio"/> <i>Elymus glaucus subsp. glaucus</i>	Western Wild-rye	Po-	-   -   -	- NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	-   -   C	- NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus stebbinsii</i>	Parish's Wheat Grass	Po-	-   -   A2	- NP	Dry slopes, chaparral, conifer forest (Jun–Jul)
<input type="radio"/> <i>Elymus trachycaulus subsp. trachycaulus</i>	Slender Wheat Grass	Po-	-   -   B	- NP	Dry to moist, open areas, forest, woodland (Jun–Aug)

Grasses & Grass-like <i>Elymus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Elymus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i> )	Beardless Wild Rye	Po-	-   -   -	- NP	Dry to moist, often saline, meadows ( Jun–Jul)
<input type="radio"/> <i>Eragrostis curvula</i> (JM93: var. <i>curvula</i> )	Weeping Love Grass	Po-	-   -   -	- NP	Roadsides (Aug–Oct)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	-   -   -	M ZP	Disturbed places (May–Jun)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i> )	Brome Fescue	Po-	-   -   -	- ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	-   -   C	- NP	Dry, open forest, moist streambanks, chaparral (May–Jun)
<input type="radio"/> <i>Festuca elmeri</i>	Elmer Fescue	Po-	-   -   A2	- NP	Moist, wooded slopes, under trees in rich soil (May–Jul)
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Po-	-   -   C	- NP	Dry, open or shady places (Jul–Sep)
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i> )	Hairy Fescue	Po-	-   -   -	- NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i> )	Rattail Sixweeks Grass	Po-	-   -   -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca octoflora</i> (JM93: <i>Vulpia</i> , var. <i>hirtella</i> , <i>octoflora</i> )	Slender Six-weeks Fescue	Po-	-   -   A2	- NA	Sandy to rocky soils, open sites (Mar–Jun)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i> )	Rye Grass	Po-	-   -   -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Po-	-   -   B	- NP	Sand dunes, grassland, subalpine forest (May–Jul)
<input type="radio"/> <i>Festuca sp. (annual)</i> (JM93: <i>Vulpia sp.</i> )	Annual Fescue	Po-	-   -   -	- NA	--
<input type="radio"/> <i>Festuca sp. (perennial)</i>	Perennial Fescue	Po-	-   -   -	-	--
<input type="radio"/> <i>Festuca temulenta</i> (JM93: <i>Lolium temulentum</i> )	Long-glumed Rye Grass	Po-	-   -   -	- ZA	Open, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Gastridium phleoides</i> (JM93: <i>G. ventricosum</i> )	Nit Grass	Po-	-   -   -	- ZA	Open, generally dry, disturbed sites (Apr–Nov)
<input type="radio"/> <i>Glyceria occidentalis</i>	Western Manna Grass	Po-	-   -   A2	- NP	Freshwater marshes, ponds and steams (Jun–Aug)
<input type="radio"/> <i>Holcus lanatus</i>	Common Velvet Grass	Po-	-   -   -	M ZP	Moist sites, roadbanks, cult fields, meadows (Jun–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. ?	California / Meadow Barley	Po-	-   -   -	- NP	
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	-   -   -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>californicum</i>	California Barley	Po-	-   -   B	- NP	Meadows, pastures, streambanks (Mar–Jul)
<input type="radio"/> <i>Hordeum depressum</i>	Low Barley	Po-	-   -   B	- NA	Moist sites, vernal pools, generally alkaline soils (Apr–May)
<input type="radio"/> <i>Hordeum jubatum</i> subsp. <i>jubatum</i> (JM93: no subsp.)	Foxtail Barley	Po-	-   -   B	- NAP	Roadsides, disturbed areas, meadows, marshes (May–Jul)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	-   -   -	- ZA	Dry to moist, disturbed sites (Apr–Jun)

Grasses & Grass-like <i>Hordeum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Hordeum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Hordeum murinum subsp. glaucum</i>	Smooth Barley	Po-	-   -   -	- ZA	Moist, generally disturbed sites (Mar–Jul)
<input type="radio"/> <i>Hordeum murinum subsp. leporinum</i>	Hare Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Hordeum murinum subsp. murinum</i>	Wall Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites (Feb–May)
<input type="radio"/> <i>Isolepis cernua (JM93: Scirpus cernuus)</i>	Low Bulrush	Cyper-	-   -   B	- NA	Sandy, sometimes brackish sea shores, bluffs, sand dunes, creeks, marshes (Late spring–winter)
<input type="radio"/> <i>Juncus balticus subsp. ater (JM93: no subsp.)</i>	Baltic Rush	Junc-	-   -   -	- NP	Moist to ± dry sites (Jul–Nov)
<input type="radio"/> <i>Juncus bufonius var. ?</i>	Toad Rush	Junc-	-   -   -	- NA	--
<input type="radio"/> <i>Juncus bufonius var. bufonius</i>	Toad Rush	Junc-	-   -   -	- NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus bufonius var. congestus</i>	Clustered Toad Rush	Junc-	-   -   -	- NA	Drying pools, shores, disturbed ground, gen saline (Apr–Aug)
<input type="radio"/> <i>Juncus dubius</i>	Mariposa Rush	Junc-	-   -   A1	- NP	Wet places (Jul–Sep)
<input type="radio"/> <i>Juncus effusus subsp. pacificus (JM93: var.)</i>	Pacific Rush	Junc-	-   -   -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus lescurii (JM93: J. lesueurii)</i>	San Francisco Rush	Junc-	-   -   A2	- NP	Salt or freshwater marshes, shores of creeks and lakes (Aug–Sep)
<input type="radio"/> <i>Juncus mexicanus</i>	Mexican Rush	Junc-	-   -   B	- NP	Common. Coast to montane meadows (Aug–Nov)
<input type="radio"/> <i>Juncus occidentalis</i>	Western Rush	Junc-	-   -   -	- NP	Moist gen sunny areas (May–Sep)
<input type="radio"/> <i>Juncus oxymiris</i>	Pointed Rush	Junc-	-   -   A2	- NP	Uncommon. Swales, montane meadows, damp, sunny ground (Jul–Aug)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	-   -   -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus phaeocephalus var. ?</i>	Brownhead Rush	Junc-	-   -   -	- NP	--
<input type="radio"/> <i>Juncus phaeocephalus var. paniculatus</i>	Panicled Brownhead Rush	Junc-	-   -   C	- NP	Wet places, coastal, inland (Jun–Sep)
<input type="radio"/> <i>Juncus phaeocephalus var. phaeocephalus</i>	Brownheaded Rush	Junc-	-   -   B	- NP	Coastal meadows, dune hollows, marsh edges (Jun–Aug)
<input type="radio"/> <i>Juncus sp.</i>	Rush	Junc-	-   -   -	-	--
<input type="radio"/> <i>Juncus tenuis</i>	Slender Rush	Junc-	-   -   -	- NP	Uncommon. Damp places (Jun–Sep)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	-   -   -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria gerardii (JM93: var. hindsii)</i>	Annual June Grass	Po-	-   -   -	- ZA	Open, disturbed sites (Apr–Jul)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	-   -   C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Lamarckia aurea</i>	Goldentop	Po-	-   -   -	- ZA	Open ground, moist seeps, rocky hillsides, sandy soil (Feb–May)
<input type="radio"/> <i>Leersia oryzoides</i>	Rice Cut Grass	Po-	-   -   A2	- NP	Marshes, streams, ponds (Aug–Oct)
<input type="radio"/> <i>Leptochloa fusca subsp. fascicularis (JM93: L. fascicularis)</i>	Bearded Sprangletop	Po-	-   -   A2	- NA	Marshes, wetlands, often wet disturbed areas, often associated with agriculture (Jun–Oct)
<input type="radio"/> <i>Luzula comosa var. comosa (JM93: no var.)</i>	Common Woodrush	Junc-	-   -   -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)



Grasses & Grass-like <i>Melica</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Melica</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Melica bulbosa</i>	Oniongrass	Po-	-   -   A1?	- NP	Dry rocky slopes, conifer forest (Jul–Aug)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	-   -   C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	-   -   -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	-   -   -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Panicum capillare</i>	Witch Grass	Po-	-   -   B	- NA	Open places, fields, roadsides (Feb–Dec)
<input type="radio"/> <i>Panicum sp.</i>	Panic Grass	Po-	-   -   -	- -	-
<input type="radio"/> <i>Parapholis incurva</i>	Sickle Grass	Po-	-   -   -	- ZA	Disturbed, well drained soils of salt marshes, generally above highest tide level (Apr–Jun)
<input type="radio"/> <i>Paspalum dilatatum</i>	Dallis Grass	Po-	-   -   -	- ZP	Disturbed areas (May–Nov)
<input type="radio"/> <i>Paspalum distichum</i>	Knot Grass	Po-	-   -   -	- ZP	Edges of lakes, ponds, rice fields, wet roadside ditches (Jun–Oct)
<input type="radio"/> <i>Paspalum sp.</i>	Dallis / Knot Grass	Po-	-   -   -	- ZP	-
<input type="radio"/> <i>Pennisetum clandestinum</i>	Kikuyu Grass	Po-	-   -   -	L ZP	Disturbed areas (All year)
<input type="radio"/> <i>Phalaris aquatica</i>	Harding Grass	Po-	-   -   -	M ZP	Disturbed areas, roadsides (Apr–Aug)
<input type="radio"/> <i>Phalaris arundinacea</i>	Reed Canary Grass	Po-	-   -   A2	- NP	Moist habitats, woodland (May–Sep)
<input type="radio"/> <i>Phalaris canariensis</i>	Canary Grass	Po-	-   -   -	- ZA	Disturbed areas (Apr–Jul)
<input type="radio"/> <i>Phalaris minor</i>	Little-seeded Canary Grass	Po-	-   -   -	- ZA	Disturbed areas, cultivated fields (Apr–Jul)
<input type="radio"/> <i>Phalaris paradoxa</i>	Hood Canary Grass	Po-	-   -   -	- ZA	Disturbed areas, cultivated field (May–Aug)
<input type="radio"/> <i>Phragmites australis</i>	Common Reed	Po-	-   -   B	N NP	Pond & lake margins, sloughs, marshes (Jul–Nov)
<input type="radio"/> <i>Pleuropogon californicus var. californicus</i> (JM93: no var.)	California Semaphore Grass	Po-	-   -   B	- NA	Wet places, vernal pools, wet grassland (Mar–Jun)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	-   -   -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa bulbosa subsp. vivipara</i> (JM93: no subsp.)	Bulbous Blue Grass	Po-	-   -   -	- ZP	Disturbed places (Mar–Jul)
<input type="radio"/> <i>Poa howellii</i>	Howell Blue Grass	Po-	-   -   B	- NA	Rocky banks, shaded slopes of woodland, chaparral, disturbed places (Apr–Jun)
<input type="radio"/> <i>Poa pratensis subsp. pratensis</i>	Kentucky Blue Grass	Po-	-   -   -	- ZP	Common. Disturbed habitats, including saline or alkaline soils (Mar–Jul)
<input type="radio"/> <i>Poa secunda subsp. secunda</i>	One-sided Blue Grass	Po-	-   -   -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Po-	-   -   -	- ZP	Common. Streambanks (May–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	-   -   -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Polypogon viridis</i> (JM93: <i>Agrostis</i> )	Water Beard Grass	Po-	-   -   -	- ZP	Common. Disturbed areas, wet areas, ponds, streambanks (May–Jun)
<input type="radio"/> <i>Puccinellia nutkaensis</i>	Alaska Alkali Grass	Po-	-   -   A1	- NP	Marshes, wet sites (Jun)

Grasses & Grass-like <i>Puccinellia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Puccinellia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Puccinellia nuttalliana</i>	Nuttall Alkali Grass	Po-	-   -   A1	- NP	Saline meadows (Jun–Sep)
<input type="radio"/> <i>Puccinellia simplex</i>	California Alkali Grass	Po-	-   -   A2	- NA	Saline flats, mineral springs (Mar–May)
<input type="radio"/> <i>Schoenoplectus acutus</i> var. <i>occidentalis</i> (JM93: <i>Scirpus</i> )	Common Tule	Cyper-	-   -   -	- NP	Common. Marshes, shores, fens, shallow lakes, often emergent (Summer)
<input type="radio"/> <i>Schoenoplectus americanus</i> (JM93: <i>Scirpus</i> )	Olney's Three-square Bulrush	Cyper-	-   -   -	- NP	Mineral-rich or brackish marshes, shores, fens, springs (Summer)
<input type="radio"/> <i>Schoenoplectus californicus</i> (JM93: <i>Scirpus</i> )	Southern Bulrush	Cyper-	-   -   -	- NP	Common. Brackish to fresh marshes, shores (Spring–summer)
<input type="radio"/> <i>Scirpus microcarpus</i>	Smallfruit Bulrush	Cyper-	-   -   -	- NP	Marshes, wet meadows, streambanks, pond margins, sometimes weedy (Jun–Jul)
<input type="radio"/> <i>Scirpus</i> , <i>Isolepis</i> , or <i>Bolboschoenus</i> sp. (JM93: <i>Scirpus</i> sp.)	Bulrush	Cyper-	-   -   -	- NAP	
<input type="radio"/> <i>Scribneria bolanderi</i>	Scribner Grass	Po-	-   -   A2	- NP	Dry, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Sorghum bicolor</i>	Sorghum	Po-	-   -   -	- ZA	Disturbed areas, roadsides (All year)
<input type="radio"/> <i>Sorghum halepense</i>	Johnson Grass	Po-	-   -   -	- ZP	Disturbed areas, ditch banks, roadsides (May–Aug)
<input type="radio"/> <i>Sorghum</i> sp.	Johnson Grass / Sorghum	Po-	-   -   -	- ZAP - -	
<input type="radio"/> <i>Sparganium eurycarpum</i> var. <i>eurycarpum</i> (JM93: subsp. <i>eurycarpum</i> )	Broad-fruited Bur-reed	Typh-	-   -   A2	- NP	Marshes, lakes, ponds, along streams (Jun–Aug)
<input type="radio"/> <i>Sparganium eurycarpum</i> var. <i>greenii</i> (JM93: <i>S. erectum</i> subsp. <i>stoloniferum</i> )	Erect Bur-reed	Typh-	-   -   A1	- NP	Generally coastal (Jun–Aug)
<input type="radio"/> <i>Spartina alterniflora</i>	Smooth Saltwater Cordgrass	Po-	-   -   -	H ZP	Coastal salt marshes (Jul–Sep)
<input type="radio"/> <i>Spartina densiflora</i>	Dense-flowered Cordgrass	Po-	-   -   -	H ZP	Coastal salt marshes (May–Aug)
<input type="radio"/> <i>Spartina foliosa</i>	California Cordgrass	Po-	-   -   B	- NP	Salt marshes, mudflats, shores (Jun–Nov)
<input type="radio"/> <i>Sporobolus airoides</i>	Alkali Sacaton	Po-	-   -   A2	- NP	Seasonally moist, alkaline areas (Apr–Oct)
<input type="radio"/> <i>Stipa cernua</i> (JM93: <i>Nassella</i> )	Nodding Needle Grass	Po-	-   -   C	- NP	Grassland, chaparral, juniper woodland (Feb–Jul)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i> )	Foothill Needle Grass	Po-	-   -   C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa miliacea</i> var. <i>miliacea</i> (JM93: <i>Piptatherum miliaceum</i> )	Smilo Grass	Po-	-   -   -	L ZP	Salt marshes, streambanks, chaparral, open woodland, disturbed (Mar–Sep)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i> )	Purple Needle Grass	Po-	-   -   C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Triglochin maritima</i>	Common Arrow-grass	Juncagin-	-   -   C	- NP	Coastal salt marshes, interior saline, brackish, alkaline marshes (Apr–Aug)
<input type="radio"/> <i>Triglochin scilloides</i> (JM93: <i>Lilaea</i> )	Flowering-quillwort	Juncagin-	-   -   C	- NA	Vernal pools, streams, ponds, lake margins (Mar–Oct)
<input type="radio"/> <i>Triglochin striata</i>	Three-ribbed Arrow-grass	Juncagin-	-   -   A2	- NP	Uncommon. Brackish to freshwater coastal marshes, springs (May–Sep)

Grasses & Grass-like <i>Trisetum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Trisetum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Trisetum canescens</i>	Tall False Oat	Po-	-   -   B	- NP	Open to shaded sites, meadows, chaparral, conifer forest (May–Aug)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	-   -   -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Typh-	-   -   -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	-   -   -	- NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)
<input type="radio"/> <i>Typha sp.</i>	Cattail	Typh-	-   -   -	- NP	- -
<input type="radio"/> <i>Zostera marina</i>	Eel-grass	Zoster-	-   -   -	- NP	Common. Shallow water, bays, estuaries (Mar–Jun)

Herbaceous <i>Acaena</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Acaena</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acaena pinnatifida</i> var. <i>californica</i>	California Acaena	Ros-	-   -   C	- NP	Coastal grassland, open, rocky slopes (Mar–May)	
<input type="radio"/> <i>Acanthomintha lanceolata</i>	Santa Clara Thornmint	Lami-	-   4.2   *A2	- NP	Arid, rocky slopes, openings in scrub or woodland, gen on serpentine (Mar–Jun)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	-   -   -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Aster-	-   -   -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)	
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i> )	Spanish Clover	Fab-	-   -   -	- NA		
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i> )	Colchita	Fab-	-   -   -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i> )	Deerweed	Fab-	-   -   -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon</i> or <i>Hosackia</i> sp. (JM93: <i>Lotus</i> sp.)	Lotus	Fab-	-   -   -	- -		
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i> )	Small-flower Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Acmispon strigosus</i> (JM93: <i>Lotus</i> )	Hairy Lotus	Fab-	-   -   A2	- NP	Coastal scrub, chaparral, foothills, deserts, roadsides, other disturbed area (Mar–Jun)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i> )	California Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Actaea rubra</i>	Baneberry	Ranuncul-	-   -   B	- NP	Deep soils, moist, open to shaded sites, mixed-evergreen or conifer forests (May–Sep)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	-   -   -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Aster-	-   -   -	- NA	Many open habitats (May–Jun)	
<input type="radio"/> <i>Agoseris retrorsa</i>	Spearleaf Native Dandelion	Aster-	-   -   B	- NP	Scrub, oak woodland, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Agoseris</i> sp.	Agoseris	Aster-	-   -   -	- NP	- -	
<input type="radio"/> <i>Alisma triviale</i> (JM93: <i>A. plantago-aquatica</i> )	Northern Water Plantain	Alismat-	-   -   B	- NP	Ponds (Spring-fall)	
<input type="radio"/> <i>Allium amplexans</i>	Narrowleaf Onion	Alli-	-   -   A2	- NP	Clays including serpentine, open or wooded places (Apr–Jul)	
<input type="radio"/> <i>Allium bolanderi</i> var. <i>bolanderi</i>	Bolander's Onion	Alli-	-   -   A1	- NP	Uncommon. Rocky clays including serpentine (May–Aug)	
<input type="radio"/> <i>Allium falcifolium</i>	Sickle Leaf Onion	Alli-	-   -   A2	- NP	Common. Heavy clays including serpentine (Apr–Jun)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	-   -   -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Allium triquetrum</i>	Three Cornered Leek	Alli-	-   -   -	- ZP	Locally common. Shady ± disturbed places (Mar–Apr)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Allium</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Allium</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Alli-	-   -   C	- NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)	
<input type="radio"/> <i>Allophyllum divaricatum</i>	Stragglng Gilia	Polemoni-	-   -   A2	- NA	Sandy areas, chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Allophyllum gilioides subsp. gilioides</i>	Purple False Gilia	Polemoni-	-   -   B	- NA	Open, sandy, gen damp or grassy areas (Apr–Jun)	
<input type="radio"/> <i>Amaranthus albus</i>	Tumbleweed	Amaranth-	-   -   -	- ZA	Disturbed areas, roadsides, riverbanks, sandy places, agricultural fields (Jun–Oct)	
<input type="radio"/> <i>Amaranthus blitoides</i>	Procumbent Pigweed	Amaranth-	-   -   C	- NA	Disturbed areas, roadsides, agricultural fields, sandy soil (Jul–Nov)	
<input type="radio"/> <i>Amaranthus californicus</i>	California Amaranth	Amaranth-	-   -   A2	- NA	Seasonally moist flats, lake margins, disturbed areas (Jul–Oct)	
<input type="radio"/> <i>Amaranthus deflexus</i>	Low Amaranth	Amaranth-	-   -   -	- ZA	Railroad right-of-ways, disturbed areas (May–Nov)	
<input type="radio"/> <i>Amaranthus powellii</i>	Powell's Amaranth	Amaranth-	-   -   A2	- NA	Agricultural fields, disturbed areas ( Jun–Oct)	
<input type="radio"/> <i>Amaranthus retroflexus</i>	Redroot Pigweed	Amaranth-	-   -   -	- ZA	Common. Wet fields, roadside ditches, waste places, agricultural fields (Jun–Nov)	
<input type="radio"/> <i>Amaranthus sp.</i>	Amaranth	Amaranth-	-   -   -	- -	- -	
<input type="radio"/> <i>Ambrosia chamissonis</i>	Beach Bur-sage	Aster-	-   -   A2	- NP	Beaches, dunes (Jul–Oct)	
<input type="radio"/> <i>Ambrosia psilostachya</i>	Western Ragweed	Aster-	-   -   -	- NP	Common. Roadsides, dry fields (Jun–Nov)	
<input type="radio"/> <i>Amsinckia grandiflora</i>	Large-flowered Fiddleneck	Boragin-	FE/CE   1B.1   *A1	- NA	Grassy slopes (Mar–May)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i> )	Common Fiddleneck	Boragin-	-   -   -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Amsinckia lunaris</i>	Bent-flowered Fiddleneck	Boragin-	-   1B.2   *A2	- NA	Gravelly slopes, grassland, openings in woodland (Mar–Jun)	
<input type="radio"/> <i>Amsinckia lycopsoides</i>	Bugloss-flowered Fiddleneck	Boragin-	-   -   B	- NA	Common. Open, grassland, foothill woodland, generally disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Amsinckia menziesii</i> (JM93: var. <i>menziesii</i> )	Small-flowered Fiddleneck	Boragin-	-   -   -	- NA	Shade-tolerant, open, disturbed areas at forest/woodland edges (May–Jul)	
<input type="radio"/> <i>Amsinckia sp.</i>	Fiddleneck	Boragin-	-   -   -	- NA	- -	
<input type="radio"/> <i>Amsinckia tessellata</i> var. ?	Devil's Lettuce	Boragin-	-   -   -	- NA	- -	
<input type="radio"/> <i>Amsinckia tessellata</i> var. <i>tessellata</i>	Desert Fiddleneck	Boragin-	-   -   B	- NA	Rocky or sandy soils (Feb–Jun)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	-   -   -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Anaphalis margaritacea</i>	Pearly Everlasting	Aster-	-   -   -	- NP	Woodland, disturbed places (Jul–Oct)	
<input type="radio"/> <i>Ancistrocarphus filagineus</i>	Woolly Fishhooks	Aster-	-   -   B	- NA	Bare or grassy, often serpentine or clay, drainages, road beds, burns, vernal moist sites (Mar–Jun)	
<input type="radio"/> <i>Anemopsis californica</i>	Yerba Mansa	Saurur-	-   -   B	- NP	Common. Saline or alkaline soil, wet or moist areas, seeps, springs (Mar–Sep)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Angelica</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Angelica</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	-   -   -	- NP	Generally wooded areas (Jun–Aug)	
<input type="radio"/> <i>Anisocarpus madioides</i> (JM93: <i>Madia</i> )	Woodland Tarweed	Aster-	-   -   A2	- NP	Moist forest, woodland (Apr–Sep)	
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	-   -   -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	-   -   -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Antirrhinum multiflorum</i>	Mayflower Snapdragon	Plantagin-	-   -   A1	- NA	Rocky or disturbed areas, burns (Apr–Aug)	
<input type="radio"/> <i>Antirrhinum vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i> (JM93: <i>A. vexillocalyculatum</i> subsp. <i>vexillo-calyculatum</i> )	Wiry Snapdragon	Plantagin-	-   -   B	- NA	Disturbed areas (Jun–Aug)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	-   -   -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Apiastrum angustifolium</i>	Wild Celery	Api-	-   -   B	- NA	Chaparral, coastal scrub (Mar–Apr)	
<input type="radio"/> <i>Apium graveolens</i>	Celery	Api-	-   -   -	- ZA	Wet places (May–Jul)	
<input type="radio"/> <i>Apocynum androsaemifolium</i>	Bitter Dogbane	Apocyn-	-   -   B	- NP	Open slopes, rocky places, with conifers, chaparral (May–Oct)	
<input type="radio"/> <i>Apocynum cannabinum</i>	Indian Hemp	Apocyn-	-   -   A2	- NP	Moist places, near streams, springs, or as weed in orchards (Apr–Oct)	
<input type="radio"/> <i>Aquilegia eximia</i>	Serpentine Columbine	Ranuncul-	-   -   A2	- NP	Generally serpentine seeps, occasionally moist ravines, mixed-evergreen or conifer forests (May–Oct)	
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	-   -   C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)	
<input type="radio"/> <i>Aralia californica</i>	Elk Clover	Arali-	-   -   B	- NP	Moist shade, canyons, stream sides (Jun–Aug)	
<input type="radio"/> <i>Aristolochia californica</i>	Dutchman's Pipe	Aristolochi-	-   -   C	- NP	Streamsides, forest, chaparral (Jan–Apr)	
<input type="radio"/> <i>Arnica discoidea</i>	Rayless Arnica	Aster-	-   -   B	- NP	Chaparral, foothill woodland (May–Jul)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	-   -   -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Artemisia dracunculus</i>	Wild Tarragon	Aster-	-   -   A1	- NP	Common. Many habitats, especially disturbed sites (Aug–Oct)	
<input type="radio"/> <i>Arthrocnemum subterminale</i> (JM93: <i>Salicornia</i> )	Glasswort Pickleweed	Chenopodi-	-   -   A2	- NP	Salt marshes, alkaline flats (Apr–Sep)	
<input type="radio"/> <i>Asarum caudatum</i>	Wild Ginger	Aristolochi-	-   -   A2	- NP	Moist places in forest (Mar–Aug)	
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	-   -   B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)	
<input type="radio"/> <i>Asclepias cordifolia</i>	Purple Milkweed	Apocyn-	-   -   A1	- NP	Rocky slopes, talus, woodland, chaparral, lava flows (May–Jul)	
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	-   -   -	- NP	Dry ground, valleys, foothills (May–Oct)	

Herbaceous <i>Asclepias</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Asclepias</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Asclepias speciosa</i>	Showy Milkweed	Apocyn-	-   -   A2	- NP	Many habitats incl fields, roadsides (May–Sep)
<input type="radio"/> <i>Asparagus officinalis</i> subsp. <i>officinalis</i>	Garden Asparagus	Lili-	-   -   -	- ZP	Disturbed places, roadsides, fields (Mar–Sep)
<input type="radio"/> <i>Aster</i> sp.	Aster	Aster-	-   -   -	- NP	- -
<input type="radio"/> <i>Astragalus asymmetricus</i>	San Joaquin Milkvetch	Fab-	-   -   B	- NP	Grassy areas, open woodland, disturbed sites (Apr–Jul)
<input type="radio"/> <i>Astragalus didymocarpus</i> var. <i>didymocarpus</i>	Two-seed Milkvetch	Fab-	-   -   A1	- NA	Grassy areas (Feb–May)
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvetch	Fab-	-   -   -	- NA	Open, grassy areas, scrub (Mar–Jul)
<input type="radio"/> <i>Astragalus oxyphysus</i>	Stanislaus Milkvetch	Fab-	-   -   A1?	- NP	Arid grassland, scrub (Mar–Jun)
<input type="radio"/> <i>Astragalus</i> sp.	Milkvetch	Fab-	-   -   -	- NAP	- -
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	-   -   -	- NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)
<input type="radio"/> <i>Atriplex cordulata</i>	Heartscale	Chenopodi-	-   1B.2   *A1	- NA	Saline or alkaline soils (Jun–Jul)
<input type="radio"/> <i>Atriplex depressa</i>	Brittlescale	Chenopodi-	-   1B.2   *A2	- NA	Alkaline or clay soils (Jun–Oct)
<input type="radio"/> <i>Atriplex joaquiniana</i> (JM93: <i>A. joaquiniana</i> )	San Joaquin Spearscale	Chenopodi-	-   1B.2   *A2	- NA	Alkaline soils (Apr–Sep)
<input type="radio"/> <i>Atriplex leucophylla</i>	Beach Saltbush	Chenopodi-	-   -   A2	- NP	Sandy soils, dunes (Apr–Oct)
<input type="radio"/> <i>Atriplex prostrata</i> (JM93: <i>A. triangularis</i> )	Fat-hen	Chenopodi-	-   -   -	- NA	Wet places, marshes (Apr–Oct)
<input type="radio"/> <i>Atriplex rosea</i>	Tumbling Orach	Chenopodi-	-   -   -	- ZA	Common. Open, disturbed places, fields (Jul–Oct)
<input type="radio"/> <i>Atriplex semibaccata</i>	Australian Saltbush	Chenopodi-	-   -   -	M ZP	Disturbed areas, scrub, woodland (Apr–Dec)
<input type="radio"/> <i>Atriplex serenana</i> var. <i>serenana</i>	Bract Scale	Chenopodi-	-   -   A1	- NA	Alkaline flats, coastal bluffs (Apr–Oct)
<input type="radio"/> <i>Atriplex</i> sp.	Saltbush / Orach	Chenopodi-	-   -   -	-	- -
<input type="radio"/> <i>Baccharis glutinosa</i> (JM93: <i>B. douglasii</i> )	Marsh Baccharis	Aster-	-   -   -	- NP	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)
<input type="radio"/> <i>Balsamorhiza deltoidea</i>	Balsamroot	Aster-	-   -   A2	- NP	Common. Grassy slopes, open forests, shrubby areas (Apr–Jul)
<input type="radio"/> <i>Balsamorhiza macrolepis</i> (JM93: var. <i>macrolepis</i> )	Bigscale Balsamroot	Aster-	-   1B.2   *A1	- NP	Open grassy or rocky slopes, valleys (Mar–Jul)
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	-   -   -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)
<input type="radio"/> <i>Bassia hyssopifolia</i>	Hyssop-leaf Bassia	Chenopodi-	-   -   -	L ZA	Disturbed sites, fields, roadsides (May–Nov)
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	-   -   -	L ZA	Disturbed grassland. (Apr–Jun)
<input type="radio"/> <i>Bellis perennis</i>	English Daisy	Aster-	-   -   -	- ZP	Damp, grassy areas (Dec–Sep)
<input type="radio"/> <i>Berula erecta</i>	Cutleaf Water-parsnip	Api-	-   -   B	- NP	Marshy areas, streams (Jul–Oct)
<input type="radio"/> <i>Bidens frondosa</i>	Sticktight	Aster-	-   -   B	- NA	Uncommon. Damp soil, esp disturbed sites (Jun–Oct)
<input type="radio"/> <i>Bidens laevis</i>	Bur-marigold	Aster-	-   -   A1	- NAP	Freshwater wetlands (Aug–Nov)

Herbaceous <i>Blennosperma</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Blennosperma</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Blennosperma nanum</i> var. <i>nanum</i>	Glue-seed	Aster-	-   -   C	- NA	Open, grassy areas, often margins of seeps or vernal pools (Jan–May)
<input type="radio"/> <i>Blepharizonia laxa</i> (JM93: <i>B. plumosa</i> subsp. <i>viscida</i> )	Stickyleaf Big Tarweed	Aster-	-   -   A2	- NA	Openings in woodland, chaparral, grassland (Jul–Nov)
<input type="radio"/> <i>Blepharizonia plumosa</i> (JM93: subsp.)	Big Tarplant	Aster-	-   1B.1   *A2	- NA	Dry slopes in grassland (Jul–Nov)
<input type="radio"/> <i>Boechera breweri</i> subsp. <i>breweri</i> (JM93: <i>Arabis breweri</i> var. <i>breweri</i> )	Brewer's Rockcress	Brassic-	-   -   A1	- NP	Rocky outcrops, ledges, talus Mar–Jul
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	-   -   -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	-   -   -	M ZA	Common. Disturbed areas, fields (Apr–Sep)
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	-   -   -	L ZA	Disturbed areas (Jan–May)
<input type="radio"/> <i>Brodiaea elegans</i> subsp. <i>elegans</i>	Harvest Brodiaea	Themid-	-   -   -	- NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)
<input type="radio"/> <i>Brodiaea terrestris</i> subsp. <i>terrestris</i>	Dwarf Brodiaea	Themid-	-   -   A2	- NP	Coastal prairie, foothill woodland (Apr–Jul)
<input type="radio"/> <i>Cakile maritima</i>	Horned Sea Rocket	Brassic-	-   -   -	L ZA	Beach dunes (May–Nov)
<input type="radio"/> <i>Calandrinia breweri</i>	Brewer Red Maids	Monti-	-   4.2   *A2	- NA	Sandy to loamy soil, disturbed sites, burns (Feb–May)
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	-   -   -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)
<input type="radio"/> <i>Callitriche marginata</i>	California Water-starwort	Callitrich-	-   -   -	- NA	Becoming stranded (often in vernal pools) (Mar–Jun)
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	-   -   -	- NP	Common. Shady to open woodland, scrub (Apr–Jun)
<input type="radio"/> <i>Calochortus argillosus</i>	Clay Mariposa Lily	Lili-	-   -   -	- NP	Hard clay from volcanic or metamorphic rocks (Apr–Jun)
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	-   -   C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)
<input type="radio"/> <i>Calochortus pulchellus</i>	Mount Diablo Fairy-lantern	Lili-	-   1B.2   *A2	- NP	Wooded slopes, rarely chaparral, gen n aspect (Apr–Jun)
<input type="radio"/> <i>Calochortus</i> sp.	Mariposa Lily	Lili-	-   -   -	- NP	- -
<input type="radio"/> <i>Calochortus splendens</i>	Splendid Lilac Mariposa Lily	Lili-	-   -   A1	- NP	Dry, often granitic soils, grassland, chaparral, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	-   -   -	- NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calochortus umbellatus</i>	Oakland Star-tulip	Lili-	-   4.2   *A2	- NP	Open chaparral or woodland, gen on serpentine (Mar–May)
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	-   -   B	- NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)



Herbaceous <i>Calycadenia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Calycadenia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Calycadenia multiglandulosa</i>	Sticky Rosinweed	Aster-	-   -   A1	- NA	Common. Gen dry, open valleys, hillsides, rocky ridges (May–Oct)	
<input type="radio"/> <i>Calycadenia truncata</i>	Rosinweed	Aster-	-   -   A2	- NA	Common. Dry, open hillsides, rocky ridges, talus, grassland, openings in foothill woodland, scrub (May–Oct)	
<input type="radio"/> <i>Calystegia malacophylla subsp. pedicellata</i>	Woolly Morning-glory	Convolvul-	-   -   A2	- NP	Dry slopes, chaparral (Apr–Jul)	
<input type="radio"/> <i>Calystegia occidentalis subsp. occidentalis</i>	Western Morning-glory	Convolvul-	-   -   -	- NP	Dry slopes, chaparral, pine woodland (May–Jul)	
<input type="radio"/> <i>Calystegia purpurata subsp. purpurata</i>	Climbing Morning-glory	Convolvul-	-   -   -	- NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia sepium subsp. limnophila</i>	Hedge Bindweed	Convolvul-	-   -   A2	- NP	Marshes, riverbanks (May–Jul)	
<input type="radio"/> <i>Calystegia sp.</i>	Morning-glory	Convolvul-	-   -   -	- NP	--	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	-   -   -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Camissonia contorta</i>	Contorted Suncup	Onagr-	-   -   B	- NA	Sandy soil, slopes, flats, often disturbed, grassland, chaparral, pinyon/juniper woodland (May–Ju)	
<input type="radio"/> <i>Camissoniopsis intermedia</i> (JM93: <i>Camissonia</i> )	Intermediate Suncup	Onagr-	-   -   A2	- NA	Shrubby slopes, especially burns (Mar–May)	
<input type="radio"/> <i>Camissoniopsis micrantha</i> (JM93: <i>Camissonia</i> )	Small-flower Suncup	Onagr-	-   -   B	- NA	Coastal dunes, beaches, sandy fields, washes (Mar–May)	
<input type="radio"/> <i>Campanula exigua</i>	Chaparral / Rock Harebell	Campanul-	-   1B.2   *A2	- NA	Talus slopes, generally serpentine soil (May–Jun)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	-   -   -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	-   -   -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	-   -   -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	-   -   -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Carduus tenuiflorus</i>	Slender Flower Thistle	Aster-	-   -   -	L ZAB	Roadsides, pastures, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	-   -   -	- NP	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Orobanch-	-   -   -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja densiflora subsp. densiflora</i>	Common Owl's-clover	Orobanch-	-   -   -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	-   -   -	- NA	Open fields, grassland (Mar–May)	
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	-   -   -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)	
<input type="radio"/> <i>Castilleja rubicundula subsp. lithospermoides</i>	Yellow Cream Sacs	Orobanch-	-   -   B	- NA	Open grassland (Apr–Jun)	
<input type="radio"/> <i>Castilleja subinclusa subsp. franciscana</i>	Franciscan Paintbrush	Orobanch-	-   -   A2	- NP	Coastal scrub (Mar–Jul)	
<input type="radio"/> <i>Castilleja wightii</i>	Seaside / Coastal Paintbrush	Orobanch-	-   -   C	- NP	Coastal scrub (Mar–Aug)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Caulanthus</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Caulanthus</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Caulanthus flavescens</i> (JM93: <i>Guillenia</i> )	Yellow-flowered Jewelflower	Brassic-	-   -   A2	- NA	Dry, exposed slopes, generally on serpentine, open hillsides (Mar–May)	
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i> )	California Mustard	Brassic-	-   -   -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun	
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	-   -   -	M ZAB	Pastures, disturbed places (Apr–Nov)	
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	-   -   -	M ZA	Disturbed fields, open woodland (Apr–Jul)	
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	-   -   -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)	
<input type="radio"/> <i>Centranthus ruber</i>	Red Valerian	Valerian-	-   -   -	- ZP	Disturbed places, rock or wall crevices, roadsides (Apr–Jul)	
<input type="radio"/> <i>Centromadia fitchii</i> (JM93: <i>Hemizonia</i> )	Fitch Spikeweed	Aster-	-   -   -	- NP	Grassland, ± alkaline flats, vernal pools, woodland, disturbed sites, serpentine (May–Nov)	
<input type="radio"/> <i>Centromadia parryi</i> subsp. <i>congdonii</i> (JM93: <i>Hemizonia</i> )	Congdon's Tarplant	Aster-	-   1B.2   *A2	- NA	Terraces, swales, floodplains, grassland, disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Centromadia parryi</i> subsp. <i>parryi</i> (JM93: <i>Hemizonia</i> )	Parry Spikeweed	Aster-	-   -   -	- NA	Grassland, coastal salt marshes, alkaline springs, seeps (Jun–Oct)	
<input type="radio"/> <i>Centromadia pungens</i> subsp. <i>pungens</i> (JM93: <i>Hemizonia</i> , subsp. <i>maritima</i> , <i>pungens</i> )	Common Spikeweed	Aster-	-   -   -	- NA	Grassland, saltbush scrub, disturbed sites (Apr–Nov)	
<input type="radio"/> <i>Cerastium fontanum</i> subsp. <i>vulgare</i>	Large Mouse-ear Chickweed	Caryophyll-	-   -   -	- ZA	Disturbed areas, grassy slopes, damp woodland, marshy ground (Spring)	
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	-   -   -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)	
<input type="radio"/> <i>Ceratophyllum demersum</i>	Hornwort	Ceratophyll-	-   -   -	- NA	Common. Ditches, lakes, ponds, pools, slow watercourses; water 0.1–4 m deep, fresh to ± brackish, medium to high nutrient levels. (Jun–Aug)	
<input type="radio"/> <i>Chaenactis glabriuscula</i> var. <i>megacephala</i>	Big-headed Yellow Pincushion	Aster-	-   -   A1	- NA	Dry, often sandy slopes, openings in chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Chamaesyce maculata</i>	Spotted Spurge	Euphorbi-	-   -   -	- ZA	Disturbed places (Apr–Oct)	
<input type="radio"/> <i>Chamaesyce ocellata</i> subsp. <i>ocellata</i>	Valley Spurge	Euphorbi-	-   -   B	- NA	Dry, sandy places (May–Sep)	
<input type="radio"/> <i>Chamaesyce serpyllifolia</i> subsp. <i>serpyllifolia</i>	Thyme-leaved Spurge	Euphorbi-	-   -   B	- NA	Common. Dry habitats (Aug–Oct)	
<input type="radio"/> <i>Chenopodium album</i>	Lamb's Quarters	Chenopodi-	-   -   -	- ZA	Common. Disturbed areas, fields (Jun–Oct)	
<input type="radio"/> <i>Chenopodium californicum</i>	California Goosefoot	Chenopodi-	-   -   -	- NP	Generally open sites, sandy to clay soils (Mar–Sep)	
<input type="radio"/> <i>Chenopodium desiccatum</i>	Aridland Goosefoot	Chenopodi-	-   -   -	- NA	Uncommon. Open places, scrub, conifer forest (Jul–Sep)	

Herbaceous <i>Chenopodium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Chenopodium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Chenopodium macrospermum</i> (JM93: var. <i>halophilum</i> )	Coast Goosefoot	Chenopodi-	-   -   -	- ZA	Wet places, marshes (Jul–Oct)
<input type="radio"/> <i>Chenopodium murale</i>	Wall Goosefoot	Chenopodi-	-   -   -	- ZA	Common. Disturbed areas, fields (All year)
<input type="radio"/> <i>Chenopodium rubrum</i> var. <i>rubrum</i> (JM93: no var.)	Red Goosefoot	Chenopodi-	-   -   A1	- NA	Open, saline places, drying mudflats Aug–Oct
<input type="radio"/> <i>Chenopodium</i> sp.	Goosefoot	Chenopodi-	-   -   -	- -	
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	-   -   -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Chloropyron molle</i> subsp. <i>molle</i> (JM93: <i>Cordylanthus mollis</i> subsp. <i>mollis</i> )	Soft Salty Bird's-beak	Orobanch-	FE/CR   1B.2   *A2	- NA	Coastal salt marshes (Jul–Nov)
<input type="radio"/> <i>Chorizanthe membranacea</i>	Pink Spineflower	Polygon-	-   -   A2	- NA	Common. Sand, gravel or rocks (Apr–Jul)
<input type="radio"/> <i>Chorizanthe polygonoides</i> var. <i>polygonoides</i>	Knotweed Spineflower	Polygon-	-   -   A1	- NA	Sand or gravel (Apr–Jun)
<input type="radio"/> <i>Cicendia quadrangularis</i>	Timwort	Gentian-	-   -   B	- NA	Open places (Mar–May)
<input type="radio"/> <i>Cichorium intybus</i>	Chicory	Aster-	-   -   -	- ZP	Common. Roadsides, disturbed places (Apr–Oct)
<input type="radio"/> <i>Cicuta douglasii</i>	Western Water-hemlock	Api-	-   -   B	- NP	Wet places, gen aquatic (Jun–Sep)
<input type="radio"/> <i>Cicuta maculata</i> var. <i>bolanderi</i>	Bolander Water Hemlock	Api-	-   2.1   *A1	- NP	Coastal wetlands ( Jul–Sep)
<input type="radio"/> <i>Cirsium brevistylum</i>	Indian Thistle	Aster-	-   -   B	- NA	Moist places (Mar–Aug)
<input type="radio"/> <i>Cirsium cymosum</i> var. <i>cymosum</i> (JM93: no var.)	Peregrine Thistle	Aster-	-   -   B	- NP	Scrubland, woodland, open forest, meadows, occ serpentine (Jun–Jul)
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Aster-	-   -   -	- NB	Disturbed areas, grassland, woodland (May–Jul)
<input type="radio"/> <i>Cirsium quercetorum</i>	Brownie Thistle	Aster-	-   -   A2	- NP	Open places, grassland, woodland (Apr–Aug)
<input type="radio"/> <i>Cirsium</i> sp.	Thistle	Aster-	-   -   -	- -	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	-   -   -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia affinis</i>	Small Clarkia	Onagr-	-   -   C	- NA	Openings in woodland, chaparral (May–Jun)
<input type="radio"/> <i>Clarkia biloba</i> subsp. <i>biloba</i>	Mule-ear Clarkia	Onagr-	-   -   A2	- NA	Foothill woodland, serpentine or not (May–Aug)
<input type="radio"/> <i>Clarkia concinna</i> subsp. ?	Red Ribbons	Onagr-	-   -   -	- NA	- -
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>automixa</i>	Santa Clara Red Ribbons	Onagr-	-   4.3   *A1	- NA	Woodland (Apr–Jun)
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>concinna</i>	Red Ribbons	Onagr-	-   -   C	- NA	Mixed-evergreen forest, woodland, coastal scrub (Apr–Jul)
<input type="radio"/> <i>Clarkia franciscana</i>	Presidio Clarkia	Onagr-	FE/CE   1B.1   *A1	- NA	Serpentine soil (May–Jun)
<input type="radio"/> <i>Clarkia gracilis</i> subsp. <i>gracilis</i>	Herald-of-summer	Onagr-	-   -   B	- NA	Common. Openings in woodland, forest (Apr–Jul)
<input type="radio"/> <i>Clarkia modesta</i>	Waltham Creek Fairy Fan	Onagr-	-   -   A2	- NA	Shady places in woodland (Apr–May)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. ?	Winecup Clarkia	Onagr-	-   -   -	- NA	- -

Herbaceous <i>Clarkia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Clarkia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>purpurea</i>	Purple Winecup Clarkia	Onagr-	-   -   A2	-	NA	Uncommon. Grassland (Apr–Jun)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	-   -   -	-	NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>viminea</i>	Large Clarkia	Onagr-	-   -   A2	-	NA	Open, grassy or shrubby places (May–Jul)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	-   -   -	-	NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	-   -   -	-	NA	- -
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	-   -   -	-	NA	Common. Woodland (Apr–Sep)
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>exigua</i>	Common Pale Claytonia	Monti-	-   -   C	-	NA	Dry or moist, disturbed bare clay to sandy soils, often serpentine (Apr–Jul)
<input type="radio"/> <i>Claytonia gypsophiloides</i>	Santa Lucia Claytonia	Monti-	-   -   A2	-	NA	Moist, bare, often stony sites, in sun or shade, often serpentine (Mar–May)
<input type="radio"/> <i>Claytonia parviflora</i> subsp. ?	Miner's Lettuce	Monti-	-   -   -	-	NA	- -
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i> )	Small-flower Claytonia	Monti-	-   -   -	-	NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	-   -   -	-	NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	-   -   -	-	NA	Vernally moist, often shady or disturbed sites (Jan–May)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i> )	Yerba Buena	Lami-	-   -   -	-	NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Clintonia andrewsiana</i>	Red-bead Lily	Lili-	-   -   A1	-	NP	Shaded, damp redwood forest (May–Jul)
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. ?	White Chinese Houses	Plantagin-	-   -   -	-	NA	- -
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. <i>bartsiiifolia</i>	White Chinese Houses	Plantagin-	-   -   A1	-	NA	Open, sandy places (Mar–Jun)
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. <i>stricta</i> (JM93: var. <i> davidsonii</i> )	Small White Chinese Houses	Plantagin-	-   -   A1	-	NA	Chaparral, open oak woodland, dry mixed woodland (May–Jul)
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	-   -   -	-	NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)
<input type="radio"/> <i>Collinsia parviflora</i>	Blue-eyed Mary	Plantagin-	-   -   A2	-	NA	Common. Moist, ± shady places, montane (Mar–Jul)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. ?	Few-flowered Collinsia	Plantagin-	-   -   -	-	NA	- -
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	-   -   C	-	NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>sparsiflora</i>	Few-flowered Collinsia	Plantagin-	-   -   C	-	NA	Grassy, sometimes disturbed or rocky places, chaparral, oak woodland, dry mixed woodland (Mar–May)
<input type="radio"/> <i>Collomia heterophylla</i>	Variable-leaf Collomia	Polemoni-	-   -   A2	-	NA	Sandy to gravelly, open areas (Apr–Jun)

Herbaceous <i>Conium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Conium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	-   -   -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	-   -   -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)
<input type="radio"/> <i>Corallorhiza maculata</i> var. <i>maculata</i> (JM93: no var.)	Spotted Coralroot	Orchid-	-   -   A2	- NP	Shaded mixed-evergreen or conifer forest, in decomposing lf litter (May–Aug)
<input type="radio"/> <i>Corallorhiza striata</i>	Striped Coralroot	Orchid-	-   -   A1	- NP	Open to shaded mixed-evergreen or conifer forest, in decomposing leaf litter (Feb–Jul)
<input type="radio"/> <i>Cordylanthus pilosus</i> subsp. <i>pilosus</i>	Hairy Bird's Beak	Orobanch-	-   -   B	- NA	Open foothill woodland, chaparral, on serpentine (Jul–Sep)
<input type="radio"/> <i>Cordylanthus rigidus</i> subsp. <i>rigidus</i>	Stiff Bird's Beak	Orobanch-	-   -   A1	- NA	Open foothill woodland, chaparral margins, conifer forest (Jul–Sep)
<input type="radio"/> <i>Corethrogyne filaginifolia</i> (JM93: <i>Lessingia</i> , var. <i>filaginifolia</i> )	Common California-aster	Aster-	-   -   C	- NP	Coastal scrub, chaparral, grassland, foothill woodland, forest (Jul–Nov)
<input type="radio"/> <i>Cotula australis</i>	Australian Brass Buttons	Aster-	-   -   -	- ZP	Common. Disturbed areas (Jan–May)
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	-   -   -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)
<input type="radio"/> <i>Crassula aquatica</i>	Water Pygmy-weed	Crassul-	-   -   C	- NA	Salt marshes, vernal pools, margins of lakes, ponds (Mar–Jun(Aug))
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	-   -   -	- NA	Open areas (Feb–May)
<input type="radio"/> <i>Crassula tillaea</i>	European Pygmy Weed	Crassul-	-   -   -	- ZA	Open, gravelly sites (Feb–May)
<input type="radio"/> <i>Crepis vesicaria</i> subsp. <i>taraxacifolia</i>	Dandelion-leaf Hawksbeard	Aster-	-   -   -	- ZAB	Sandy clearings, hillsides, disturbed places (Feb–Oct)
<input type="radio"/> <i>Cressa truxillensis</i>	Alkali Weed	Convolvul-	-   -   -	- NP	Saline and alkaline substrates (May–Oct)
<input type="radio"/> <i>Crocidium multicaule</i>	Spring Gold	Aster-	-   -   A1	- NA	Sandy soils, grassland, open woodland (Feb–Jun)
<input type="radio"/> <i>Croton californicus</i>	Croton	Euphorbi-	-   -   A2	- NP	Sandy soils, dunes, washes (Apr–Jul)
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i> )	Turkey-mullein	Euphorbi-	-   -   -	- NA	Dry, open, often disturbed areas (May–Oct)
<input type="radio"/> <i>Cryptantha flaccida</i>	Weak-stemmed Cryptantha	Boragin-	-   -   -	- NA	Common; semi-barren, gravelly, loose soils, rocky sites, washes, slopes, ridges (Apr–Jul)
<input type="radio"/> <i>Cryptantha intermedia</i> var. <i>intermedia</i> (JM93: no var.)	Intermediate Cryptantha	Boragin-	-   -   A1	- NA	Sandy to gravelly flats, slopes, generally granitic or serpentine-based, grassland, chaparral, foothill woodland, occasionally open conifer forest (May–Jul)
<input type="radio"/> <i>Cryptantha micromeres</i>	Minute-flowered Cryptantha	Boragin-	-   -   A1	- NA	Open sites, disturbed, coarse soils, chaparral, woodland, burns (Mar–Jul)
<input type="radio"/> <i>Cryptantha muricata</i> var. <i>muricata</i> (JM93: no var.)	Prickly Cryptantha	Boragin-	-   -   A2	- NA	Coastal scrub, chaparral, foothill woodland, open conifer forest (Mar–Jul)

Herbaceous <i>Cryptantha</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Cryptantha</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Cryptantha rattanii</i> (JM93: <i>C. decipiens</i> )	Deceptive Cryptantha	Boragin-	-   -   A2	- NA	Rocky, gravelly slopes, grassland, coastal scrub, chaparral, foothill woodland (Apr–Jul)	
<input type="radio"/> <i>Cryptantha</i> sp.	Cryptantha	Boragin-	-   -   -	- NA	-	
<input type="radio"/> <i>Cryptantha torreyana</i> var. <i>pumila</i> (JM93: no var.)	Dwarf Cryptantha	Boragin-	-   -   A2	- NA	Generally chaparral, foothill woodland (Apr–Jun)	
<input type="radio"/> <i>Cuscuta californica</i> var. <i>californica</i>	California Dodder	Convolvul-	-   -   A2	- NA	On herbs, shrubs, roadsides, chaparral, grassland, yellow-pine forest (May–Sep)	
<input type="radio"/> <i>Cuscuta pacifica</i> var. <i>pacifica</i> (JM93: <i>C. salina</i> var. <i>major</i> )	Goldenthread	Convolvul-	-   -   B	- NA	Gen on Salicornia, Jaumea in coastal salt marshes, tidal flats (Jul–Oct)	
<input type="radio"/> <i>Cuscuta subinclusa</i>	Canyon Dodder	Convolvul-	-   -   B	- NP	Generally on herbs, shrubs, in forests near streams, river canyon bottoms, salt marshes (Mar–Oct(Dec?))	
<input type="radio"/> <i>Cymbalaria muralis</i>	Kenilworth Ivy	Plantagin-	-   -   -	- ZP	Rock walls, shady, disturbed areas (May–Sep)	
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	-   -   -	M ZP	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	-   -   -	- NP	Chaparral, woodland (Feb–May)	
<input type="radio"/> <i>Datisca glomerata</i>	Durango Root	Datisc-	-   -   A2	- NP	Dry streambeds or washes (May–Jul)	
<input type="radio"/> <i>Datura wrightii</i>	Wright Jimson Weed	Solan-	-   -   -	- NAP	Sandy or gravelly open areas (Apr–Oct)	
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	-   -   -	- NA	Rocky or sandy places (Apr–Jun)	
<input type="radio"/> <i>Deinandra corymbosa</i> (JM93: <i>Hemizonia</i> , subsp. <i>corymbosa</i> )	Coast Spikeweed	Aster-	-   -   A2	- NA	Grassland, openings in scrub or woodland, dunes, disturbed sites (Mar–Nov)	
<input type="radio"/> <i>Deinandra lobbii</i> (JM93: <i>Hemizonia</i> )	Lobb Tarweed	Aster-	-   -   B	- NA	Grassland, open woodland, sagebrush scrub, disturbed areas (May–Dec)	
<input type="radio"/> <i>Delairea odorata</i> (JM93: <i>Senecio mikanioides</i> )	German Ivy	Aster-	-   -   -	H ZP	Shady, ± disturbed places, riparian woodland, coastal scrub (Nov–Mar)	
<input type="radio"/> <i>Delphinium californicum</i> subsp. <i>californicum</i>	California Larkspur	Ranuncul-	-   -   A2	- NP	Generally slopes in dense chaparral, w side of coast ranges (Apr–Jun)	
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Ranuncul-	-   -   C	- NP	Open coastal grassland, chaparral (Mar–May)	
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>hesperium</i>	Western Larkspur	Ranuncul-	-   -   B	- NP	Oak woodland, w slope coast ranges (Mar–Jun)	
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>pallescens</i>	Pale-flowered Western Larkspur	Ranuncul-	-   -   C	- NP	Oak woodland, e slope coast ranges (Mar–May)	
<input type="radio"/> <i>Delphinium nudicaule</i>	Red Or Orange Larkspur	Ranuncul-	-   -   C	- NP	Moist talus, wooded, rocky slopes (Mar–Jun)	
<input type="radio"/> <i>Delphinium patens</i> subsp. <i>patens</i>	Spreading Larkspur	Ranuncul-	-   -   -	- NP	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Delphinium</i> sp.	Larkspur	Ranuncul-	-   -   -	- NP	-	
<input type="radio"/> <i>Delphinium variegatum</i> subsp. <i>variegatum</i>	Royal Larkspur	Ranuncul-	-   -   C	- NP	Grassland, open oak woodland (Mar–May)	
<input type="radio"/> <i>Descurainia sophia</i>	Feather-leaf Tansy Mustard	Brassic-	-   -   -	L ZA	Common. Disturbed areas, fields, canyon bottoms, desert (May–Aug)	

Herbaceous <i>Dicentra</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Dicentra</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Dicentra formosa</i>	Pacific Bleeding Heart	Papaver-	-   -   A2	- NP	Damp, shaded areas (Mar–Jul)
<input type="radio"/> <i>Dichelostemma capitatum subsp. capitatum</i>	Blue Dicks	Themid-	-   -   -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	-   -   -	- NP	Open woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Dichelostemma sp.</i>	Blue Dicks/ookow	Themid-	-   -   -	- NP	- -
<input type="radio"/> <i>Digitalis purpurea</i>	Foxglove	Plantagin-	-   -   -	L ZB	Acid soil in open woodland, disturbed areas (May–Jul)
<input type="radio"/> <i>Dipsacus sativus</i>	Fuller's Teasel	Dipsac-	-   -   -	M ZB	Disturbed areas, fields, vacant lots, pastures (May–Jul)
<input type="radio"/> <i>Dittrichia graveolens</i>	Stinkwort	Aster-	-   -   -	M ZA	Disturbed areas (Sep–Nov)
<input type="radio"/> <i>Dodecatheon clevelandii subsp. ?</i>	Shooting Star	Primul-	-   -   -	- NP	- -
<input type="radio"/> <i>Dodecatheon clevelandii subsp. patulum</i>	Padre Shooting Star	Primul-	-   -   C	- NP	Moist places, often on serpentine or in ± alkaline sites (Mar–May)
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	-   -   -	- NP	Gen in shady sites (Mar–Jul)
<input type="radio"/> <i>Downingia ornata subsp. eximia</i>	Horned Downingia	Campanul-	-   -   A1x	- NA	Vernal pools, roadside ditches (Apr–May)
<input type="radio"/> <i>Downingia pulchella</i>	Flatface Downingia	Campanul-	-   -   B	- NA	Vernal pools, roadside ditches (Apr–Jun)
<input type="radio"/> <i>Downingia sp.</i>	Downingia	Campanul-	-   -   -	- NA	- -
<input type="radio"/> <i>Draba cuneifolia (JM93: var. integrifolia)</i>	Wedge-leaf Whitlow Grass	Brassic-	-   -   A1	- NA	Open or disturbed places (May)
<input type="radio"/> <i>Draba verna</i>	Spring Whitlow Grass	Brassic-	-   -   B	- NA	Open or disturbed areas (Feb–May)
<input type="radio"/> <i>Dryocallis glandulosa var. glandulosa (JM93: Potentilla glandulosa ssp. glandulosa)</i>	Sticky Cinquefoil	Ros-	-   -   -	- NP	Gen ± shady or moist areas (May–Jul)
<input type="radio"/> <i>Duchesnea indica var. indica (JM93: no var.)</i>	Mock-strawberry	Ros-	-   -   -	- ZP	Disturbed areas (May–Aug)
<input type="radio"/> <i>Dudleya cymosa subsp. cymosa</i>	Spreading Dudleya	Crassul-	-   -   A1	- NP	Rocky outcrops, talus slopes, less often shaded canyon slopes (May–Jul)
<input type="radio"/> <i>Dudleya cymosa subsp. paniculata</i>	Common / Hot Rock Dudleya	Crassul-	-   -   B	- NP	Uncommon. Rocky outcrops, canyons (May–Jun)
<input type="radio"/> <i>Dudleya farinosa</i>	Powdery Dudleya	Crassul-	-   -   A1	- NP	Common: Coastal soils and cliffs (Jun–Aug)
<input type="radio"/> <i>Dysphania ambrosioides (JM93: Chenopodium)</i>	Mexican Tea	Chenopodi-	-   -   -	- ZAP	Disturbed places (Jul–Sep)
<input type="radio"/> <i>Dysphania multifida (JM93: Chenopodium)</i>	Cutleaf Goosefoot	Chenopodi-	-   -   -	- ZA	Disturbed areas (Jul–Oct)
<input type="radio"/> <i>Echinodorus berteroi</i>	Upright Burhead	Alismat-	-   -   A2	- NA	Ponds, ditches (Mid-summer–fall)
<input type="radio"/> <i>Eclipta prostrata</i>	False Daisy	Aster-	-   -   A1	- NA	Damp places (All year)
<input type="radio"/> <i>Ehrendorferia chrysantha (JM93: Dicentra)</i>	Golden Eardrops	Papaver-	-   -   A2	- NP	Dry slopes, burns, disturbed areas (Apr–Sep)
<input type="radio"/> <i>Eichhornia crassipes</i>	Water Hyacinth	Pontederi-	-   -   -	H ZP	Locally abundant. Ponds, sloughs, waterways (Jun–Oct)
<input type="radio"/> <i>Elatine californica</i>	California Waterwort	Elatin-	-   -   A1	- NA	Pools, ponds, rice fields, streambanks (Mar–Aug)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Emmenanthe</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Emmenanthe</i></b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
○ <i>Emmenanthe penduliflora</i> var. <i>penduliflora</i>	Whispering Bells	Boragin-	-   -   B	- NP	Chaparral to creosote-bush scrub, rocky, sandy, decomposed granite, serpentine soils (Apr–Jul)	
○ <i>Enemion occidentale</i> (JM93: <i>Isopyrum</i> )	Western Rue-anemone	Ranuncul-	-   -   A1	- NP	Shaded slopes, chaparral, oak woodland, conifer forest (Mar–May)	
○ <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	-   -   -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)	
○ <i>Epilobium campestre</i> (JM93: <i>E. pygmaeum</i> )	Smooth Willowherb	Onagr-	-   -   B	- NA	Vernal pools, clay mud flats (May–Sep)	
○ <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	-   -   -	- NP	Dry slopes, ridges (Jun–Dec)	
○ <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Onagr-	-   -   -	- NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)	
○ <i>Epilobium ciliatum</i> subsp. <i>watsonii</i>	Watson Willowherb	Onagr-	-   -   B	- NP	Moist coastal bluffs, streamsides, ± disturbed sites (May–Oct)	
○ <i>Epilobium densiflorum</i>	Dense-flower Willowherb	Onagr-	-   -   C	- NA	Streambanks, outwashes, seasonal moist flats (May–Oct)	
○ <i>Epilobium minutum</i>	Chaparral Willowherb	Onagr-	-   -   -	- NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)	
○ <i>Epilobium</i> sp.	Willowherb	Onagr-	-   -   -	- NA	-	
○ <i>Epilobium torreyi</i>	Narrowleaf Willowherb	Onagr-	-   -   A2	- NA	Seasonally moist streambanks, seeps, roadside ditches (May–Aug)	
○ <i>Epipactis gigantea</i>	Stream / Brook Orchid	Orchid-	-   -   A1	- NP	Seeps, wet meadows, streambank (Mar–Oct)	
○ <i>Epipactis helleborine</i>	Broad-leaved Helleborine Orchid	Orchid-	-   -   -	- ZP	Generally dry slopes, roadcuts, mixed-conifer forest (Apr–Dec)	
○ <i>Eremalche parryi</i> subsp. <i>parryi</i>	Parry's Mallow	Malv-	-   -   A1x	- NA	Grassland, scrub, foothill woodland (Mar–May)	
○ <i>Erethogon boothii</i> subsp. <i>decorticans</i> (JM93: <i>Camissonia</i> )	Shredding Evening-primrose	Onagr-	-   -   A1	- NA	Open, generally steep and rocky, especially shale slopes (Mar–Jun)	
○ <i>Erigeron bonariensis</i> (JM93: <i>Conyza</i> )	Flax-leaved Horseweed	Aster-	-   -   -	- ZA	Disturbed sites (All year)	
○ <i>Erigeron canadensis</i> (JM93: <i>Conyza</i> )	Horseweed	Aster-	-   -   -	- ZA	Disturbed places (All year)	
○ <i>Erigeron foliosus</i> var. <i>foliosus</i>	Leafy Fleabane	Aster-	-   -   -	- NP	Open, rocky grassland, chaparral, forest (May–Aug)	
○ <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Aster-	-   -   -	- NP	Grassy dunes, chaparral, oak woodland (May–Oct)	
○ <i>Erigeron petrophilus</i> var. <i>petrophilus</i>	Rock-loving Fleabane	Aster-	-   -   A2	- NP	Rocky foothills to montane forest, sometimes on serpentine (May–Sep)	
○ <i>Erigeron philadelphicus</i> var. <i>philadelphicus</i> (JM93: no var.)	Philadelphia Fleabane	Aster-	-   -   C	- NBP	Streamsides, other moist habitats (May–Jun)	
○ <i>Erigeron</i> sp. (JM93: <i>Conyza</i> )	Fleabane Daisy	Aster-	-   -   -	-	-	
○ <i>Erigeron sumatrensis</i> (JM93: <i>Conyza bilbaoana</i> )	Tropical Horseweed	Aster-	-   -   -	- ZA	Disturbed sites (All year)	



Herbaceous <i>Eriogonum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Eriogonum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Eriogonum angulosum</i>	Angle-stem Wild Buckwheat	Polygon-	-   -   A2	- NA	Common. Clay (All year)
<input type="radio"/> <i>Eriogonum gracile</i> var. <i>gracile</i>	Slender Woolly Wild Buckwheat	Polygon-	-   -   B	- NA	Common. Sand (All year)
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>caninum</i>	Tiburon Buckwheat	Polygon-	-   1B.2   *A1	- NA	Serpentine (May–Oct)
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>luteolum</i>	Golden-carpet Wild Buckwheat	Polygon-	-   -   A2	- NA	Common. Serpentine (Jul–Nov)
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	-   -   -	- NP	--
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	-   -   A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Eriogonum roseum</i>	Wand Wild Buckwheat	Polygon-	-   -   B	- NA	Common. Sand or gravel (May–Nov)
<input type="radio"/> <i>Eriogonum</i> sp.	Wild Buckwheat	Polygon-	-   -   -	- NAP	--
<input type="radio"/> <i>Eriogonum umbellatum</i> var. <i>smallianum</i> (JM93: var. <i>bahiiiforme</i> )	Small's Sulphur Flower	Polygon-	-   -   *A2	- NP	Serpentine (Jul–Sep)
<input type="radio"/> <i>Eriogonum wrightii</i> var. <i>trachygonum</i>	Rough-node Bastard-sage	Polygon-	-   -   A2	- NP	Common. Gravel (Jul–Oct)
<input type="radio"/> <i>Eriophyllum lanatum</i> var. <i>achilleoides</i> (JM93: var. <i>achillaeoides</i> )	Common Woolly Sunflower	Aster-	-   -   C	- NP	Dry, often rocky sites, chaparral, forest (May–Jul)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium brachycarpum</i>	Southern European Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	-   -   -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	-   -   -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Eryngium articulatum</i>	Beethistle	Api-	-   -   A1	- NP	Seasonally wet areas on lake and stream margins, marshes (Jun–Aug)
<input type="radio"/> <i>Eryngium jepsonii</i> (JM93: <i>E. aristulatum</i> var. <i>aristulatum</i> )	Button Celery	Api-	-   -   -	- NP	Moist clay soil, vernal pools, lake shores, drying lakes, wet depressions (Apr–Aug)
<input type="radio"/> <i>Eryngium</i> sp.	Eryngo	Api-	-   -   -	- NP	--
<input type="radio"/> <i>Erysimum capitatum</i> var. <i>capitatum</i> (JM93: var. <i>capitatum</i> , var. <i>angustatum</i> )	Western Wallflower	Brassic-	-   -   C	- NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	-   -   -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Eucrypta chrysanthemifolia</i> var. <i>chrysanthemifolia</i>	Common Eucrypta	Boragin-	-   -   B	- NA	Roadsides, burns, coastal bluffs, ravines (Mar–Jun)
<input type="radio"/> <i>Euphorbia crenulata</i>	Chinese Caps	Euphorbi-	-   -   -	- NA	Common. Dry places (Mar–Aug)
<input type="radio"/> <i>Euphorbia lathyris</i>	Caper Spurge	Euphorbi-	-   -   -	- ZAB	Disturbed areas (All year)
<input type="radio"/> <i>Euphorbia oblongata</i>	European Spurge	Euphorbi-	-   -   -	L ZP	Disturbed areas (Jun–Aug)
<input type="radio"/> <i>Euphorbia peplus</i>	Petty Spurge	Euphorbi-	-   -   -	- ZA	Common. Disturbed areas (Feb–Aug)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	-   -   -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i> )	Broadleaf Aster	Aster-	-   -   -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)

Herbaceous <i>Euthamia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Euthamia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Euthamia occidentalis</i>	Western Goldenrod	Aster-	-   -   -	- NP	Marshes, streambanks, meadows (Jul–Nov)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	-   -   -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Fragaria vesca</i>	Wood Strawberry	Ros-	-   -   -	- NP	Gen partial shade in forest (Jan–Jul)
<input type="radio"/> <i>Frankenia salina</i>	Alkali Heath	Frankeni-	-   -   -	- NP	Salt marshes, alkali flats (Apr–Sep)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: <i>var. affinis</i> )	Checker Lily	Lili-	-   -   -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Fritillaria agrestis</i>	Stinkbells	Lili-	-   4.2   *A2	- NP	Clay (gen serpentine) banks, depressions (Mar–Jun)
<input type="radio"/> <i>Fritillaria liliacea</i>	Fragrant Fritillary	Lili-	-   1B.2   *A1	- NP	Heavy soil, open hills, fields near coast (Feb–Apr)
<input type="radio"/> <i>Fumaria officinalis</i>	Fumitory	Papaver-	-   -   -	- ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Galium andrewsii</i> subsp. <i>gatense</i>	Phlox-leaf Serpentine Bedstraw	Rubi-	-   4.2   *A2	- NP	Dry, rocky places in serpentine soil, chaparral or open oak/pine woodland (Apr–Jun)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	-   -   -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	-   -   -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	-   -   -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)
<input type="radio"/> <i>Galium parisiense</i>	Wall Bedstraw	Rubi-	-   -   -	- ZA	Warm, dry, generally rocky soil to moist areas, coastal-sage scrub, chaparral, grassy hillsides with oaks, roadsides (Apr–Aug)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	-   -   -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Galium</i> sp.	Bedstraw	Rubi-	-   -   -	- -	-
<input type="radio"/> <i>Galium trifidum</i> subsp. <i>columbianum</i> (JM93: <i>var. pacificum</i> )	Trifid Bedstraw	Rubi-	-   -   A2	- NP	Wet places, yellow-pine forest (Jul–Aug)
<input type="radio"/> <i>Galium triflorum</i>	Sweet-scent Bedstraw	Rubi-	-   -   B	- NP	Damp, shady forest (May–Jul)
<input type="radio"/> <i>Gamochaeta ustulata</i> (JM93: <i>Gnaphalium purpureum</i> )	Purple Cudweed	Aster-	-   -   -	- NA	Dunes, bluffs, fields, disturbed sites (Apr–Jul)
<input type="radio"/> <i>Geranium carolinianum</i>	Carolina Geranium	Gerani-	-   -   -	- NA	Open to shaded sites, grassland, scrub, forest (Feb–Aug)
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	-   -   -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	-   -   -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Geranium robertianum</i>	Herb Robert	Gerani-	-   -   -	- ZAB	Open to shaded sites (Apr–Sep)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>achilleifolia</i>	California Gilia	Polemoni-	-   -   B	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>multicaulis</i>	Many-stem California Gilia	Polemoni-	-   -   -	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Feb–Jun)
<input type="radio"/> <i>Gilia capitata</i> subsp. ?	Globe Gilia	Polemoni-	-   -   -	- NA	-

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Gilia</i>	Sorted Alphabetically by Scientific Name					<i>Gilia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Gilia capitata</i> subsp. <i>staminea</i>	Pale Gilia	Polemoni-	-   -   B	- NA	Sandhills, flats (Mar–May)	
<input type="radio"/> <i>Gilia clivorum</i>	Purplespot Gilia	Polemoni-	-   -   C	- NA	Common. Open, grassy areas (Feb–Jun)	
<input type="radio"/> <i>Gilia</i> sp.	Gilia	Polemoni-	-   -   -	- NA	--	
<input type="radio"/> <i>Gilia tricolor</i> subsp. ?	Bird's-eye Gilia	Polemoni-	-   -   -	- NA	--	
<input type="radio"/> <i>Gilia tricolor</i> subsp. <i>diffusa</i>	Diffuse Bird's-eye Gilia	Polemoni-	-   -   B	- NA	Open grassland, hills, valleys (Mar–Jul)	
<input type="radio"/> <i>Gilia tricolor</i> subsp. <i>tricolor</i>	Bird's-eye Gilia	Polemoni-	-   -   B	- NA	Open, grassland, hills, valleys (Mar–May)	
<input type="radio"/> <i>Githopsis diffusa</i> subsp. <i>robusta</i>	Southern Bluecup	Campanul-	-   -   A2	- NA	Shaded or disturbed areas, burns (Apr–Jun)	
<input type="radio"/> <i>Githopsis specularioides</i>	Looking-glass Bluecup	Campanul-	-   -   B	- NA	Chaparral, oak woodland (Apr–May)	
<input type="radio"/> <i>Glaux maritima</i>	Sea Milkwort	Myrsin-	-   -   A1	- NP	Coastal salt marshes, saline meadows (May–Jul)	
<input type="radio"/> <i>Glinus lotoides</i>	Damascisa	Mollugin-	-   -   -	- ZA	Uncommon. Moist or seasonally dry margins of wetlands (Jun–Nov)	
<input type="radio"/> <i>Glycyrrhiza lepidota</i>	Wild Licorice	Fab-	-   -   C	- NP	In colonies, moist, gen open, disturbed areas incl streambanks, roadsides, alkaline soils or not (May–Jul)	
<input type="radio"/> <i>Gnaphalium</i> or <i>Pseudognaphalium</i> sp. (JM93: <i>Gnaphalium</i> sp.)	Everlasting Cudweed	Aster-	-   -   -	-	--	
<input type="radio"/> <i>Gnaphalium palustre</i>	Lowland Cudweed	Aster-	-   -   -	- NA	Arroyos, sandy streambeds, pond edges, potholes (May–Oct)	
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i> )	Great Valley Gumplant	Aster-	-   -   -	- NP	Sandy or saline bottomland, roadsides (May–Nov)	
<input type="radio"/> <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i> )	Hairy Gumplant	Aster-	-   -   C	- NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)	
<input type="radio"/> <i>Grindelia</i> sp.	Gumplant	Aster-	-   -   -	- NP	--	
<input type="radio"/> <i>Grindelia stricta</i> var. <i>platyphylla</i>	Pacific Gumplant	Aster-	-   -   A1	- NP	Coastal bluffs, dunes (All year)	
<input type="radio"/> <i>Helenium bigelovii</i>	Bigelow Sneezeweed	Aster-	-   -   A2	- NP	Wet meadows, marshes, bogs, fens, streambanks, lake margins (Jul–Aug)	
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	-   -   -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)	
<input type="radio"/> <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	-   1B.2   *A2	- NP	Open, grassy sites (Apr–Jun)	
<input type="radio"/> <i>Helianthus annuus</i>	Common Sunflower	Aster-	-   -   -	- NP	Disturbed areas, scrub, grassland, many other habitats (Jun–Oct)	
<input type="radio"/> <i>Helianthus californicus</i>	California Sunflower	Aster-	-   -   C	- NP	Springs, marshes, streambanks, canyons (Jul–Oct)	
<input type="radio"/> <i>Heliotropium curassavicum</i> var. <i>oculatum</i> (JM93: no var.)	Seaside Heliotrope	Boragin-	-   -   -	- NP	Moist to dry, saline to alkaline soils, gen near water (Feb–Oct)	
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i> )	Bristly Ox-tongue	Aster-	-   -   -	L ZAB	Common. Disturbed areas (All year)	
<input type="radio"/> <i>Hemizonia congesta</i> subsp. ?	Yellow Hayfield Tarweed	Aster-	-   -   -	- NA	--	

Herbaceous <i>Hemizonia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Hemizonia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>lutescens</i> (JM93: subsp. <i>congesta</i> )	Yellow Hayfield Tarweed	Aster-	-   -   C	- NA	Grassland, barrens, openings in chaparral and woodland, often on serpentine (Apr–Dec)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Aster-	-   -   -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)
<input type="radio"/> <i>Hemizonia</i> , <i>Centromadia</i> , or <i>Deinandra</i> sp. (JM93: <i>Hemizonia</i> sp.)	Spikeweed	Aster-	-   -   -	- NAP	- -
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i> )	Cow Parsnip	Api-	-   -   -	- NP	Moist places, wooded or open (Apr–Jul)
<input type="radio"/> <i>Herniaria hirsuta</i> var. <i>cinerea</i> (JM93: subsp.)	Ash-color Rupturewort	Caryophyll-	-   -   -	- ZA	Disturbed areas, alkaline hills, clay flats (Spring–fall)
<input type="radio"/> <i>Hesperevax acaulis</i> var. <i>ambusticola</i>	Dwarf Fire Evax	Aster-	-   -   A2	- NA	Clearings, barren slopes, burns (Apr–Jun)
<input type="radio"/> <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Aster-	-   -   -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	-   -   -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)
<input type="radio"/> <i>Hesperolinon breweri</i>	Brewer's Western Flax	Lin-	-   1B.2   *A2	- NA	Chaparral or grassland, occ on serpentine (May–Jun)
<input type="radio"/> <i>Hesperolinon californicum</i>	California Flax	Lin-	-   -   A2	- NA	Rocky areas, chaparral, grassland, occasionally on serpentine (Apr–Jul)
<input type="radio"/> <i>Hesperolinon</i> sp.	Flax	Lin-	-   -   -	- NA	
<input type="radio"/> <i>Heterotheca grandiflora</i>	Telegraph Weed	Aster-	-   -   -	- NA	Disturbed areas, dry streambeds, sand dunes (Jun–Oct(± all year))
<input type="radio"/> <i>Heterotheca oregona</i> var. <i>rudis</i>	Inland Oregon Goldenaster	Aster-	-   -   A1	- NP	Seasonally dry streambed (Jul–Oct)
<input type="radio"/> <i>Heterotheca oregona</i> var. <i>scaberrima</i>	Sticky Oregon Goldenaster	Aster-	-   -   A1	- NP	Seasonally dry streambeds (Jul–Oct)
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. ?	Sessileflower Goldenaster	Aster-	-   -   -	- NP	- -
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. <i>bolanderi</i>	Bolander Goldenaster	Aster-	-   -   C	- NP	Dunes, headlands, grassy coastal slopes (Jun–Sep)
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. <i>echioides</i>	Bristly Goldenaster	Aster-	-   -   -	- NP	Grassland, scrub, woodland, open forest, disturbed sites (Jul–Oct)
<input type="radio"/> <i>Heterotheca</i> sp.	Goldenaster	Aster-	-   -   -	- NP	- -
<input type="radio"/> <i>Heuchera micrantha</i>	Small-flower Alumroot	Saxifrag-	-   -   C	- NP	Moist, rocky banks and cliffs (Apr–Jul)
<input type="radio"/> <i>Hieracium albiflorum</i>	White Hawkweed	Aster-	-   -   B	- NP	Forest (May–Sep)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	-   -   -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Hoita macrostachya</i>	Leather Root	Fab-	-   -   B	- NP	Streamsides, marshes, spring-moist places (May–Aug)
<input type="radio"/> <i>Holocarpa heermannii</i>	Heermann Tarplant	Aster-	-   -   -	- NA	Grassland (May–Nov)
<input type="radio"/> <i>Holocarpa macradenia</i>	Santa Cruz Tarplant	Aster-	FT/CE   1B.1   *A1	- NA	Grassy areas, clay soil (Jun–Nov)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Holocarpa</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Holocarpa</i></b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Holocarpa obconica</i>	San Joaquin Tarplant	Aster-	-   -   B	- NA	Grassland (Apr–Nov)	
<input type="radio"/> <i>Holocarpa sp.</i>	Tarplant	Aster-	-   -   -	- NA	-	
<input type="radio"/> <i>Holocarpa virgata subsp. virgata</i>	Sticky Tarplant	Aster-	-   -   -	- NA	Grassland (May–Nov)	
<input type="radio"/> <i>Horkelia californica var. californica (JM93: subsp. californica)</i>	Few-lobed Calif. Horkelia	Ros-	-   -   A1	- NP	Grassy openings, edges of coastal scrub, especially n slopes (Mar–Sep)	
<input type="radio"/> <i>Horkelia californica var. elata (JM93: subsp. dissita)</i>	Many-lobed Calif. Horkelia	Ros-	-   -   A2	- NP	Shady meadow edges, seasonal streams, open chaparral (Jun–Sep)	
<input type="radio"/> <i>Horkelia californica var. frondosa (JM93: subsp. frondosa)</i>	Unlobed California Horkelia	Ros-	-   -   C	- NP	Coastal scrub, canyons, poison-oak thickets (May–Oct)	
<input type="radio"/> <i>Horkelia sp.</i>	Horkelia	Ros-	-   -   -	- NP	-	
<input type="radio"/> <i>Hydrocotyle ranunculoides</i>	Floating Marsh Pennywort	Arali-	-   -   A1	- NP	Lake margins, ponds, slow-moving streams (Mar–Aug)	
<input type="radio"/> <i>Hydrocotyle sp.</i>	Water / Marsh Pennywort	Arali-	-   -   -	- NP	-	
<input type="radio"/> <i>Hydrocotyle verticillata</i>	Whorled Marsh Pennywort	Arali-	-   -   A2	- NP	Lake margins, ponds, slow-moving streams, canals, seeps, springs, marshes (Apr–Sep)	
<input type="radio"/> <i>Hydrophyllum occidentale</i>	Western Waterleaf	Boragin-	-   -   A2	- NP	Moist, shaded slopes, woodland, meadows, streambanks, chaparral (May–Jul)	
<input type="radio"/> <i>Hypericum perforatum</i>	Klamathweed	Hyperic-	-   -   -	M ZP	Open, disturbed areas in many plant communities (May–Aug)	
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	-   -   -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	-   -   -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)	
<input type="radio"/> <i>Idahoia scapigera</i>	Flat-pod	Brassic-	-   -   A1	- NA	Moist ledges, slopes, meadows, foothills (Mar–May)	
<input type="radio"/> <i>Iris douglasiana</i>	Douglas Iris	Irid-	-   -   A2	- NP	Common. Grassy places, esp near coast (May–Jul)	
<input type="radio"/> <i>Iris longipetala</i>	Coast Iris	Irid-	-   4.2   *A1	- NP	Moist, coastal prairie or open coastal forest (Mar–Jun)	
<input type="radio"/> <i>Iris pseudacorus</i>	Yellow Flag	Irid-	-   -   -	L ZP	Common. Pond margins, estuaries (Apr–Jun)	
<input type="radio"/> <i>Iris sp.</i>	Iris	Irid-	-   -   -	-	-	
<input type="radio"/> <i>Iva axillaris (JM93: subsp. robustior)</i>	Poverty Weed	Aster-	-   -   -	- NP	Seasonally wet, saline habitats, roadsides (Apr–Oct)	
<input type="radio"/> <i>Jaumea carnosa</i>	Fleshy Jaumea	Aster-	-   -   C	- NP	Coastal salt marshes, bases of sea cliffs (Apr–Dec)	
<input type="radio"/> <i>Kickxia elatine</i>	Sharp Point Fluvellin	Plantagin-	-   -   -	- ZA	Disturbed, open places (Apr–Oct)	
<input type="radio"/> <i>Kopsiopsis strobilacea (JM93: Boschniakia)</i>	Cone Flower / Ground Cone	Orobanch-	-   -   A1	- NP	Open woodland, chaparral, generally on <i>Arctostaphylos</i> (Apr–Jun)	
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	-   -   -	- ZA	Roadsides, grassland (Jul–Nov)	

Herbaceous <i>Lactuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lactuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	-   -   -	- ZA	Abundant. Disturbed places (May–Oct)	
<input type="radio"/> <i>Lactuca sp.</i>	Wild Lettuce	Aster-	-   -   -	- ZA	- -	
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	-   -   -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)	
<input type="radio"/> <i>Laennecia coulteri</i> (JM93: <i>Conyza</i> )	Coulter's Horseweed	Aster-	-   -   A1	- NA	Disturbed sites, clayey or sandy soils, often seasonally wet, alkaline (May–Nov)	
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: <i>subsp. ramosissima</i> )	Common Hare-leaf	Aster-	-   -   -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)	
<input type="radio"/> <i>Lamium amplexicaule</i>	Clasping Henbit	Lami-	-   -   -	- ZAB	Disturbed sites, cult or abandoned fields (Apr–Sep)	
<input type="radio"/> <i>Lamium purpureum</i>	Red Henbit	Lami-	-   -   -	- ZA	Uncommon. Disturbed sites, meadows (Apr–Sep)	
<input type="radio"/> <i>Lasthenia californica subsp. californica</i> (JM93: <i>no subsp.</i> )	California Goldfields	Aster-	-   -   -	- NA	Many habitats (Feb–Jun)	
<input type="radio"/> <i>Lasthenia glabrata subsp. glabrata</i>	Yellow-ray Goldfields	Aster-	-   -   B	- NA	Saline places, vernal pools (Mar–May)	
<input type="radio"/> <i>Lasthenia microglossa</i>	Small-ray Goldfields	Aster-	-   -   B	- NA	Shaded slopes of woodland, chaparral, desert scrub (Mar–May)	
<input type="radio"/> <i>Lathyrus jepsonii var. californicus</i>	Bluff Pea	Fab-	-   -   B	- NP	Forest, open areas (Apr–Aug)	
<input type="radio"/> <i>Lathyrus jepsonii var. jepsonii</i>	Delta Tule Pea	Fab-	-   1B.2   *A2	- NP	Coastal, estuarine marshes (Apr–Aug)	
<input type="radio"/> <i>Lathyrus latifolius</i>	Perennial Sweet Pea	Fab-	-   -   -	- ZP	Disturbed areas, esp roadsides (Apr–May)	
<input type="radio"/> <i>Lathyrus sp.</i>	Pea	Fab-	-   -   -	-	- -	
<input type="radio"/> <i>Lathyrus vestitus var. vestitus</i>	Pale Purple Pacific Pea	Fab-	-   -   -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)	
<input type="radio"/> <i>Layia chrysanthemoides</i>	Chrysanthemum Tidytops	Aster-	-   -   A2	- NA	Grassy or open heavy soil, sometimes ± alkaline (Mar–Jun)	
<input type="radio"/> <i>Layia gaillardiioides</i>	Yellow Woodland Tidytops	Aster-	-   -   A2	- NA	Open or semi-shaded slopes, in sandy or clayey soil (including serpentine) (Mar–Aug)	
<input type="radio"/> <i>Layia hieracioides</i>	Tall Tidytops	Aster-	-   -   A2	- NA	Open, semi-shady, or disturbed sites, in light soil (Apr–Jul)	
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	-   -   C	- NA	Many habitats (Feb–Jul)	
<input type="radio"/> <i>Legenere limosa</i>	Legenere	Campanul-	-   1B.1   *A1	- NA	Wet areas, vernal pools, ponds (May–Jun)	
<input type="radio"/> <i>Lemna gibba</i>	Swollen Duckweed	Lemn-	-   -   -	- NP	Common. Fresh or brackish water -	
<input type="radio"/> <i>Lemna minor</i>	Common Lesser Duckweed	Lemn-	-   -   -	- NP	Common. Freshwater (Aug)	
<input type="radio"/> <i>Lemna sp.</i>	Duckweed	Lemn-	-   -   -	- NP	- -	
<input type="radio"/> <i>Lepidium dictyotum</i> (JM93: <i>var. dictyotum</i> )	Alkali Peppergrass	Brassic-	-   -   B	- NA	Saline soils, dry stream beds, roadsides, sandy flats, fields, meadows, dried pools (Mar–Jun)	
<input type="radio"/> <i>Lepidium didymum</i> (JM93: <i>Coronopus didymus</i> )	Lesser Swine Cress	Brassic-	-   -   -	- ZA	Common. Disturbed areas, fields, pastures (Mar–Jul)	

Herbaceous <i>Lepidium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lepidium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
○ <i>Lepidium draba</i> (JM93: <i>Cardaria</i> )	Heart-podded Hoary Cress	Brassic-	-   -   -	M ZP	Disturbed areas, saline soils, pastures, fields (Apr–Aug)	
○ <i>Lepidium latifolium</i>	Perennial Peppergrass	Brassic-	-   -   -	H ZP	Pastures, disturbed areas, fields, grassland, saline meadows, streambanks, sagebrush scrub, edge of marshes (Jun–Sep)	
○ <i>Lepidium latipes</i> (JM93: var. <i>latipes</i> )	Dwarf Peppergrass	Brassic-	-   -   B	- NA	Alkaline soils, vernal pool margins, salt marsh edges, pastures (Mar–Jun)	
○ <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i> )	Threadleaf Peppergrass	Brassic-	-   -   -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
○ <i>Lepidium oxycarpum</i>	Sharp-pod Peppergrass	Brassic-	-   -   B	- NA	Fields, vernal pool margins, alkaline flats (Mar–May)	
○ <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	-   -   -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)	
○ <i>Leptosiphon acicularis</i> (JM93: <i>Linanthus</i> )	Bristly Leptosiphon	Polemoni-	-   4.2   *A1	- NA	Grassy areas, woodland, chaparral (Apr–May)	
○ <i>Leptosiphon ambiguus</i> (JM93: <i>Linanthus</i> )	Serpentine Leptosiphon	Polemoni-	-   4.2   *A2	- NA	Grassy areas gen serpentine (Apr–May)	
○ <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i> )	Pinklobe Leptosiphon	Polemoni-	-   -   C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)	
○ <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i> )	Bicolor Leptosiphon	Polemoni-	-   -   -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)	
○ <i>Leptosiphon ciliatus</i> (JM93: <i>Linanthus</i> )	Whisker Brush	Polemoni-	-   -   B	- NA	Common. Open or wooded areas (Mar–Jul)	
○ <i>Leptosiphon parviflorus</i> (JM93: <i>Linanthus</i> )	Common Linanthus	Polemoni-	-   -   B	- NA	Abundant. Open or wooded areas (Mar–Jun)	
○ <i>Lessingia nemaclada</i>	Funnel-flower Lessingia	Aster-	-   -   A1	- NA	Open fields, roadsides, woodland borders, occasionally on serpentine soil (Jul–Oct)	
○ <i>Lessingia</i> sp.	Lessingia	Aster-	-   -   -	- NA	--	
○ <i>Lewisia rediviva</i> var. <i>rediviva</i> (JM93: no var.)	Bitter Root	Monti-	-   -   A2	- NP	Rocky, sandy ground, open conifer woodland, scrub (Mar–Jun)	
○ <i>Lilaeopsis masonii</i>	Mason's Lilaeopsis	Api-	CR   1B.1   *A2	- NP	Intertidal marshes, streambanks. (Jun–Aug)	
○ <i>Lilium pardalinum</i> subsp. <i>pardalinum</i>	Leopard Lily	Lili-	-   -   A2	- NP	Moist places (drier along coast) (May–Aug)	
○ <i>Limonium californicum</i>	Western Marsh-rosemary	Plumbagin-	-   -   B	- NP	Common; coastal dunes, salt marshes (Jul–Dec)	
○ <i>Limosella acaulis</i>	Southern Mudwort	Scrophulari-	-   -   A2	- NA	Wet, muddy places, generally fresh water (May–Oct)	
○ <i>Limosella australis</i> (JM93: <i>L. subulata</i> )	Delta Mudwort	Scrophulari-	-   2.1   -	- ZA	Muddy or sandy intertidal flats, brackish water (Apr)	
○ <i>Linanthus dichotomus</i> subsp. ? (JM93: no subsp.)	Evening Snow	Polemoni-	-   -   A2	- NA	--	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Linanthus</i>	Sorted Alphabetically by Scientific Name					<i>Linanthus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. <i>dichotomus</i> (JM93: no subsp.)	Evening Snow	Polemoni-	-   -   A2	- NA	Common. Drying openings, especially serpentine (Apr–Jun)	
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. <i>meridianus</i> (JM93: no subsp.)	Day-blooming Evening Snow	Polemoni-	-   -   A2	- NA	Drying openings, especially serpentine (Apr–Jun)	
<input type="radio"/> <i>Linanthus</i> or <i>Leptosiphon</i> sp. (JM93: <i>Linanthus</i> sp.)	Linanthus	Polemoni-	-   -   -	- NA	--	
<input type="radio"/> <i>Linum bienne</i>	Narrow-leaf Flax	Lin-	-   -   -	- ZP	Garden escape; grassland, woodland, disturbed places, esp coastal (Mar–Jun)	
<input type="radio"/> <i>Linum</i> sp.	Flax	Lin-	-   -   -	-	--	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	-   -   -	- NP	Open, grassy slopes (Mar–Apr)	
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	-   -   -	- NP	Shaded slopes (Feb–Jun)	
<input type="radio"/> <i>Lithophragma parviflorum</i> var. <i>parviflorum</i>	Prairie Starflower	Saxifrag-	-   -   B	- NP	Common. Open areas (Mar–Jul)	
<input type="radio"/> <i>Lithophragma</i> sp.	Woodland Star	Saxifrag-	-   -   -	- NP	--	
<input type="radio"/> <i>Lobularia maritima</i>	Sweet Alyssum	Brassic-	-   -   -	L ZP	Disturbed areas, fields (Mar–Oct)	
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i> )	California Fluffweed	Aster-	-   -   -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i> )	Daggerleaf Cottonrose	Aster-	-   -   -	- ZA	Bare or grassy openings, burns (Mar–Jul)	
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Api-	-   -   B	- NP	Woodland, brushy slopes (Apr–Jun)	
<input type="radio"/> <i>Lomatium caruifolium</i> var. <i>caruifolium</i>	Alkali Desertparsley	Api-	-   -   B	- NP	Wet, clay depressions, open grassland (Mar–May)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	-   -   -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Api-	-   -   -	- NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium nudicaule</i>	Swollen-stem Biscuit Root	Api-	-   -   -	- NP	Rocky slopes, flats, gen pine woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium</i> sp.	Biscuit Root	Api-	-   -   -	- NP	--	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	-   -   -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lotus corniculatus</i>	Bird's-foot Trefoil	Fab-	-   -   -	- ZP	Open, disturbed areas (Jun–Sep)	
<input type="radio"/> <i>Ludwigia hexapetala</i>	Uruguayan Primrose-willow	Onagr-	-   -   A1?	H NP	Lake margins, wetlands (May–Dec)	
<input type="radio"/> <i>Ludwigia peploides</i> subsp. <i>peploides</i>	Yellow Water Primrose	Onagr-	-   -   B	- NP	Lakeshores, streambanks, seasonal wetlands (May–Oct)	
<input type="radio"/> <i>Ludwigia</i> sp.	Water Primrose	Onagr-	-   -   -	- NP	--	
<input type="radio"/> <i>Lunaria annua</i>	Money Plant	Brassic-	-   -   -	- ZA	Disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Lupinus adsurgens</i>	Drew's Silky Lupine	Fab-	-   -   -	- NP	Dry slopes, montane forest (May–Jul)	
<input type="radio"/> <i>Lupinus affinis</i>	Sky Lupine	Fab-	-   -   A1	- NA	Uncommon. Open areas (Mar–May)	



<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Lupinus</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Lupinus</i></b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus concinnus</i>	Bajada Lupine	Fab-	-   -   A1	- NA	Common. Open or disturbed areas, burns (Mar–May)	
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	-   -   -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	
<input type="radio"/> <i>Lupinus latifolius</i> var. <i>latifolius</i>	Broadleaf Lupine	Fab-	-   -   -	- NP	Moist areas, open woodland (Apr–Jul)	
<input type="radio"/> <i>Lupinus luteolus</i>	Butter Lupine	Fab-	-   -   A1	- NA	Clearings, open or disturbed areas (May–Aug)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus pachylobus</i>	Big Pod Lupine	Fab-	-   -   B	- NA	Uncommon. Open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Lupinus</i> sp.	Lupine	Fab-	-   -   -	- NAP	--	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lupinus variicolor</i>	Varied Lupine	Fab-	-   -   A1	- NP	Coastal terraces, beaches (Apr–Jul)	
<input type="radio"/> <i>Lycopus americanus</i>	Cutleaf Bugleweed	Lami-	-   -   A1	- NP	Moist areas, marshes, streambanks (Aug–Sep)	
<input type="radio"/> <i>Lycopus asper</i>	Bugleweed	Lami-	-   -   A1	- NP	Moist areas, marshes, streambanks (Jun–Oct)	
<input type="radio"/> <i>Lycopus</i> sp.	Water Horehound / Bugleweed	Lami-	-   -   -	- NP	--	
<input type="radio"/> <i>Lythrum californicum</i>	California Loosestrife	Lythr-	-   -   B	- NP	Marshes, pond and stream margins (Apr–Sep)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	-   -   -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Lythrum</i> sp.	Loosestrife	Lythr-	-   -   -	-	--	
<input type="radio"/> <i>Lythrum tribracteatum</i>	Three-bract Loosestrife	Lythr-	-   -   -	- ZA	Wet areas, drying ponds (May–Jun)	
<input type="radio"/> <i>Madia anomala</i>	Plump-seed Tarweed	Aster-	-   -   A1	- NA	Openings in woodland or chaparral (Apr–Jun)	
<input type="radio"/> <i>Madia exigua</i>	Thread-stem Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites, in sandy or clayey soils, including serpentine (Apr–Jul)	
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	-   -   -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia</i> or <i>Anisocarpus</i> sp. (JM93: <i>Madia</i> sp.)	Tarweed	Aster-	-   -   -	- NA	--	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i> )	Fat False Solomon's Seal	Rusc-	-   -   -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i> )	Starry False Solomon's Seal	Rusc-	-   -   -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	

Herbaceous <i>Malacothrix</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Malacothrix</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Malacothrix coulteri</i>	Snake's Head Dandelion	Aster-	-   -   A1	- NA	Sandy, open areas, in coastal-sage scrub, grassland, desert (Mar–May)
<input type="radio"/> <i>Malacothrix floccifera</i>	Woolly / Flocked Dandelion	Aster-	-   -   A2	- NA	In loose soil of open burns, slides, in chaparral, pinyon/juniper woodland, yellow-pine forest (Mar–Nov)
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	-   -   -	- ZA	Disturbed places (Mar–Jun)
<input type="radio"/> <i>Malva parviflora</i>	Cheeseweed	Malv-	-   -   -	- ZA	Common. Disturbed places (Mar–May)
<input type="radio"/> <i>Malva pseudolavatera</i> (JM93: <i>Lavatera cretica</i> )	Cretan Mallow	Malv-	-   -   -	- ZAB	Uncommon. Disturbed places on coastal bluffs, dunes, occ inland (Apr–Jun)
<input type="radio"/> <i>Malva sp.</i>	Mallow	Malv-	-   -   -	- -	
<input type="radio"/> <i>Malvella leprosa</i>	Alkal-mallow	Malv-	-   -   -	- NP	Valleys, gen saline (Apr–Nov)
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i> )	California Man-root	Cucurbit-	-   -   -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)
<input type="radio"/> <i>Marah oregana</i> (JM93: <i>M. oreganus</i> )	Coast Man-root	Cucurbit-	-   -   C	- NP	Shrubby or open areas, forest edges (Mar–May)
<input type="radio"/> <i>Marah sp.</i>	Manroot	Cucurbit-	-   -   -	- NP	- -
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	-   -   -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i> )	Pineapple Weed	Aster-	-   -   -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)
<input type="radio"/> <i>Meconella californica</i>	California Meconella	Papaver-	-   -   A1	- NA	Open, rocky areas (Mar–May)
<input type="radio"/> <i>Medicago arabica</i>	Spotted Bur Clover	Fab-	-   -   -	- ZA	Disturbed and agricultural areas, fields, woodland (Mar–Jun)
<input type="radio"/> <i>Medicago lupulina</i>	Black Medick	Fab-	-   -   -	- ZA	Disturbed and agricultural areas, forest, mtns (Apr–Jul)
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	-   -   -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Medicago sativa</i>	Alfalfa	Fab-	-   -   -	- ZP	Disturbed, agricultural areas (Apr–Oct)
<input type="radio"/> <i>Melilotus albus</i> (JM93: <i>M. alba</i> )	White Sweetclover	Fab-	-   -   -	- ZAB	Locally abundant. Pastures, open disturbed sites (May–Sep)
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i> )	Sourclover	Fab-	-   -   -	- ZAB	Open, disturbed areas (Apr–Oct)
<input type="radio"/> <i>Melilotus officinalis</i>	Yellow Sweetclover	Fab-	-   -   -	- ZAB	Open fields, disturbed sites, cultivated (May–Aug)
<input type="radio"/> <i>Melissa officinalis</i>	Lemon Balm	Lami-	-   -   -	- ZP	Moist sites, meadows, fields (Jun–Sep)
<input type="radio"/> <i>Mentha arvensis</i>	Field Mint	Lami-	-   -   B	- NP	Moist places, fields (Jul–Oct)
<input type="radio"/> <i>Mentha piperita</i>	Peppermint	Lami-	-   -   -	- ZP	Moist places, fields (Jul–Oct)
<input type="radio"/> <i>Mentha pulegium</i>	Pennyroyal	Lami-	-   -   -	M ZP	Moist places, fields (Jul–Oct)
<input type="radio"/> <i>Mentha sp.</i>	Mint	Lami-	-   -   -	- -	

Herbaceous <i>Mentha</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Mentha</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Mentha spicata</i> (JM93: var. <i>spicata</i> )	Spearmint	Lami-	-   -   -	- ZP	Moist places, marshes, lakeshores, fields (Jul–Oct)	
<input type="radio"/> <i>Mentzelia affinis</i>	Hydra Blazing Star	Loas-	-   -   A2	- NA	Rocky or gray-white soils in grassland, woodland, creosote-bush scrub (Apr–May)	
<input type="radio"/> <i>Mentzelia dispersa</i>	Nevada Blazing Star	Loas-	-   -   A2	- NA	Loamy to sandy or rocky slopes, roadsides (May–Aug)	
<input type="radio"/> <i>Mentzelia lindleyi</i>	Lindley Blazing Star	Loas-	-   -   A2	- NA	Rocky, open slopes, coastal-sage scrub, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i> )	California Saxifrage	Saxifrag-	-   -   -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Micropus amphibolus</i>	Mount Diablo Cottonweed	Aster-	-   3.2   *A1	- NA	Openings on slopes, ridges, shallow soils, sedimentary or volcanic rocks (Mar–Jun)	
<input type="radio"/> <i>Micropus californicus</i> var. ?	Slender Cottonweed	Aster-	-   -   -	- NA	--	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Aster-	-   -   -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris acuminata</i>	Sierra Foothill Silverpuffs	Aster-	-   -   B	- NA	Grassy, open rocky or clay soil (Apr–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. ?	Douglas Silverpuffs	Aster-	-   -   -	- NA	--	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	-   -   -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris elegans</i>	Nodding Silverpuffs	Aster-	-   -   A2	- NA	Generally inland clay grassland, often near vernal pools (Apr–Jun)	
<input type="radio"/> <i>Microseris</i> sp.	Silverpuffs	Aster-	-   -   -	- NA	--	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i> )	Slender Annual Phlox	Polemoni-	-   -   -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus cardinalis</i>	Scarlet Monkeyflower	Phrym-	-   -   C	- NP	Moist to wet places along streams, seepage areas (May–Sep)	
<input type="radio"/> <i>Mimulus douglasii</i>	Mickey Mouse Monkeyflower	Phrym-	-   -   A2	- NA	Bare clay, serpentine or granitic soils; generally along upper banks of small creeks (Feb–Apr)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	-   -   -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Mimulus latidens</i>	Broadtooth Monkeyflower	Phrym-	-   -   A2	- NA	Vernally wet depressions (Apr–Jun)	
<input type="radio"/> <i>Mimulus pilosus</i>	Downy Monkeyflower	Phrym-	-   -   A2	- NA	Moist, sandy areas, especially by small streams, disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Minuartia californica</i>	California Sandwort	Caryophyll-	-   -   A2	- NA	Gravelly, sandy slopes, grassy ridges, chaparral, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Minuartia douglasii</i>	Douglas' Stitchwort	Caryophyll-	-   -   B	- NA	Dry, rocky, slopes, flats in chaparral, oak and pine woodland, often serpentine (Spring–early summer)	
<input type="radio"/> <i>Minuartia pusilla</i>	Annual Sandwort	Caryophyll-	-   -   A2	- NA	Plains, open pine forest, chaparral slopes (Spring–summer)	

Herbaceous <i>Mirabilis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Mirabilis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Mirabilis laevis</i> var. <i>crassifolia</i> (JM93: <i>M. californica</i> )	Wishbone Bush	Nyctagin-	-   -   A1	- NP	Grassy areas, chaparral, dunes, dry rocky areas, washes Dec–Jun)	
<input type="radio"/> <i>Moehringia macrophylla</i>	Big-leaf Sandwort	Caryophyll-	-   -   A2	- NP	Moist, shaded slopes, rocky ridges, summits, pine, oak forests, serpentine (Spring–summer)	
<input type="radio"/> <i>Monardella douglasii</i> (JM93: subsp. <i>douglasii</i> )	Douglas Coyotemint	Lami-	-   -   A2	- NA	Grassland, oak woodland, chaparral, serpentine outcrops (May–Nov)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>M. globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i> )	Coyote-mint	Lami-	-   -   -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Monolopia major</i>	Cupped Monolopia	Aster-	-   -   B	- NA	Grassland, bare clay (Feb–Jul)	
<input type="radio"/> <i>Montia fontana</i>	Water Chickweed	Monti-	-   -   C	- NA	Common. Ponds, streams, vernal pools, seeps (Jul–Aug)	
<input type="radio"/> <i>Muilla maritima</i>	Common Muilla	Themid-	-   -   B	- NP	Coastal scrub, grassland, woodland, valleys (Mar–Jun)	
<input type="radio"/> <i>Myosotis latifolia</i>	Broadleaved Forget-me-not	Boragin-	-   -   -	L ZP	Moist, disturbed, shady places (Feb–Jul)	
<input type="radio"/> <i>Myriophyllum sibiricum</i>	Siberian Water-milfoil	Halorag-	-   -   -	- NP	Ponds, streams, lakes (Jun–Sep)	
<input type="radio"/> <i>Myriophyllum</i> sp.	Water-milfoil	Halorag-	-   -   -	- -	- -	
<input type="radio"/> <i>Najas marina</i>	Holly-leaved Water-nymph	Hydrocharit-	-   -   -	- NA	Ponds, lakes, marshes, rivers (Jul–Aug)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i> )	Water Cress	Brassic-	-   -   -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia atractylodes</i>	Holly-leaf Navarretia	Polemoni-	-   -   A2	- NA	Open, rocky or sandy areas (May–Jul)	
<input type="radio"/> <i>Navarretia intertexta</i> subsp. <i>intertexta</i>	Needle-leaved Navarretia	Polemoni-	-   -   A1	- NA	Open, seasonally wet areas, meadows, vernal pools (May–Jul)	
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	-   -   C	- NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)	
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Polemoni-	-   -   C	- NA	Open, slopes, gravel, clay (May–Jul)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	-   -   -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	-   -   -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Boragin-	-   -   -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Nemophila parviflora</i> var. <i>parviflora</i>	Small-flower Nemophila	Boragin-	-   -   C	- NA	Woodland, forest, roadsides, slopes (Mar–Jul)	
<input type="radio"/> <i>Nemophila pedunculata</i>	Meadow Nemophila	Boragin-	-   -   B	- NA	Common. Ocean bluffs, grassland, slopes, meadows, sandbars, fields, woodland, streambanks (Feb–Jul)	
<input type="radio"/> <i>Nicotiana attenuata</i>	Coyote Tobacco	Solan-	-   -   A1	- NA	Open, well-drained slopes (May–Oct)	

Herbaceous <i>Nicotiana</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Nicotiana</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Nicotiana quadrivalvis</i>	Indian Tobacco	Solan-	-   -   B	- NA	Open, well-drained washes, slopes (May–Oct)
<input type="radio"/> <i>Nuttallanthus texanus</i> (JM93: <i>Linaria canadensis</i> )	Blue Toadflax	Plantagin-	-   -   B	- NA	Sand or gravel (Mar–May)
<input type="radio"/> <i>Nymphaea mexicana</i>	Yellow Or Banana Waterlily	Nymphae-	-   -   -	- ZP	Lakes, ponds, slow streams (Apr–Jul)
<input type="radio"/> <i>Nymphaea odorata</i>	Fragrant Or White Waterlily	Nymphae-	-   -   -	- ZP	Quiet waters, ponds, edges of lakes (Apr–Aug)
<input type="radio"/> <i>Oenanthe sarmentosa</i>	Water Parsley	Api-	-   -   -	- NP	Streams, marshes, ponds, gen aquatic (Jun–Oct)
<input type="radio"/> <i>Oenothera deltoides</i> subsp. <i>howellii</i>	Antioch Dunes Evening-primrose	Onagr-	FE/CE   1B.1   *A2	- NP	Sandy bluffs, dunes (Mar–Sep)
<input type="radio"/> <i>Oenothera elata</i> subsp. <i>hirsutissima</i>	Hairy Evening-primrose	Onagr-	-   -   -	- NBP	Moist places, generally inland (Jun–Sep)
<input type="radio"/> <i>Oenothera elata</i> subsp. <i>hookeri</i>	Hooker Evening-primrose	Onagr-	-   -   B	- NB	Moist, coastal, slightly inland, sandy bluffs (Jun–Sep)
<input type="radio"/> <i>Orobanche bulbosa</i>	Chaparral Broom-rape	Orobanch-	-   -   A2	- NP	Openings in chaparral, generally on <i>Adenostoma fasciculatum</i> (Apr–Jun)
<input type="radio"/> <i>Orobanche californica</i> subsp. <i>jepsonii</i>	Jepson Broomrape	Orobanch-	-   -   A1	- NP	Uncommon. Gen dry flats, slopes. Root parasite on sunflower family (Jul–Sep)
<input type="radio"/> <i>Orobanche fasciculata</i>	Clustered Broomrape	Orobanch-	-   -   B	- NP	Dry, gen bare places. Root parasite on various shrubs (Apr–Jul)
<input type="radio"/> <i>Orobanche uniflora</i>	Naked Broom-rape	Orobanch-	-   -   C	- NP	Generally moist places, on herbs, especially <i>Sedum</i> , <i>Saxifragaceae</i> , <i>Asteraceae</i> (Apr–Jul)
<input type="radio"/> <i>Orobanche vallicola</i>	Sharp-lobe Broom-rape	Orobanch-	-   -   A2	- NP	Riparian woodland, on <i>Sambucus nigra</i> (Jul–Sep)
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i> )	Sweet-cicely	Api-	-   -   -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Osmorhiza brachypoda</i>	California Sweet Cicely	Api-	-   -   B	- NP	Moist ravines, conifer forest, woodland (Mar–May)
<input type="radio"/> <i>Oxalis corniculata</i>	Creeping Wood Sorrel	Oxalid-	-   -   -	- ZP	Disturbed areas (Most of Year)
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	-   -   -	M ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)
<input type="radio"/> <i>Oxalis pilosa</i> (JM93: <i>O. albicans</i> subsp. <i>pilosa</i> )	Hairy Wood Sorrel	Oxalid-	-   -   A2	- NP	Coastal grassland, scrub, chaparral (Feb–Sep)
<input type="radio"/> <i>Oxalis</i> sp.	Wood Sorrel	Oxalid-	-   -   -	- -	
<input type="radio"/> <i>Packera breweri</i> (JM93: <i>Senecio</i> )	Brewer's Ragwort	Aster-	-   -   B	- NP	Common. Seasonally damp grassland, oak savanna, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Papaver californicum</i>	Fire Poppy	Papaver-	-   -   A2	- NA	Burns, disturbed areas, open woodland (Apr–May)
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i> )	Wind Poppy	Papaver-	-   -   -	- NA	Grassy areas, openings in chaparral (Apr–May)
<input type="radio"/> <i>Pectocarya penicillata</i>	Northern Pectocarya	Boragin-	-   -   A1	- NA	Disturbed sites, roadsides (Feb–May)
<input type="radio"/> <i>Pectocarya pusilla</i>	Little Pectocarya	Boragin-	-   -   B	- NA	Dry, semi-barren sites in grassland, chaparral, woodland, roadsides (Mar–Jun)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Pedicularis</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Pedicularis</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	-   -   -	- NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Pediomelum californicum</i>	Indian Breadroot	Fab-	-   -   A1	- NP	Open chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Penstemon centranthifolius</i>	Scarlet Bugler	Plantagin-	-   -   A2	- NP	Dry, open chaparral or oak woodland (Apr–Jul)	
<input type="radio"/> <i>Penstemon heterophyllus</i> var. <i>heterophyllus</i>	Foothill Penstemon	Plantagin-	-   -   B	- NP	Grassland, chaparral, forest openings (Apr–Jul)	
<input type="radio"/> <i>Pentachaeta exilis</i> subsp. <i>exilis</i>	Meager Pentachaeta	Aster-	-   -   A1	- NA	Grassland, foothill woodland (Mar–May)	
<input type="radio"/> <i>Perideridia californica</i>	California Yampah	Api-	-   -   A2	- NP	Damp soil by streams (Apr–May)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	-   -   -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Perideridia oregana</i>	Oregon Yampah	Api-	-   -   A2	- NP	Open flats or slopes, pine/oak woodland (Jul–Aug)	
<input type="radio"/> <i>Perideridia</i> sp.	Yampah	Api-	-   -   -	- NP	- -	
<input type="radio"/> <i>Persicaria amphibia</i> (JM93: <i>Polygonum amphibium</i> var. <i>emersum</i> , <i>stipulaceum</i> )	Water Smartweed	Polygon-	-   -   -	- NP	Shallow lakes, streams, shores (Jun–Nov)	
<input type="radio"/> <i>Persicaria hydropiperoides</i> (JM93: <i>Polygonum</i> )	False Waterpepper	Polygon-	-   -   A2	- NP	Wet banks, shallow water, marshes, moist prairies (Jun–Oct)	
<input type="radio"/> <i>Persicaria lapathifolia</i> (JM93: <i>Polygonum lapathifolium</i> )	Willow Weed	Polygon-	-   -   C	- NA	Moist places, disturbed areas, roadsides, fields (Jun–Oct)	
<input type="radio"/> <i>Persicaria maculosa</i> (JM93: <i>Polygonum persicaria</i> )	Lady's Thumb	Polygon-	-   -   -	- ZA	Moist disturbed areas, field (Jun–Nov)	
<input type="radio"/> <i>Persicaria punctata</i> (JM93: <i>Polygonum punctatum</i> )	Water Smartweed	Polygon-	-   -   C	- NAP	Shallow water, shores, marshes, floodplain forest (Jun–Nov)	
<input type="radio"/> <i>Petunia parviflora</i>	Wild Petunia	Solan-	-   -   A1	- NA	Open washes, dry streambeds (Apr–Aug)	
<input type="radio"/> <i>Phacelia breweri</i>	Brewer Phacelia	Boragin-	-   -   A2	- NP	Rocky soils, slopes, chaparral, oak woodland (Mar–Jun)	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	-   -   -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia ciliata</i>	Field Phacelia	Boragin-	-   -   B	- NA	Clay or gravelly slopes in grassland, fields (Feb–Jun)	
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	-   -   -	- NA	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Phacelia divaricata</i>	Divaricate Phacelia	Boragin-	-   -   A2	- NA	Open areas, chaparral, woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Phacelia heterophylla</i> subsp. <i>virgata</i>	Varileaf Phacelia	Boragin-	-   -   -	- NBP	Slopes, flats, roadsides (May–Jul)	
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	-   -   -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Phacelia malvifolia</i>	Stinging Phacelia	Boragin-	-   -   A2	- NA	Sandy or gravelly soils, scrub, conifer forest (Apr–Jul)	
<input type="radio"/> <i>Phacelia nemoralis</i> subsp. <i>nemoralis</i>	Bristly Phacelia	Boragin-	-   -   B	- NBP	Moist slopes, streambanks, mixed-evergreen forest (Apr–Jul)	

Herbaceous <i>Phacelia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Phacelia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Phacelia ramosissima</i> (JM93: var. <i>ramosissima</i> , <i>latifolia</i> )	Branched Phacelia	Boragin-	-   -   A2	- NP	Diverse habitats incl salt marshes, canyons (Apr–Oct)	
<input type="radio"/> <i>Phacelia rattanii</i>	Rattan Phacelia	Boragin-	-   -   A1	- NA	Shaded crevices, steep slopes (May–Jul)	
<input type="radio"/> <i>Phacelia tanacetifolia</i>	Tansy-leaf Phacelia	Boragin-	-   -   B	- NA	Sandy to gravelly slopes, open areas (Mar–May)	
<input type="radio"/> <i>Pholistoma auritum</i> var. <i>auritum</i>	Fiesta Flower	Boragin-	-   -   B	- NA	Ocean bluffs, talus slopes, woodland, streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Pholistoma membranaceum</i>	White Fiesta Flower	Boragin-	-   -   -	- NA	Beaches, bluffs, ravines, wooded slopes, desert washes (Feb–May)	
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>macrophyllum</i> (JM93: <i>P. macrophyllum</i> )	Big-leaf Mistletoe	Visc-	-   -   C	- NP	On trees other than <i>Quercus</i> (especially <i>Alnus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Platanus</i> , <i>Populus</i> , <i>Robinia</i> , <i>Salix</i> ) (Dec–Mar)	
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>tomentosum</i> (JM93: <i>P. villosum</i> )	Oak Mistletoe	Visc-	-   -   -	- NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)	
<input type="radio"/> <i>Phyla lanceolata</i>	Fog-fruit	Verben-	-   -   A1	- NP	Wet places, marshes (May–Nov)	
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i> )	Lemon Verbena	Verben-	-   -   -	- NP	Wet places, pond margins (May–Nov)	
<input type="radio"/> <i>Phyla</i> sp.	Verbena / Fog-fruit	Verben-	-   -   -	- NP	- -	
<input type="radio"/> <i>Piperia elegans</i> subsp. <i>elegans</i> (JM93: no subsp.)	Elegant Rein Orchid	Orchid-	-   -   B	- NP	Generally dry, open sites, scrub, conifer forest (May–Sep)	
<input type="radio"/> <i>Piperia elongata</i>	Wood Rein-orchid	Orchid-	-   -   A2	- NP	Generally dry sites, scrub, chaparral, mixed-evergreen or conifer forest (May–Jul)	
<input type="radio"/> <i>Piperia</i> sp.	Rein Orchid	Orchid-	-   -   -	- NP	- -	
<input type="radio"/> <i>Piperia unalascensis</i>	Alaska Rein Orchid	Orchid-	-   -   A1	- NP	Generally dry sites, scrub, woodland, forest (May–Aug)	
<input type="radio"/> <i>Plagiobothrys acanthocarpus</i>	Adobe Popcornflower	Boragin-	-   -   B	- NA	Vernal pools, moist clay soil (Mar–May)	
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Boragin-	-   -   -	- NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)	
<input type="radio"/> <i>Plagiobothrys canescens</i> var. <i>canescens</i> (JM93: no var.)	Valley Popcornflower	Boragin-	-   -   C	- NA	Common. Grassland, woodland, coastal scrub, desert scrub, roadsides (Mar–Jun)	
<input type="radio"/> <i>Plagiobothrys fulvus</i> var. <i>campestris</i> (JM93: no var.)	Field Popcornflower	Boragin-	-   -   B	- NA	Grassland, open woodland, loamy, sandy, or gravelly soil (Mar–May)	
<input type="radio"/> <i>Plagiobothrys humistratus</i>	Low Popcornflower	Boragin-	-   -   B	- NA	Vernal pools, wet places, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys infectivus</i>	Dye Popcornflower	Boragin-	-   -   A1	- NA	Open grassland, friable clay to sandy soils (Mar–May)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	-   -   -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	-   -   -	-	- -	

Herbaceous <i>Plagiobothrys</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Plagiobothrys</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Plagiobothrys stipitatus</i> var. <i>micranthus</i>	Valley Popcornflower	Boragin-	-   -   -	- NA	Common. Vernal pools, wet sites in grassland, conifer forest, sagebrush scrub (Apr–Jul)
<input type="radio"/> <i>Plagiobothrys stipitatus</i> var. <i>stipitatus</i>	Showy Great Popcornflower	Boragin-	-   -   C	- NA	Vernal pools, wet sites in grassland (Mar–Jun)
<input type="radio"/> <i>Plagiobothrys trachycarpus</i>	Rough-nutlet Popcornflower	Boragin-	-   -   B	- NA	Shallow vernal pools, wet places in grassland, scrub, chaparral, woodland (Mar–May)
<input type="radio"/> <i>Plantago coronopus</i>	Cut-leaf Plantain	Plantagin-	-   -   -	- NA	Coastal bluffs, salt marshes, trampled places, chaparral, grassy flats (Apr–Jul)
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	-   -   -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	-   -   -	L ZA	Common. Disturbed areas (Apr–Aug)
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Plantagin-	-   -   -	- ZAP	Disturbed areas (Apr–Sep)
<input type="radio"/> <i>Plantago subnuda</i>	Coast Plantain	Plantagin-	-   -   A1	- NP	Wet meadows, ditches, coastal bluffs, marshes (May–Sep)
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	-   -   -	- NA	Open grassland, sandy soil, burns (Mar–May)
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: <i>subsp. ciliosa, insignis</i> )	Longspur Plectritis	Valerian-	-   -   B	- NA	Common. Open, partly shaded slopes (Mar–Jun)
<input type="radio"/> <i>Plectritis congesta</i> subsp. <i>brachystemon</i> (JM93: <i>P. brachystemon</i> )	Pale Plectritis	Valerian-	-   -   B	- NA	Common. Coastal bluffs, open, partly shaded slopes (Mar–Jun)
<input type="radio"/> <i>Plectritis congesta</i> subsp. <i>congesta</i>	Sea Blush	Valerian-	-   -   A2	- NA	Coastal bluffs, open, partly shaded slopes (Mar–Jun)
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	-   -   -	- NA	Common. Open, partly shaded slopes (Mar–Jun)
<input type="radio"/> <i>Pluchea odorata</i> var. <i>odorata</i> (JM93: <i>no var.</i> )	Saltmarsh-fleabane	Aster-	-   -   B	- NP	Moist, often saline valley bottoms (Jun–Nov)
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	-   -   -	- NA	Grassy, brushy areas (Mar–Jun)
<input type="radio"/> <i>Polycarpon tetraphyllum</i>	Four-leaved Allseed	Caryophyll-	-   -   -	- ZA	Disturbed shaded areas, roadsides (Spring–fall)
<input type="radio"/> <i>Polygala californica</i>	California Milkwort	Polygal-	-   -   A1	- NP	Coastal prairie and forest, chaparral, occasionally on serpentine (Apr–Jul)
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i> )	Knotweed	Polygon-	-   -   -	- ZA	Disturbed places (May–Nov)
<input type="radio"/> <i>Polygonum</i> or <i>Persicaria</i> sp. (JM93: <i>Polygonum</i> sp.)	Knotweed / Smartweed	Polygon-	-   -   -	- --	
<input type="radio"/> <i>Portulaca oleracea</i>	Purslane	Portulac-	-   -   -	- ZP	Disturbed soil (Late spring-early fall)
<input type="radio"/> <i>Potamogeton crispus</i>	Crisp-leaved Pondweed	Potamogeton-	-   -   -	M ZP	Uncommon. Shallow water, ponds, streams. (Jul–Sep)
<input type="radio"/> <i>Potamogeton illinoensis</i>	Shining Pondweed	Potamogeton-	-   -   -	- NP	Lakes, ponds, streams (Jun–Aug)
<input type="radio"/> <i>Potamogeton nodosus</i>	Long-leaved Pondweed	Potamogeton-	-   -   -	- NP	Shallow water, lakes, ponds, streams (May–Aug)



Herbaceous <i>Potamogeton</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Potamogeton</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Potamogeton</i> or <i>Stuckenia</i> sp. (JM93: <i>Potamogeton</i> sp.)	Pondweed	Potamogeton-	-   -   -	- NP	--	
<input type="radio"/> <i>Potentilla anserina</i> subsp. <i>pacifica</i>	Pacific Cinquefoil	Ros-	-   -   A1	- NP	Coastal wetlands, often brackish (Mar–Oct)	
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i> )	Hooker Fairy Bells	Lili-	-   -   C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)	
<input type="radio"/> <i>Prosartes smithii</i> (JM93: <i>Disporum</i> )	Large-flower Fairy Bells	Lili-	-   -   A1?	- NP	Moist, shady coastal, montane forest (Mar–Jun)	
<input type="radio"/> <i>Prunella vulgaris</i> var. ?	European Selfheal	Lami-	-   -   -	- ZP	--	
<input type="radio"/> <i>Prunella vulgaris</i> var. <i>lanceolata</i>	Narrow-leaved Selfheal	Lami-	-   -   A1	- NP	Moist areas, gen conifer forest, woodland (May–Sep)	
<input type="radio"/> <i>Prunella vulgaris</i> var. <i>vulgaris</i>	European Selfheal	Lami-	-   -   -	- ZP	Moist, disturbed sites (Jun–Sep)	
<input type="radio"/> <i>Pseudognaphalium beneolens</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>beneolens</i> )	Scented White Everlasting	Aster-	-   -   C	- NB	Dry, open slopes, disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Pseudognaphalium biolettii</i> (JM93: <i>Gnaphalium bicolor</i> )	Two-color Everlasting	Aster-	-   -   A2	- NP	Rocky slopes, roadsides, dunes, coastal scrub, chaparral, oak woodland (Apr–Jun)	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i> )	California Everlasting	Aster-	-   -   -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i> )	Weedy Cudweed	Aster-	-   -   -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)	
<input type="radio"/> <i>Pseudognaphalium microcephalum</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>microcephalum</i> )	White Everlasting	Aster-	-   -   B	- NB	Grassy hillsides, gravelly canyon bottoms, chaparral, coastal-sage scrub (Jun–Aug)	
<input type="radio"/> <i>Pseudognaphalium ramosissimum</i> (JM93: <i>Gnaphalium</i> )	Pink Everlasting	Aster-	-   -   -	- NB	Dry, open slopes, woodland, sandy fields, dunes (Jul–Sep)	
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i> )	Cotton-batting Plant	Aster-	-   -   -	- NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)	
<input type="radio"/> <i>Psilocarphus brevissimus</i> var. <i>brevissimus</i>	Dwarf Woollyheads	Aster-	-   -   B	- NA	Drying edges of vernal pools, mud flats, drainages (May–Jun)	
<input type="radio"/> <i>Psilocarphus oregonus</i>	Oregon Woollyheads	Aster-	-   -   B	- NA	Drying clay of vernal pools, drainages, moist rocky slopes (Mar–Jul)	
<input type="radio"/> <i>Psilocarphus tenellus</i> (JM93: var. <i>tenellus</i> )	Slender Woolly-marbles	Aster-	-   -   -	- NA	Common. Dry, seasonally moist slopes, flats, burns, trails, rarely vernal pools (Mar–Jul)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	-   -   -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	-   -   B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)	
<input type="radio"/> <i>Ranunculus aquatilis</i> var. <i>diffusus</i> (JM93: var. <i>capillaceus</i> )	Thread-leaf Water Buttercup	Ranuncul-	-   -   -	- NP	Ponds, lakes, streams (Mar–Sep)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Ranunculus</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Ranunculus</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	-   -   -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus canus</i> var. <i>canus</i> (JM93: no var.)	Sacramento Valley Buttercup	Ranuncul-	-   -   B	- NP	Grassland, open oak woodlan (Mar–Jul)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	-   -   -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus lobbii</i>	Lobb's Aquatic Buttercup	Ranuncul-	-   4.2   *A2	- NA	Ponds (Mar–May)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	-   -   -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)	
<input type="radio"/> <i>Ranunculus occidentalis</i> var. <i>occidentalis</i> (JM93: no var.)	Western Buttercup	Ranuncul-	-   -   A2	- NP	Grassy slopes in meadows or open woodland (Mar–Jul)	
<input type="radio"/> <i>Ranunculus orthorhynchus</i> var. <i>bloomeri</i>	Bloomer Buttercup	Ranuncul-	-   -   A1	- NP	Meadows, marshy areas (Mar–May)	
<input type="radio"/> <i>Ranunculus repens</i>	Creeping Buttercup	Ranuncul-	-   -   -	L ZP	Meadows, road banks, disturbed places, edges of marshes, streams (All year)	
<input type="radio"/> <i>Ranunculus sceleratus</i> var. <i>sceleratus</i>	Cursed Crowsfoot	Ranuncul-	-   -   -	- NA	Ponds, riverbanks (Apr–Jun,Oct)	
<input type="radio"/> <i>Ranunculus</i> sp.	Buttercup	Ranuncul-	-   -   -	- -		
<input type="radio"/> <i>Raphanus raphanistrum</i>	Jointed Charlock	Brassic-	-   -   -	- ZA	Disturbed areas, fields (Apr–Jul)	
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	-   -   -	L ZA	Disturbed areas, fields (May–Jul)	
<input type="radio"/> <i>Rigiopappus leptocladus</i>	Bristle-head	Aster-	-   -   B	- NA	Grassy sites, openings among shrubs or trees (Apr–Jul)	
<input type="radio"/> <i>Rorippa curvisiliqua</i>	Yellow Water Cress	Brassic-	-   -   A2	- NA	Uncommon. Streambanks, marshy ground, seepage areas, mud flats, meadows (May–Oct)	
<input type="radio"/> <i>Rorippa palustris</i> subsp. <i>palustris</i> (JM93: var. <i>occidentalis</i> )	Marsh Water Cress	Brassic-	-   -   A2	- NA	Generally wet places (Mar–Sep)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	-   -   -	M ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex conglomeratus</i>	Green Dock	Polygon-	-   -   -	- ZP	Common. Moist places (May–Aug)	
<input type="radio"/> <i>Rumex crassus</i> (JM93: <i>R. salicifolius</i> var. <i>crassus</i> )	Dense Willow Dock	Polygon-	-   -   A2	- NP	Coastal dunes, sandy shores, marshes (Feb–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	-   -   -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex fueginus</i> (JM93: <i>R. maritimus</i> )	Golden Dock	Polygon-	-   -   A2	- NA	Riparian, disturbed places, shores, marshes, bogs, wet meadows (May–Aug)	
<input type="radio"/> <i>Rumex obtusifolius</i>	Bitter Dock	Polygon-	-   -   -	- ZP	Moist places (May–Sep)	
<input type="radio"/> <i>Rumex occidentalis</i>	Western Dock	Polygon-	-   -   A1	- NP	Uncommon. Wet habitats (May–Aug)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	-   -   -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i> )	Thin-leaf Willow Dock	Polygon-	-   -   B	- NP	Wet places, margins, rocky slopes (May–Jul)	
<input type="radio"/> <i>Rumex</i> sp.	Dock	Polygon-	-   -   -	- -		

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Rumex</i>	Sorted Alphabetically by Scientific Name					<i>Rumex</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Rumex transitorius</i> (JM93: <i>R. salicifolius</i> var. <i>transitorius</i> )	Ovate Willow Dock	Polygon-	-   -   -	- NP	Coastal dunes, wet margins of meadows (Apr–Jun)	
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	-   -   -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Ruppia maritima</i>	Straight Ditch-grass	Ruppi-	-   -   -	- NP	Brackish or saline waters, often coastal, marshes, ponds, sloughs (Mar–Aug)	
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Caryophyll-	-   -   -	- NA	Sandy disturbed areas, river bars, streamsides (Spring–early summer)	
<input type="radio"/> <i>Sagina decumbens</i> subsp. <i>occidentalis</i>	Western Pearlwort	Caryophyll-	-   -   C	- NA	Dry streams, chaparral, grassy areas, rock outcrops, vernal pools (Spring–early summer)	
<input type="radio"/> <i>Sagittaria latifolia</i>	Wappato / Tule Potato / Arrowhead	Alismat-	-   -   A1	- NP	Ponds, slow streams (Jul–Aug)	
<input type="radio"/> <i>Salicornia depressa</i> (JM93: <i>S. europaea</i> )	Slender Glasswort	Chenopodi-	-   -   -	- NA	Salt marshes, alkaline flats (Jul–Sep)	
<input type="radio"/> <i>Salicornia pacifica</i> (JM93: <i>S. virginica</i> )	Pacific Pickleweed	Chenopodi-	-   -   -	- NP	Salt marshes, alkaline flats (Jul–Nov)	
<input type="radio"/> <i>Salsola soda</i>	Fleshy Russian Thistle	Chenopodi-	-   -   -	M ZA	Uppermost intertidal zone, saline or muddy flats, open areas in salt marshes (Jul–Oct)	
<input type="radio"/> <i>Salsola tragus</i>	Russian Thistle	Chenopodi-	-   -   -	L ZAP	Common. Disturbed places (Jul–Oct)	
<input type="radio"/> <i>Salvia columbariae</i>	Chia	Lami-	-   -   B	- NA	Dry, disturbed sites, chaparral, coastal-sage scrub (Mar–Jun)	
<input type="radio"/> <i>Salvia spathacea</i>	California Hummingbird Sage	Lami-	-   -   A1	- NP	Common, oak woodland, chaparral, coastal-sage scrub, slopes (Mar–May)	
<input type="radio"/> <i>Samolus parviflorus</i>	Seaside Brookweed	Theophrast-	-   -   A1	- NP	Moist sites (Spring–summer)	
<input type="radio"/> <i>Sanicula arctopoides</i>	Footsteps-of-spring	Api-	-   -   A1x	- NP	Open coastal bluffs, headlands, dunes (Feb–May)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	-   -   -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	-   -   -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	-   -   -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula laciniata</i>	Fringe-leaf / Coast Sanicle	Api-	-   -   A2	- NP	Coastal, open or shrubby slopes, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula tuberosa</i>	Turkey Pea Sanicle	Api-	-   -   C	- NP	Open gravelly meadows, chaparral, woodland, pine forest (Mar–Jul)	
<input type="radio"/> <i>Scabiosa atropurpurea</i>	Pincushion Flower	Dipsac-	-   -   -	- ZA	Disturbed areas (Mar–Nov)	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	-   -   -	- ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica</i> , <i>floribunda</i> )	California Figwort	Scrophulari-	-   -   -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Scutellaria californica</i>	California Skullcap	Lami-	-   -   A2	- NP	Open sites, scrub, woodland (Jun–Jul)	
<input type="radio"/> <i>Scutellaria siphocampyloides</i>	Grayleaf Skullcap	Lami-	-   -   A1	- NP	Open sites, seeps, dry stream beds, scrub, woodland (May–Jul)	

Herbaceous <i>Scutellaria</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Scutellaria</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberous Skullcap	Lami-	-   -   C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)
<input type="radio"/> <i>Sedella pentandra</i> (JM93: <i>Parvisedum pentandrum</i> )	Annual Stonecrop	Crassul-	-   -   A2	- NA	Compacted ground, slate, shale, sandstone or serpentine outcrops (Mar–Jun)
<input type="radio"/> <i>Sedum spathulifolium</i>	Broadleaf Stonecrop	Crassul-	-   -   B	- NP	Outcrops, often in shade (Apr–Aug)
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	-   -   C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)
<input type="radio"/> <i>Senecio glomeratus</i> (JM93: <i>Erechtites glomerata</i> )	Cutleaf Burnweed	Aster-	-   -   -	M ZA	Disturbed sites, mostly coastal (Apr–Sep)
<input type="radio"/> <i>Senecio hydrophilus</i>	Water Ragwort	Aster-	-   -   A1	- NP	Marshes, swampy places, standing water, alkaline sites (May–Sep)
<input type="radio"/> <i>Senecio minimus</i> (JM93: <i>Erechtites minima</i> )	Coastal Burnweed	Aster-	-   -   -	M ZA	Disturbed coastal sites (Jun–Sep)
<input type="radio"/> <i>Senecio sylvaticus</i>	Woodland Ragwort	Aster-	-   -   -	- ZA	Disturbed woodland, open, sandy sites (Mar–Sep)
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	-   -   -	- ZA	Common. Disturbed areas (Feb–Jul)
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	-   -   -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)
<input type="radio"/> <i>Sidalcea diploscypha</i>	Fringed Checkerbloom	Malv-	-   -   A2	- NA	Grassland, open woodland, valleys, occasionally near vernal pools, generally serpentine (Apr–May)
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. ?	Common Checkerbloom	Malv-	-   -   -	- NP	- -
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>laciniata</i>	Geranium-leaved Checkerbloom	Malv-	-   -   C	- NP	Grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>malviflora</i>	Common Checkerbloom	Malv-	-   -   B	- NP	Coastal prairie, scrub, open forest (Mar–Jul)
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	-   -   -	- ZA	Fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Silene laciniata</i> subsp. <i>californica</i> (JM93: <i>S. californica</i> )	California Pink	Caryophyll-	-   -   B	- NP	Chaparral, oak woodland, conifer forest, serpentine or not (Spring–summer)
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	-   -   -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)
<input type="radio"/> <i>Sinapis arvensis</i>	Charlock	Brassic-	-   -   -	L ZA	Disturbed areas (Mar–Sep)
<input type="radio"/> <i>Sisymbrium irio</i>	London Rocket	Brassic-	-   -   -	M ZA	Disturbed areas, fields, pastures (Jan–Apr)
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	-   -   -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)
<input type="radio"/> <i>Sisymbrium</i> sp.	Rocket / Mustard	Brassic-	-   -   -	- ZA	- -
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	-   -   -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)
<input type="radio"/> <i>Sium suave</i>	Hemlock Waterparsnip	Api-	-   -   A1	- NP	Swamps, marshes, streambanks (Jul–Aug)
<input type="radio"/> <i>Solanum americanum</i>	Small-flowered Nightshade	Solan-	-   -   -	- NAP	Open, often disturbed places (Apr–Nov)
<input type="radio"/> <i>Solanum nigrum</i>	Black Nightshade	Solan-	-   -   -	- ZA	Disturbed places (Mar–Oct)
<input type="radio"/> <i>Solanum</i> sp.	Nightshade	Solan-	-   -   -	-	- -

Herbaceous <b>Solanum</b>	<b>Grouped by Growth Form</b>					Herbaceous <b>Solanum</b>
<b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Solanum xanti</i>	Purple Nightshade	Solan-	-   -   A1?	- NP	Shrubland, oak/pine woodland, conifer forest (Feb–Jun)	
<input type="radio"/> <i>Soleirolia soleirolii</i>	Baby's Tears	Urtic-	-   -   -	- ZA	Damp, shaded places (Mar–Jun)	
<input type="radio"/> <i>Solidago elongata</i> (JM93: <i>S. canadensis</i> subsp. <i>elongata</i> )	West Coast Canada Goldenrod	Aster-	-   -   -	- NP	Meadows, thickets (May–Oct)	
<input type="radio"/> <i>Solidago sp.</i>	Goldenrod	Aster-	-   -   -	- NP	--	
<input type="radio"/> <i>Solidago velutina</i> subsp. <i>californica</i> (JM93: <i>S. californica</i> )	California Goldenrod	Aster-	-   -   -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	-   -   -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	-   -   -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	-   -   -	- ZA	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Spergula arvensis</i> (JM93: subsp. <i>arvensis</i> )	Stickwort	Caryophyll-	-   -   -	- ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Spergularia bocconi</i>	Boccone's Sand-spurry	Caryophyll-	-   -   -	- ZA	Salt marshes, alkaline areas, sandy soils (Spring)	
<input type="radio"/> <i>Spergularia macrotheca</i> var. <i>leucantha</i>	Short-style Sand-spurry	Caryophyll-	-   -   A2	- NP	Alkaline soils, floodplains, vernal pools, meadows, marshy ground (Apr–Jun)	
<input type="radio"/> <i>Spergularia macrotheca</i> var. <i>macrotheca</i>	Sticky Sand-spurry	Caryophyll-	-   -   A2	- NP	Salt flats, marshes, dunes, rocky outcrops, sandy or rocky coastal bluffs, gravelly ridges, alkaline fields (Spring–fall)	
<input type="radio"/> <i>Spergularia marina</i>	Saltmarsh Sand-spurrey	Caryophyll-	-   -   -	- NA	Mud flats, alkaline fields, sandy river bottoms, sandy coasts, salt marshes (Mar–Sep)	
<input type="radio"/> <i>Spergularia media</i> var. <i>media</i> (JM93: no var.)	Greater Sea-spurrey	Caryophyll-	-   -   -	- ZAP	Salt flats, salt marshes, sandy beaches (Summer–fall)	
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	-   -   -	- ZAP	Forest, meadows, mud flats, disturbed	
<input type="radio"/> <i>Spergularia sp.</i>	Sand-spurry	Caryophyll-	-   -   -	-	--	
<input type="radio"/> <i>Spiranthes porrifolia</i>	Western Ladies Tresses	Orchid-	-   -   A1	- NP	Wet meadows, freshwater marshes, seeps, grassland, oak woodland (Jun–Sep)	
<input type="radio"/> <i>Stachys ajugoides</i> (JM93: var. <i>ajugoides</i> )	Bugle Hedge-nettle	Lami-	-   -   A2	- NA	Moist, open places, gen remain wet into summer (Jun–Sep)	
<input type="radio"/> <i>Stachys albens</i>	White-stem Hedge-nettle	Lami-	-   -   B	- NP	Swamps, seeps (May–Oct)	
<input type="radio"/> <i>Stachys bullata</i>	California Hedge-nettle	Lami-	-   -   A1?	- NP	Dry slopes near coast (Mar–Sep)	
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Lami-	-   -   C	- NP	Streambanks, springs, pine/oak forest (Jun–Oct)	
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i> )	Common Rigid Hedge-nettle	Lami-	-   -   -	- NP	Moist to ± dry places (Mar–Oct)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Stebbinsoseris</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Stebbinsoseris</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Stebbinsoseris heterocarpa</i>	Grassland Silverpuffs	Aster-	-   -   B	- NA	Open, sometimes disturbed sites, rocky to clay soils, generally inland (Apr–Jun)	
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	-   -   -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	-   -   C	- NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)	
<input type="radio"/> <i>Stellaria pallida</i>	Lesser Chickweed	Caryophyll-	-   -   -	- ZA	Oak woodland, streambanks, grassy hills, flats, disturbed areas (Spring)	
<input type="radio"/> <i>Stephanomeria elata</i>	Tall Wreath Plant	Aster-	-   -   A2	- NA	Chaparral openings, grassy meadows, roadside embankments (Jul–Nov)	
<input type="radio"/> <i>Stephanomeria virgata subsp. pleurocarpa</i>	Twiggy Wreath Plant	Aster-	-   -   C	- NA	Chaparral openings, grassland (Jun–Nov)	
<input type="radio"/> <i>Streptanthus glandulosus subsp. glandulosus</i> (JM93: incl. <i>S. albidus subsp. peramoenus</i> )	Bristly Jewel Flower	Brassic-	-   -   B	- NA	Serpentine, bare slopes, chaparral & woodland openings (Apr–Jul)	
<input type="radio"/> <i>Stuckenia pectinata</i> (JM93: <i>Potamogeton pectinatus</i> )	Fennel-leaf Pondweed	Potamogeton-	-   -   -	- NP	Common. Ponds, lakes, marshes, streams (May–Jul)	
<input type="radio"/> <i>Symphyotrichum chilense</i> (JM93: <i>Aster chilensis</i> )	Common California Wild Aster	Aster-	-   -   -	- NP	Grassland, salt marshes, disturbed places (Jun–Oct)	
<input type="radio"/> <i>Symphyotrichum lentum</i> (JM93: <i>Aster lentus</i> )	Suisun Marsh Aster	Aster-	-   1B.2   *A2	- NP	Marshes (May–Nov)	
<input type="radio"/> <i>Symphyotrichum subulatum</i> var. <i>parviflorum</i> (JM93: <i>Aster subulatus</i> var. <i>ligulatus</i> )	Annual Saltmarsh Aster	Aster-	-   -   -	- NA	Marshes, disturbed places (Jul–Oct)	
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	-   -   -	- ZBP	Abundant. Esp disturbed areas (All year)	
<input type="radio"/> <i>Taraxia ovata</i> (JM93: <i>Camissonia</i> )	Golden Eggs Suncup	Onagr-	-   -   C	- NP	Grassy fields, gen clay soil (Mar–Jun)	
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	-   -   C	- NP	Chaparral, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Tellima grandiflora</i>	Fringe Cups	Saxifrag-	-   -   B	- NP	Moist slopes (Apr–Jul)	
<input type="radio"/> <i>Tetragonia tetragonioides</i>	New Zealand Spinach	Aizo-	-   -   -	L ZA	Common. Sand dunes, bluffs, margins of coastal wetlands (Apr–Sep)	
<input type="radio"/> <i>Tetrapteron graciliflorum</i> (JM93: <i>Camissonia graciliflora</i> )	Hill Sun Cup	Onagr-	-   -   A2	- NA	Open or shrubby slopes, generally clay soils, grassland, oak and Joshua-tree woodland (Mar–Apr)	
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	-   -   -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	-   -   -	- NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)	
<input type="radio"/> <i>Thysanocarpus laciniatus</i> var. <i>laciniatus</i> (JM93: no var.)	Narrowleaf Fringepod	Brassic-	-   -   B	- NA	Oak woodland, rocky ridges, slopes, chaparral, washes (Mar–May)	
<input type="radio"/> <i>Thysanocarpus radians</i>	Ribbed Fringepod	Brassic-	-   -   A2	- NA	Moist slopes, pastures, open meadows, fields (Mar–Apr)	

Herbaceous <i>Tolmiea</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Tolmiea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Tolmiea diplomenziesii</i> (JM93: <i>T. menziesii</i> )	Pig-a-back Plant	Saxifrag-	-   -   A1	- NP	Moist streambanks (May–Aug)	
<input type="radio"/> <i>Tonella tenella</i>	Small-flowered Tonella	Plantagin-	-   -   A2	- NA	Moist, shaded places in chaparral, oak and mixed woodland (Mar–Jun)	
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	-   -   -	M ZA	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	-   -   -	- ZA	Disturbed places (Apr–Jun)	
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i> )	Common Star Lily	Melanthi-	-   -   -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)	
<input type="radio"/> <i>Tradescantia fluminensis</i>	Spiderwort	Commelin-	-   -   -	- ZP	Shaded woodland, streambanks (Apr–Jul)	
<input type="radio"/> <i>Tragopogon dubius</i>	Yellow Salsify	Aster-	-   -   -	- ZAB	Uncommon. Disturbed places (May–Sep)	
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	-   -   -	- ZBP	Common. Disturbed places (Mar–Nov)	
<input type="radio"/> <i>Trianthema portulacastrum</i>	Horse-purslane	Aizo-	-   -   A1	- NA	Uncommon. Moist or seasonally dry wetlands, disturbed areas (Jun–Nov)	
<input type="radio"/> <i>Tribulus terrestris</i>	Puncture Vine	Zygophyll-	-   -   -	- ZA	Dry, disturbed areas incl roadsides, railways, vacant lots (Apr–Oct)	
<input type="radio"/> <i>Trichostema lanceolatum</i>	Vinegar Weed	Lami-	-   -   -	- NA	Dry, open, gen disturbed habitats (Jun–Nov)	
<input type="radio"/> <i>Trientalis latifolia</i>	Starflower	Myrsin-	-   -   C	- NP	Shaded places, esp woodland (Apr–Jul)	
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i> )	Rancheria Clover	Fab-	-   -   C	- NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)	
<input type="radio"/> <i>Trifolium angustifolium</i>	Narrow-leaved Clover	Fab-	-   -   -	- ZA	Disturbed areas (Late spring)	
<input type="radio"/> <i>Trifolium barbigerum</i> (JM93: var. <i>barbigerum</i> )	Bearded Clover	Fab-	-   -   B	- NP	Wet meadows, open, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Trifolium bifidum</i> var. ?	Pinole Clover	Fab-	-   -   -	- NA	- -	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>bifidum</i>	Pinole Clover	Fab-	-   -   B	- NA	Open, grassy areas, oak chaparral, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Fab-	-   -   -	- NA	Open, grassy areas, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium campestre</i>	Hop Clover	Fab-	-   -   -	- ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	-   -   -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Fab-	-   -   -	- NA	Grassland, coastal woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>depauperatum</i>	Dwarf Sack Clover	Fab-	-   -   B	- NA	Wet meadows, grassland, roadsides, open spring-moist, heavy soils (Mar–May)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Fab-	-   -   -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium dichotomum</i> (JM93: <i>T. albopurpureum</i> var. <i>dichotomum</i> )	Branched Indian Clover	Fab-	-   -   B	- NA	Coastal dunes, open slopes, meadows, oak woodland, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	-   -   -	- ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium fragiferum</i>	Strawberry Clover	Fab-	-   -   -	- ZP	Roadsides, gen in saline soil (Late spring)	

Herbaceous <i>Trifolium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Trifolium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Trifolium fucatum</i>	Bull Clover	Fab-	-   -   C	- NA	Locally abundant. Moist, open grassland, marshes, roadsides, occasionally saline or serpentine soils (Apr–Jun)
<input type="radio"/> <i>Trifolium glomeratum</i>	Clustered Clover	Fab-	-   -   -	- ZA	Uncommon. Disturbed areas (Mar–May)
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i> )	Pinpoint Clover	Fab-	-   -   -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Trifolium grayi</i> (JM93: <i>T. barbigerum</i> var. <i>andrewsii</i> )	Gray's Clover	Fab-	-   -   A1?	- NA	Wet meadows, foothill slopes, pine woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	-   -   -	M ZA	Disturbed areas, roadsides (Apr–May)
<input type="radio"/> <i>Trifolium incarnatum</i>	Crimson Clover	Fab-	-   -   -	- ZA	Uncommon. Disturbed areas (May–Aug)
<input type="radio"/> <i>Trifolium macraei</i>	Macrae's Clover	Fab-	-   -   A2	- NA	Disturbed areas, dunes (Mar–May)
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	-   -   -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	-   -   -	- NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)
<input type="radio"/> <i>Trifolium obtusiflorum</i>	Clammy Clover	Fab-	-   -   B	- NA	Moist disturbed areas, gravel bars, marshes (Apr–Jul)
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Fab-	-   -   C	- NA	Woody or shrubby slopes, roadsides (Mar–Jun)
<input type="radio"/> <i>Trifolium olivaceum</i> (JM93: <i>T. albopurpureum</i> var. <i>olivaceum</i> )	Olive Clover	Fab-	-   -   B	- NA	Grassy slopes, valley meadows, disturbed areas (Apr–May)
<input type="radio"/> <i>Trifolium pratense</i>	Red Clover	Fab-	-   -   -	- ZP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Trifolium repens</i>	White Clover	Fab-	-   -   -	- ZP	Disturbed areas (Feb–Dec)
<input type="radio"/> <i>Trifolium sp.</i>	Clover	Fab-	-   -   -	- -	
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Fab-	-   -   -	- ZA	Meadows, roadsides, disturbed areas (Mar–Apr)
<input type="radio"/> <i>Trifolium variegatum</i> var. ? (JM93: no var.)	White-tip Clover	Fab-	-   -   -	- NA	- -
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	-   -   -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Trifolium wormskioldii</i>	Cow Clover	Fab-	-   -   A2	- NA	Beaches to mtn meadows, ridges, generally open moist or marshy places (May–Oct)
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	-   -   -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)
<input type="radio"/> <i>Trillium ovatum</i> subsp. <i>ovatum</i>	Western White Trillium	Melanthi-	-   -   A2	- NP	Redwood, mixed-evergreen forest on moist wooded slopes (Feb–Apr)
<input type="radio"/> <i>Triodanis biflora</i>	Small Venus-looking-glass	Campanul-	-   -   A2	- NA	Disturbed areas (Apr–Jun)
<input type="radio"/> <i>Triphysaria eriantha</i> subsp. <i>eriantha</i>	Yellow Johnny-tuck	Orobanch-	-   -   -	- NA	Grassland, foothills (Mar–May)
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	-   -   -	- NA	Grassland (Apr–Jun)



Herbaceous <i>Triphysaria</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Triphysaria</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Triphysaria versicolor</i> subsp. <i>faucibarbata</i>	Smooth Owl's Clover	Orobanch-	-   -   A2	- NA	Grassland (Apr–Jun)
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	-   -   -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	-   -   -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)
<input type="radio"/> <i>Tropaeolum majus</i>	Garden Nasturtium	Tropaeol-	-   -   -	- ZAP	Disturbed areas, moist or shaded areas in coastal scrub, wooded flats or slopes (Mar–Jun)
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Brassic-	-   -   B	- NP	Grassy banks, open fields, roadsides, pastures (Mar–May)
<input type="radio"/> <i>Turritis glabra</i> (JM93: <i>Arabis</i> , var. <i>glabra</i> )	Tower Mustard	Brassic-	-   -   C	- NP	Open fields, meadows, slopes (Apr–Jul)
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	-   -   -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)
<input type="radio"/> <i>Urospermum picroides</i>	False Hawkbit	Aster-	-   -   -	- ZAP	Uncommon. Disturbed places (Apr–Jul)
<input type="radio"/> <i>Urtica dioica</i> subsp. <i>holosericea</i>	Hoary Nettle	Urtic-	-   -   -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	-   -   -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)
<input type="radio"/> <i>Verbascum thapsus</i>	Woolly Mullein	Scrophulari-	-   -   -	L ZB	Roadsides, streambanks, disturbed areas (May–Sep)
<input type="radio"/> <i>Verbena bonariensis</i>	Tall Vervain	Verben-	-   -   -	- ZAB	Disturbed, often wet places, fields (Jun–Oct)
<input type="radio"/> <i>Verbena lasiostachys</i> var. ?	Western Vervain	Verben-	-   -   -	- NP	- -
<input type="radio"/> <i>Verbena lasiostachys</i> var. <i>lasiostachys</i>	Western Vervain	Verben-	-   -   -	- NP	Open, dry to wet places (May–Sep)
<input type="radio"/> <i>Verbena lasiostachys</i> var. <i>scabrida</i>	Robust Vervain	Verben-	-   -   -	- NP	Open, dry to wet places (May–Sep)
<input type="radio"/> <i>Veronica americana</i>	American Brooklime	Plantagin-	-   -   C	- NP	Common. Moist to wet soil, springs, slow streams, meadows, lakeshores (May–Aug)
<input type="radio"/> <i>Veronica anagallis-aquatica</i>	Water Speedwell	Plantagin-	-   -   -	- ZP	Wet meadows, streambanks, slow streams (May–Sep)
<input type="radio"/> <i>Veronica arvensis</i>	Common Speedwell	Plantagin-	-   -   -	- ZA	Meadows (Apr–Jul)
<input type="radio"/> <i>Veronica catenata</i>	Chain Speedwell	Plantagin-	-   -   -	- ZP	Wet meadows, slow streams (Jul–Sep)
<input type="radio"/> <i>Veronica peregrina</i> subsp. <i>xalapensis</i>	Purslane Speedwell	Plantagin-	-   -   -	- NA	Moist places (Apr–Aug)
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	-   -   -	- ZA	Wet, disturbed areas, fields (Feb–May)
<input type="radio"/> <i>Veronica</i> sp.	Speedwell	Plantagin-	-   -   -	-	- -
<input type="radio"/> <i>Vicia americana</i> subsp. <i>americana</i> (JM93: var. <i>americana</i> )	American Vetch	Fab-	-   -   -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Vicia benghalensis</i>	One-sided Purple Vetch	Fab-	-   -   -	- ZP	Grassland, roadsides, disturbed areas (Mar–Jun)

Herbaceous <i>Vicia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Vicia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Vicia disperma</i>	Two-seeded Vetch	Fab-	-   -   -	- ZA	Roadsides, urban weed (Mar–May)	
<input type="radio"/> <i>Vicia gigantea</i>	Giant Vetch	Fab-	-   -   C	- NP	Coastal shrub, coastal forest, chaparral (Mar–Aug)	
<input type="radio"/> <i>Vicia hassei</i>	Slender Vetch	Fab-	-   -   A2	- NA	Coastal scrub, chaparral, oak woodland (Mar–May)	
<input type="radio"/> <i>Vicia ludoviciana subsp. ludoviciana (JM93: var. ludoviciana)</i>	Deerpea Vetch	Fab-	-   -   -	- NA	Slopes, canyons, streambanks, in grassland, coastal-sage scrub, chaparral, oak woodland, riparian woodland (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. ?</i>	Narrowleaf / Spring Vetch	Fab-	-   -   -	- ZA	--	
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sp.</i>	Vetch	Fab-	-   -   -	-	--	
<input type="radio"/> <i>Vicia villosa subsp. ?</i>	Vetch	Fab-	-   -   -	- ZAB	--	
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)	
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Apocyn-	-   -   -	M ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))	
<input type="radio"/> <i>Viola adunca var. adunca (JM93: no var.)</i>	Western Dog Violet	Viol-	-   -   A1	- NP	Vernally moist meadows, damp streambanks, meadow edges in conifer forest, generally shade (Apr–Aug)	
<input type="radio"/> <i>Viola douglasii</i>	Douglas' Violet	Viol-	-   -   A1	- NP	Vernally moist flats, grassy slopes, often on serpentine (Feb–Jul)	
<input type="radio"/> <i>Viola glabella</i>	Stream / Smooth Yellow Violet	Viol-	-   -   A2	- NP	Moist to wet generally shady places in forest, streambanks, etc (Mar–Aug)	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	-   -   C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	
<input type="radio"/> <i>Viola purpurea subsp. purpurea</i>	Mountain Violet	Viol-	-   -   A2	- NP	In openings or beneath shrubs, generally in yellow-pine forest or higher (Mar–Jul)	
<input type="radio"/> <i>Viola purpurea subsp. quercetorum</i>	Oak Violet	Viol-	-   -   A2	- NP	Dry, grassy or brushy slopes, chaparral, generally below yellow-pine forest (Feb–Jul)	
<input type="radio"/> <i>Viola sempervirens</i>	Redwood / Evergreen Violet	Viol-	-   -   A1	- NP	Shady areas in coastal forest (Jan–Jul)	
<input type="radio"/> <i>Viola sheltonii</i>	Shelton's Violet	Viol-	-   -   A2	- NP	Fir, pine, or oak woodland, rich or gravelly soil (Mar–Jul)	
<input type="radio"/> <i>Viola sp.</i>	Violet	Viol-	-   -   -	- NP	--	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	-   -   -	- NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	-   -   C	- NP	Gen shady sites (Mar–Jun)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Wyethia</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Wyethia</b>
<input checked="" type="checkbox"/> <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="checkbox"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	-   -   -	-	NP	Open grassland, woodland, scrub (Mar–May(Aug))
<input type="checkbox"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	-   -   -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)
<input type="checkbox"/> <i>Xanthium strumarium</i>	Cocklebur	Aster-	-   -   -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)
<input type="checkbox"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Api-	-   -   C	-	NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)
<input type="checkbox"/> <i>Zannichellia palustris</i>	Horned-pondweed	Zannichelli-	-   -   -	-	NP	Streams, ponds, lakes (Mar–Nov)
<input type="checkbox"/> <i>Zantedeschia aethiopica</i>	Calla-lily	Ar-	-   -   -	L	ZP	Disturbed areas, near former habitations (Mar–Jun)
<input type="checkbox"/> <i>Zeltnera davyi (JM93: Centaurium)</i>	Davy's Centaury	Gentian-	-   -   B	-	NA	Moist coastal bluffs, dunes, open forest (May–Aug)
<input type="checkbox"/> <i>Zeltnera muehlenbergii (JM93: Centaurium)</i>	Monterey Centaury	Gentian-	-   -   -	-	NA	Moist coastal bluffs, forest openings (Jun–Aug)
<input type="checkbox"/> <i>Zeltnera sp. (JM93: Centaurium)</i>	Centaury	Gentian-	-   -   -	-	NA	- -

<b>Woody Acacia</b>	<b>Grouped by Growth Form Sorted Alphabetically by Scientific Name</b>					<b>Woody Acacia</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Acacia cyclops</i>	Western Coastal Wattle	Fab-	-   -   -	- ZP	Uncommon. Disturbed areas, coastal dunes (Dec–Mar)	
<input type="radio"/> <i>Acacia dealbata</i>	Silver Wattle	Fab-	-   -   -	M ZP	Locally common. Disturbed areas, often roadsides (Feb–Apr)	
<input type="radio"/> <i>Acacia longifolia</i>	Sydney Golden Wattle	Fab-	-   -   -	- ZP	Uncommon. Disturbed places, especially sandy coastal areas (Jan–Apr)	
<input type="radio"/> <i>Acacia melanoxylon</i>	Blackwood Acacia	Fab-	-   -   -	L ZP	Uncommon. Disturbed areas (Feb–Mar)	
<input type="radio"/> <i>Acacia sp.</i>	Acacia	Fab-	-   -   -	- ZP	--	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	-   -   -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Acer negundo (JM93: var. californicum)</i>	Box Elder	Sapind-	-   -   -	- NP	Streamsides, bottomland. (Mar–Apr)	
<input type="radio"/> <i>Adenostoma fasciculatum var. fasciculatum (JM93: no var.)</i>	Chamise	Ros-	-   -   -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	-   -   -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Ailanthus altissima</i>	Tree Of Heaven	Simaroub-	-   -   -	M ZP	Disturbed areas, grassland, oak woodland, riparian areas (Jun)	
<input type="radio"/> <i>Allenrolfea occidentalis</i>	Iodine Bush	Chenopodi-	-   -   A2	- NP	Flats, hummocks, in alkaline soils (Jun–Aug)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	-   -   -	- NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Alnus rubra</i>	Red Alder	Betul-	-   -   A2	- NP	Wet places, especially after logging (Feb–Mar)	
<input type="radio"/> <i>Alnus sp.</i>	Alder	Betul-	-   -   -	- NP	--	
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Ros-	-   -   C	- NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	-   -   C	- NP	Conifer, oak forests (Mar–May)	
<input type="radio"/> <i>Arctostaphylos auriculata</i>	Mount Diablo Manzanita	Eric-	-   1B.3   *A2	- NP	Sandstone, upland chaparral near coast. Mount Diablo and vicinity (Feb–Mar)	
<input type="radio"/> <i>Arctostaphylos crustacea subsp. crustacea (JM93: A. tomentosa subsp. crustacea)</i>	Brittle-leaf Manzanita	Eric-	-   -   C	- NP	Chaparral, conifer forest (Feb–Apr)	
<input type="radio"/> <i>Arctostaphylos glandulosa subsp. glandulosa</i>	Eastwood Manzanita	Eric-	-   -   B	- NP	Chaparral, conifer forest (Jan–Apr)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Eric-	-   -   C	- NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Arctostaphylos manzanita subsp. laevigata</i>	Contra Costa Manzanita	Eric-	-   1B.2   *A2	- NP	Chaparral, rocky outcrops (Feb–May)	
<input type="radio"/> <i>Arctostaphylos manzanita subsp. manzanita</i>	Common / Giant Manzanita	Eric-	-   -   C	- NP	Chaparral, conifer forest (Feb–May)	
<input type="radio"/> <i>Arctostaphylos pallida</i>	Pallid Manzanita	Eric-	FT/CE   1B.1   *A1	- NP	Siliceous shales, slopes, ridges, chaparral (Jan–Mar)	
<input type="radio"/> <i>Arctostaphylos sp.</i>	Manzanita	Eric-	-   -   -	- NP	--	
<input type="radio"/> <i>Argyranthemum foeniculaceum</i>	Canary Islands Marguerite	Aster-	-   -   -	- ZP	Escape from cult in disturbed coastal areas (Mar–Aug)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Artemisia</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Artemisia</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	-   -   -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Atriplex lentiformis</i> (JM93: <i>subsp. lentiformis</i> )	Big Saltbush	Chenopodi-	-   -   A2	- NP	Alkaline or saline washes, dry lakes, scrub (Jul–Oct)	
<input type="radio"/> <i>Baccharis pilularis subsp. consanguinea</i> (JM93: <i>no subsp.</i> )	Coyote Brush	Aster-	-   -   -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Baccharis salicifolia subsp. salicifolia</i> (JM93: <i>no subsp.</i> )	Mule Fat	Aster-	-   -   -	- NP	Riparian woodland, canyon bottoms, disturbed sites, often forming thickets (All year)	
<input type="radio"/> <i>Berberis aquifolium var. dictyota</i>	Hollyleaf Oregon Grape	Berberid-	-   -   A2	- NP	Slopes, canyons, conifer forest, oak woodland, chaparral (Mar–May)	
<input type="radio"/> <i>Berberis nervosa</i>	Longleaf Oregon Grape	Berberid-	-   -   A1	- NP	Conifer forest (Mar–Jun)	
<input type="radio"/> <i>Berberis pinnata subsp. pinnata</i>	Shinyleaf Oregon-grape	Berberid-	-   -   C	- NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
<input type="radio"/> <i>Berberis sp.</i>	Oregon Grape	Berberid-	-   -   -	- NP	--	
<input type="radio"/> <i>Brickellia californica</i>	California Brickellbush	Aster-	-   -   B	- NP	Dry, rocky hillsides, canyons, sea bluffs (Jul–Dec)	
<input type="radio"/> <i>Buddleja davidii</i>	Butterfly Bush	Scrophulari-	-   -   -	- ZP	Disturbed, generally ± damp soil (May–Sep)	
<input type="radio"/> <i>Carpobrotus chilensis</i>	Sea Fig	Aizo-	-   -   -	M ZP	Common on coastal sandy shores (All year)	
<input type="radio"/> <i>Carpobrotus edulis</i>	Freeway Iceplant	Aizo-	-   -   -	H ZP	Common. Many coastal habitats, esp. sand (All year)	
<input type="radio"/> <i>Carpobrotus sp.</i>	Hottentot / Sea Fig	Aizo-	-   -   -	- ZP	--	
<input type="radio"/> <i>Ceanothus cuneatus var. cuneatus</i>	Buckbrush	Rhamn-	-   -   C	- NP	Sandy to rocky flats, slopes, ridges (Feb–May)	
<input type="radio"/> <i>Ceanothus oliganthus var. sorediatus</i>	Jim Brush	Rhamn-	-   -   -	- NP	Slopes, ridges, chaparral, conifer forest (Jan–May)	
<input type="radio"/> <i>Ceanothus thyrsiflorus var. thyrsiflorus</i> (JM93: <i>no var.</i> )	Blue Blossom	Rhamn-	-   -   -	- NP	Bluffs, slopes, canyons, chaparral, coastal scrub, closed-cone-pine forest (Mar–Jun)	
<input type="radio"/> <i>Cephalanthus occidentalis</i> (JM93: <i>var. californicus</i> )	California Button Willow	Rubi-	-   -   B	- NP	Lake, stream edges (May–Sep)	
<input type="radio"/> <i>Cercis occidentalis</i>	Western Redbud	Fab-	-   -   A1	- NP	Dry, shrubby slopes, canyons, streambanks, chaparral, foothill woodland, yellow-pine forest (Mar–May)	
<input type="radio"/> <i>Cercocarpus betuloides var. betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	-   -   C	- NP	Dry, rocky slopes, chaparral (Mar–May)	
<input type="radio"/> <i>Chrysolepis chrysophylla var. minor</i>	Golden Chinquapin	Fag-	-   -   A2	- NP	Conifer forest, closed-cone-pine forest, chaparral (Jun–Sep)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	-   -   -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	-   -   C	- NP	Along streams, wet places (Jun–Sep)	
<input type="radio"/> <i>Clematis sp.</i>	Virgin's Bower / Pipestems	Ranuncul-	-   -   -	- NP	--	
<input type="radio"/> <i>Cornus glabrata</i>	Brown Dogwood	Corn-	-   -   A1	- NP	Generally moist places (May–Jun)	

<b>Woody Cornus</b>	<b>Grouped by Growth Form Sorted Alphabetically by Scientific Name</b>						<b>Woody Cornus</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Cornus sericea</i> subsp. ?	American Dogwood	Corn-	-   -   -	-	NP	--	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Corn-	-   -   C	-	NP	Generally moist places (May–Jul)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Corn-	-   -   C	-	NP	Generally moist habitats (May–Jul)	
<input type="radio"/> <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i> )	California Hazelnut	Betul-	-   -   -	-	NP	Common. Many habitats, esp moist, shady places (Jan–Mar)	
<input type="radio"/> <i>Cotoneaster pannosus</i> (JM93: <i>C. pannosa</i> )	Silverleaf Cotoneaster	Ros-	-   -   -	M	ZP	Disturbed places, mixed-evergreen forest (May–Jul)	
<input type="radio"/> <i>Crataegus monogyna</i>	Hawthorn	Ros-	-   -   -	L	ZP	Naturalized in scattered places (May–Aug)	
<input type="radio"/> <i>Cytisus scoparius</i>	Scotch Broom	Fab-	-   -   -	H	ZP	Common. Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Dendromecon rigida</i>	Bush Poppy	Papaver-	-   -   A2	-	NP	Dry slopes, washes, esp recent burns (Apr–Jun)	
<input type="radio"/> <i>Dirca occidentalis</i>	Western Leatherwood	Thymelae-	-   1B.2   *A2	-	NP	Gen n facing slopes, mixed-evergreen forest to chaparral, gen fog belt (Nov–Mar)	
<input type="radio"/> <i>Elaeagnus angustifolia</i>	Russian Olive		-   -   -	M	ZP	Uncommon. Disturbed, sometimes moist places (May–Jun)	
<input type="radio"/> <i>Ericameria arborescens</i>	Golden-fleece	Aster-	-   -   A2	-	NP	Woodland, open forest, chaparral, especially after fire (Aug–Nov)	
<input type="radio"/> <i>Ericameria linearifolia</i>	Interior Goldenbush	Aster-	-   -   -	-	NP	Dry slopes, valleys, foothill and desert woodland, saltbush and creosote-bush scrub (Mar–May)	
<input type="radio"/> <i>Ericameria nauseosa</i> var. <i>mohavensis</i> (JM93: <i>Chrysothamnus nauseosus</i> subsp. <i>mohavensis</i> )	Mojave Rabbitbrush	Aster-	-   -   A2	-	NP	Common. Dry scrub (Aug–Oct)	
<input type="radio"/> <i>Eriodictyon californicum</i>	California Yerba Santa	Boragin-	-   -   -	-	NP	Slopes, fields, roadsides, woodland, chaparral (Apr–Jul)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. ?	California Buckwheat	Polygon-	-   -   -	-	NP	--	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>foliolosum</i>	Leafy California Buckwheat	Polygon-	-   -   A2	-	NP	Gravel (All year)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	Mojave Desert Calif. Buckwheat	Polygon-	-   -   A1	-	NP	Sand, gravel or rocks (All year)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	-   -   -	-	NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Eriophyllum jepsonii</i>	Jepson's Woolly Sunflower	Aster-	-   4.3   *A2	-	NP	Dry oak woodland (Apr–Jun)	
<input type="radio"/> <i>Eriophyllum staechadifolium</i>	Seaside Woolly Sunflower	Aster-	-   -   A2	-	NP	Dunes, sea bluffs, coastal scrub, canyons (Apr–Sep)	
<input type="radio"/> <i>Eucalyptus camaldulensis</i>	River Red Gum	Myrt-	-   -   -	L	ZP	Common. Disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	-   -   -	M	ZP	Common. Disturbed areas (Oct–Jan)	
<input type="radio"/> <i>Eucalyptus</i> sp.	Gum	Myrt-	-   -   -	-	ZP	--	
<input type="radio"/> <i>Eucalyptus tereticornis</i>	Forest Red Gum	Myrt-	-   -   -	-	ZP	Uncommon. Disturbed coastal areas (May–Aug)	
<input type="radio"/> <i>Eucalyptus viminalis</i>	Manna Gum	Myrt-	-   -   -	-	ZP	Uncommon. Disturbed areas (Jul–Sep)	
<input type="radio"/> <i>Euonymus occidentalis</i> var. <i>occidentalis</i>	Western Burning Bush	Celastr-	-   -   A1	-	NP	Shaded streambanks, canyons (Apr–Jun)	

<b>Woody Forestiera</b>	<b>Grouped by Growth Form Sorted Alphabetically by Scientific Name</b>					<b>Woody Forestiera</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Forestiera pubescens</i>	Desert Olive	Ole-	-   -   A2	- NP	Streambanks, canyons, washes (Mar–Apr)	
<input type="radio"/> <i>Frangula californica subsp. californica</i> (JM93: <i>Rhamnus</i> )	California Coffee Berry	Rhamn-	-   -   -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Frangula californica subsp. tomentella</i> (JM93: <i>Rhamnus tomentella subsp. tomentella</i> )	Hoary Coffee Berry	Rhamn-	-   -   B	- NP	Chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Fraxinus dipetala</i>	California Ash	Ole-	-   -   A2	- NP	Canyons, slopes, chaparral, oak/pine woodland (Apr–Jun)	
<input type="radio"/> <i>Fraxinus latifolia</i>	Oregon Ash	Ole-	-   -   B	- NP	Canyons, streambanks, woodland (Mar–May)	
<input type="radio"/> <i>Fraxinus sp.</i>	Ash	Ole-	-   -   -	- NP	-	
<input type="radio"/> <i>Garrya elliptica</i>	Coast Silk Tassel	Garry-	-   -   B	- NP	Seacliffs, sand dunes, chaparral, foothill-pine woodland (Jan–Mar)	
<input type="radio"/> <i>Garrya fremontii</i>	Green-leaf Silk Tassel	Garry-	-   -   A1	- NP	Chaparral, foothill woodland, montane forest (Jan–Apr)	
<input type="radio"/> <i>Gaultheria shallon</i>	Salal	Eric-	-   -   A1	- NP	Moist forest margins (Apr–Jul)	
<input type="radio"/> <i>Genista monspessulana</i>	French Broom	Fab-	-   -   -	H ZP	Common. Disturbed places. (Mar–Jun)	
<input type="radio"/> <i>Grindelia stricta var. angustifolia</i>	Marsh Gumplant	Aster-	-   -   C	- NP	Tidal wetlands (May–Dec)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	-   -   C	- NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Hedera helix</i> (JM93: <i>subsp. helix</i> )	English Ivy	Arali-	-   -   -	H ZP	Woodland, open, disturbed areas (Aug–Nov)	
<input type="radio"/> <i>Helianthemum scoparium</i>	Peak Rush-rose	Cist-	-   -   B	- NP	Dry sandy or rocky soil of hills, slopes, ridges (Feb–Sep)	
<input type="radio"/> <i>Hesperocyparis macrocarpa</i> (JM93: <i>Cupressus</i> )	Monterey Cypress	Cupress-	-   -   -	N NP	Closed-cone-pine/cypress forests -	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	-   -   -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor var. discolor</i> (JM93: <i>no var.</i> )	Oceanspray	Ros-	-   -   -	- NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Ilex aquifolium</i>	English Holly	Aquifoli-	-   -   -	M ZP	Cool, wooded areas (May-Jun)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica var. hindsii</i> )	N. California Black Walnut	Jugland-	-   1B.1   *A2	- NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Juniperus californica</i>	California Juniper	Cupress-	-   -   B	- NP	Dry slopes, flats, pinyon/juniper woodland -	
<input type="radio"/> <i>Keckiella corymbosa</i>	Redwood / Red Rock Penstemon	Plantagin-	-   -   A1	- NP	Rocky slopes in conifer or hardwood forests, (chaparral) (Jun–Oct)	
<input type="radio"/> <i>Lepechinia calycina</i>	Pitcher Sage	Lami-	-   -   B	- NP	Common. Rocky slopes; chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: <i>var. vacillans</i> )	Hairy Vine Honeysuckle	Caprifoli-	-   -   -	- NP	Canyons, streamsides, woodland (May–Jun)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Lonicera</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Lonicera</b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Lonicera interrupta</i>	Chaparral Vine Honeysuckle	Caprifoli-	-   -   -	-	NP	Dry slopes, ridges, floodplains, oak woodland, chaparral (Apr–May)
<input type="radio"/> <i>Lonicera involucrata</i> var. <i>ledebourii</i>	Black Twinberry Honeysuckle	Caprifoli-	-   -   B	-	NP	Moist places (May–Jul)
<input type="radio"/> <i>Lonicera japonica</i>	Japanese Honeysuckle	Caprifoli-	-   -   -	-	ZP	Disturbed places (May–Jul)
<input type="radio"/> <i>Lonicera subspicata</i> var. <i>denudata</i>	Southern Honeysuckle	Caprifoli-	-   -   B	-	NP	Chaparral slopes (Jun–Jul)
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	-   -   -	-	NP	- -
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>albifrons</i>	Bush Lupine	Fab-	-   -   -	-	NP	Common. Chaparral, foothill woodland (Mar–Jun)
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>collinus</i>	Bay Area Silver Lupine	Fab-	-   -   -	-	NP	Cliffs, forest openings (Mar–Jun)
<input type="radio"/> <i>Lupinus arboreus</i>	Yellow Bush Lupine	Fab-	-   -   A2	N	NP	Coastal bluffs, dunes, or more inland (Apr–Jul)
<input type="radio"/> <i>Malacothamnus fasciculatus</i> var. <i>nuttallii</i> (JM93: no var.)	Chaparral Mallow	Malv-	-   -   -	-	NP	Open chaparral (May–Jul)
<input type="radio"/> <i>Malacothamnus fremontii</i>	Fremont Bush-mallow	Malv-	-   -   B	-	NP	Open chaparral to pine woodland (May–Jul)
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	-   -   -	-	NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)
<input type="radio"/> <i>Morella californica</i> (JM93: <i>Myrica</i> )	California Bayberry	Myric-	-   -   A2	-	NP	Coastal dunes and scrub, closed-cone pine, redwood forest (Mar–Apr)
<input type="radio"/> <i>Myoporum laetum</i>	New Zealand Myoporum	Scrophulari-	-   -   -	M	ZP	Open areas in grassland, scrub, riparian habitats, generally coastal (Early spring)
<input type="radio"/> <i>Nicotiana glauca</i>	Tree Tobacco	Solan-	-   -   -	M	ZP	Open, disturbed flats or slopes (Apr–Aug)
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	-   -   -	-	NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)
<input type="radio"/> <i>Olea europaea</i>	Olive	Ole-	-   -   -	L	ZP	Generally waif, persisting from cultivation (Feb–Jun)
<input type="radio"/> <i>Parthenocissus vitacea</i>	Woodbine	Vit-	-   -   -	-	NP	Uncommon. Hillsides, thickets, ravines, open woodland, roadsides (Apr–Jun)
<input type="radio"/> <i>Phoenix canariensis</i>	Canary Island Date Palm	Arec-	-   -   -	L	ZP	Uncommon. Near development, disturbed areas (Oct–Apr)
<input type="radio"/> <i>Physocarpus capitatus</i>	Pacific Ninebark	Ros-	-   -   C	-	NP	Moist banks, n-facing slopes, mixed-conifer forest (May–Jul)
<input type="radio"/> <i>Pickeringia montana</i> var. <i>montana</i>	Chaparral Pea	Fab-	-   -   B	-	NP	Chaparral, open woodland (May–Aug)
<input type="radio"/> <i>Pinus attenuata</i>	Knobcone Pine	Pin-	-   -   A2	-	NP	Closed-cone-pine forest, chaparral -
<input type="radio"/> <i>Pinus coulteri</i>	Coulter Pine	Pin-	-   -   A2	-	NP	Chaparral, lower mixed-conifer, mixed-hardwood forests -
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	-   -   -	-	NP	Closed-cone-pine forest, oak woodland -



<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b><i>Pinus</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Pinus</i></b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Pinus sabiniana</i>	Gray Pine	Pin-	-   -   -	- NP	Foothill woodland, n oak woodland, chaparral, infertile soils in mixed-conifer and hardwood forests -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	-   -   B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Populus fremontii</i> subsp. <i>fremontii</i>	Fremont Cottonwood	Salic-	-   -   -	- NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)	
<input type="radio"/> <i>Populus trichocarpa</i> (JM93: <i>P. balsamifera</i> subsp. <i>trichocarpa</i> )	Black Cottonwood	Salic-	-   -   C	- NP	Alluvial bottomland, streamsides (Feb–Apr)	
<input type="radio"/> <i>Prunus cerasifera</i>	Cherry Plum	Ros-	-   -   -	L ZP	Roadsides, streambanks, chaparral as waif (Feb–Mar)	
<input type="radio"/> <i>Prunus dulcis</i>	Almond	Ros-	-   -   -	- ZP	Canyons, roadsides, grassland (as waif) (Feb–Mar)	
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Ros-	-   -   C	- NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Prunus ilicifolia</i> subsp. <i>ilicifolia</i>	Holly-leaved Cherry	Ros-	-   -   C	- NP	Canyons, slopes, scrubland, woodland (Apr–May)	
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Ros-	-   -   C	- NP	Mixed-evergreen or conifer forest (Mar–May)	
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Ros-	-   -   -	- NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Ptelea crenulata</i>	Western Hop Tree	Rut-	-   -   B	- NP	Scrub, woodland (Apr–May)	
<input type="radio"/> <i>Pyracantha angustifolia</i>	Woolly Firethorn	Ros-	-   -   -	L ZP	Disturbed areas, fencerows, abandoned fields, roadsides (Feb–Jun)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	-   -   A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	-   -   C	- NP	Dry slopes, chaparral (Feb–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	-   -   -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	-   -   -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	
<input type="radio"/> <i>Quercus durata</i> var. <i>durata</i>	Leather Oak	Fag-	-   -   A2	- NP	Chaparral, generally serpentine (Apr–May)	
<input type="radio"/> <i>Quercus garryana</i> var. <i>garryana</i>	Oregon / Garry Oak	Fag-	-   -   B	- NP	Slopes, mixed-evergreen or conifer forest (Apr–Jun)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	-   -   -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	-   -   B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus palmeri</i>	Palmer Oak	Fag-	-   -   A2	- NP	Uncommon. Rocky slopes, flats (Apr–May)	
<input type="radio"/> <i>Quercus</i> sp.	Oak	Fag-	-   -   -	- NP	--	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>frutescens</i>	Dwarf Interior Live Oak	Fag-	-   -   C	- NP	Valleys, chaparral (Mar–May)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Quercus</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Quercus</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	-   -   -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Rhamn-	-   -   C	- NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	-   -   C	- NP	Chaparral, montane forest (Mar–Jun)	
<input type="radio"/> <i>Rhus aromatica</i> (JM93: <i>R. trilobata</i> )	Skunk Bush	Anacardi-	-   -   C	- NP	Slopes, washes, scrub (Mar–May)	
<input type="radio"/> <i>Ribes amarum</i>	Bitter Gooseberry	Grossulari-	-   -   A2	- NP	Chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	-   -   -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes divaricatum</i> var. <i>pubiflorum</i>	Straggle Gooseberry	Grossulari-	-   -   B	- NP	Uncommon. Coastal bluffs, forest edges (Mar–May)	
<input type="radio"/> <i>Ribes malvaceum</i> var. <i>malvaceum</i>	Chaparral Currant	Grossulari-	-   -   B	- NP	Chaparral, oak woodland (Oct–Apr)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	-   -   -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes sanguineum</i> var. <i>glutinosum</i>	Red-flowering Currant	Grossulari-	-   -   C	- NP	Many habitats (Feb–Apr)	
<input type="radio"/> <i>Ribes</i> sp.	Gooseberry / Currant	Grossulari-	-   -   -	- NP	- -	
<input type="radio"/> <i>Ribes speciosum</i>	Fuchsia-flowered Gooseberry	Grossulari-	-   -   A1	- NP	Coastal-sage scrub, chaparral (Jan–May)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Fab-	-   -   -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	-   -   -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	-   -   -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa</i> sp.	Rose	Ros-	-   -   -	- NP	- -	
<input type="radio"/> <i>Rosa spithamea</i>	Coast Ground Rose	Ros-	-   -   C	- NP	Open forest, chaparral, especially after fire (Apr–Aug)	
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i> )	Himalayan Blackberry	Ros-	-   -   -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	-   -   -	- NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="radio"/> <i>Rubus spectabilis</i>	Salmonberry	Ros-	-   -   A1	- NP	Moist areas, especially edges of woodland, streambanks (Mar–Jun)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	-   -   -	- NP	Open, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Salic-	-   -   -	- NP	Common. Floodplains, sandy gravel (Apr–May)	
<input type="radio"/> <i>Salix gooddingii</i>	Goodding's Black Willow	Salic-	-   -   B	- NP	Common. Streamsides, marshes, seepage areas, washes, meadows (Mar–Apr)	
<input type="radio"/> <i>Salix laevigata</i>	Red Willow	Salic-	-   -   -	- NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)	
<input type="radio"/> <i>Salix lasiandra</i> var. <i>lasiandra</i> (JM93: <i>S. lucida</i> subsp. <i>Lasiandra</i> )	Pacific Willow	Salic-	-   -   -	- NP	Wet meadows, lakeshores, riverbanks (May–Jun)	

<b>Woody</b>	<b>Grouped by Growth Form</b>						<b>Woody</b>
<b>Salix</b>	<b>Sorted Alphabetically by Scientific Name</b>						<b>Salix</b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	-   -   -	-	NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="radio"/> <i>Salix scouleriana</i>	Scouler Willow	Salic-	-   -   A2	-	NP	Common. Dry to moist forest, meadows, springs, swamps (Feb–Jun)	
<input type="radio"/> <i>Salix sitchensis</i>	Sitka Willow	Salic-	-   -   -	-	NP	Common. Tidal swamps, marshes, springs, streambeds (Mar)	
<input type="radio"/> <i>Salix sp.</i>	Willow	Salic-	-   -   -	-	?P	- -	
<input type="radio"/> <i>Salvia mellifera</i>	Black Sage	Lami-	-   -   -	-	NP	Coastal-sage scrub, lower chaparral (Mar–Jun)	
<input type="radio"/> <i>Sambucus nigra subsp. caerulea (JM93: S. mexicana)</i>	Blue Elderberry	Adox-	-   -   -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="radio"/> <i>Sambucus racemosa var. racemosa</i>	Red Elderberry	Adox-	-   -   A1	-	NP	Moist places (May–Jul)	
<input type="radio"/> <i>Schinus molle</i>	Pepper Tree	Anacardi-	-   -   -	L	ZP	Washes, slopes, abandoned fields (Jun–Aug)	
<input type="radio"/> <i>Senecio flaccidus var. douglasii</i>	Douglas' Threadleaf Ragwort	Aster-	-   -   A2	-	NP	Generally disturbed dry, open, sandy or rocky sites (Jun–Oct)	
<input type="radio"/> <i>Sequoia sempervirens</i>	Redwood	Cupress-	-   -   B	-	NP	Redwood forest -	
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	-   -   -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="radio"/> <i>Spartium junceum</i>	Spanish Broom	Fab-	-   -   -	H	ZP	Disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Suaeda sp.</i>	Seepweed / Seablite	Chenopodi-	-   -   -	-	NP	- -	
<input type="radio"/> <i>Symphoricarpos albus var. laevigatus</i>	Snowberry	Caprifoli-	-   -   -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="radio"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	-   -   -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="radio"/> <i>Tamarix aphylla</i>	Athel	Tamaric-	-   -   -	L	ZP	Uncommon. Washes, roadsides (May–Nov)	
<input type="radio"/> <i>Tamarix parviflora</i>	Smallflower Tamarisk	Tamaric-	-   -   -	H	ZP	Common. Washes, streambanks, slopes, roadsides (Mar–Apr)	
<input type="radio"/> <i>Tamarix ramosissima</i>	Saltcedar	Tamaric-	-   -   -	H	ZP	Common. Washes, streambanks (Apr–Aug)	
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	-   -   -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="radio"/> <i>Ulmus parvifolia</i>	Chinese Elm	Ulm-	-   -   -	-	ZP	Streams, springs, wetlands, roadsides, disturbed areas (Aug–Oct)	
<input type="radio"/> <i>Umbellularia californica</i>	California Bay	Laur-	-   -   -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)	
<input type="radio"/> <i>Vaccinium ovatum</i>	California Huckleberry	Eric-	-   -   B	-	NP	Edges, clearings in conifer forest (Mar–May)	
<input type="radio"/> <i>Viburnum ellipticum</i>	Oval-leaved Viburnum	Adox-	-   2.3   *A2	-	NP	Chaparral, yellow-pine forest, generally n-facing slopes (Jun–Aug)	
<input type="radio"/> <i>Vitis californica</i>	California Wild Grape	Vit-	-   -   -	-	NP	Streamsides, springs, canyons (May–Jun)	
<input type="radio"/> <i>Vitis vinifera</i>	Wine Grape	Vit-	-   -   -	-	ZP	Abandoned fields, roadsides (May–Jun)	

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Acer negundo* var. *californicum* (no var.)

*Adenostoma fasciculatum* (var. *fasciculatum*)

*Agoseris grandiflora* (var. *grandiflora*)

*Agoseris heterophylla* (var. *cryptopleura*)

*Agrostis viridis* (*Polypogon*)

*Alisma plantago-aquatica* (*A. triviale*)

*Amsinckia menziesii* var. *intermedia* (*A. intermedia*)

*Amsinckia menziesii* var. *menziesii* (no var.)

*Antirrhinum vexillo-calyculatum* ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

*Arabis breweri* var. *breweri* (*Boechera breweri* subsp. *breweri*)

*Arabis glabra* var. *glabra* (*Turritis glabra*)

*Arctostaphylos tomentosa* ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

*Aster chilensis* (*Symphyotrichum chilense*)

*Aster lentus* (*Symphyotrichum lentum*)

*Aster radulinus* (*Eurybia radulina*)

*Aster subulatus* var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

*Atriplex joaquiniana* (*A. joaquinana*)

*Atriplex lentiformis* ssp. *lentiformis* (no subsp.)

*Atriplex triangularis* (*A. prostrata*)

*Azolla mexicana* (*A. microphylla*)

*Baccharis douglasii* (*B. glutinosa*)

*Baccharis pilularis* (subsp. *consanguinea*)

*Baccharis salicifolia* (subsp. *salicifolia*)

*Balsamorhiza macrolepis* var. *macrolepis* (no var.)

*Blepharizonia plumosa* ssp. *plumosa* (no subsp.)

*Blepharizonia plumosa* ssp. *viscida* (*Blepharizonia laxa*)

*Boschniakia strobilacea* (*Kopsiopsis*)

*Bromus catharticus* (var. *catharticus*)

*Camissonia boothii* ssp. *decorticans* (*Eremothera*)

*Camissonia graciliflora* (*Tetrapteron graciliflorum*)

*Camissonia intermedia* (*Camissoniopsis*)

*Camissonia micrantha* (*Camissoniopsis*)

*Camissonia ovata* (*Taraxia*)

*Cardaria draba* (*Lepidium*)

*Carduus pycnocephalus* (subsp. *pycnocephalus*)

*Carex deweyana* ssp. *leptopoda* (*C. leptopoda*)

*Carex lanuginosa* (*C. pellita*)

*Carex ovalis* (*C. leporina*)

*Ceanothus thrysiflorus* (var. *thrysiflorus*)

*Centaurium davyi* (*Zeltnera*)

*Centaurium muehlenbergii* (*Zeltnera*)

*Cephalanthus occidentalis* var. *californicus* (no var.)

*Chamomilla suaveolens* (*Matricaria discoidea*)

*Chenopodium ambrosioides* (*Dysphania*)

*Chenopodium macrospermum* var. *halophilum* (no var.)

*Chenopodium multifidum* (*Dysphania multifida*)

*Chenopodium rubrum* (var. *rubrum*)

*Chrysothamnus nauseosus* ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

*Cirsium cymosum* (var. *cymosum*)

*Claytonia parviflora* ssp. *parviflora* (subsp. *utahensis*)

*Collinsia bartsifolia* var. *davidsonii* (var. *stricta*)

*Collinsia heterophylla* (var. *heterophylla*)

*Conyza bilbaoana* (*Erigeron sumatrensis*)

*Conyza bonariensis* (*Erigeron*)

*Conyza canadensis* (*Erigeron*)

*Conyza coulteri* (*Laennecia*)

*Corallorhiza maculata* (var. *maculata*)

*Cordylanthus mollis* ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

*Coronopus didymus* (*Lepidium didymum*)

*Corylus cornuta* var. *californica* (subsp. *californica*)

*Cotoneaster pannosa* (*C. pannosus*)

*Cryptantha decipiens* (*C. rattanii*)

*Cryptantha intermedia* (var. *intermedia*)

*Cryptantha muricata* (var. *muricata*)

*Cryptantha torreyana* (var. *pumila*)

*Cupressus macrocarpa* (*Hesperocyparis*)

*Cuscuta salina* var. *major* (*C. pacifica* var. *pacifica*)

*Cynara cardunculus* (subsp. *flavescens*)

*Danthonia californica* var. *californica* (no var.)

*Dicentra chrysantha* (*Ehrendorferia*)

*Disporum hookeri* (*Prosartes*)

*Disporum smithii* (*Prosartes*)

*Draba cuneifolia* var. *integrifolia* (no var.)

*Duchesnea indica* (no var.)

*Epilobium pygmaeum* (*E. campestre*)

*Eragrostis curvula* var. *curvula* (no var.)

*Erechtites glomerata* (*Senecio glomeratus*)

*Erechtites minima* (*Senecio minimus*)

*Eremocarpus setigerus* (*Croton*)

*Erigeron philadelphicus* (var. *philadelphicus*)

*Eriogonum umbellatum* var. *bahiiforme* (var. *smallianum*)

*Eriophyllum lanatum* var. *achillaeoides* (var. *achilleoides*)

*Eryngium aristulatum* var. *aristulatum* (*E. jepsonii*)

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

*Erysimum capitatum* var. *angustatum* (var. *capitatum*)

*Filago californica* (*Logfia filaginoides*)

*Filago gallica* (*Logfia*)

*Fritillaria affinis* var. *affinis* (no var.)

*Galium trifidum* var. *pacificum* (subsp. *columbianum*)

*Gastridium ventricosum* (*G. phleoides*)

*Gnaphalium bicolor* (*Pseudognaphalium biolettii*)

*Gnaphalium californicum* (*Pseudognaphalium*)

*Gnaphalium canescens* ssp. *beneolens*  
(*Pseudognaphalium beneolens*)

*Gnaphalium canescens* ssp. *microcephalum*  
(*Pseudognaphalium microcephalum*)

*Gnaphalium luteo-album* (*Pseudognaphalium luteoalbum*)

*Gnaphalium purpureum* (*Gamochaeta ustulata*)

*Gnaphalium ramosissimum* (*Pseudognaphalium*)

*Gnaphalium stramineum* (*Pseudognaphalium*)

*Grindelia camporum* var. *camporum* (no var.)

*Grindelia hirsutula* var. *davyi* (*G. camporum*)

*Grindelia hirsutula* var. *hirsutula* (no var.)

*Guillenia flavescens* (*Caulanthus*)

*Guillenia lasiophylla* (*Caulanthus lasiophyllus*)

*Hedera helix* ssp. *helix* (no subsp.)

*Heliotropium curassavicum* (var. *oculatum*)

*Hemizonia congesta* ssp. *congesta* (subsp. *lutescens*)

*Hemizonia corymbosa* ssp. *corymbosa* (*Deinandra corymbosa*)

*Hemizonia fitchii* (*Centromadia*)

*Hemizonia lobbii* (*Deinandra*)

*Hemizonia parryi* ssp. *congdonii* (*Centromadia*)

*Hemizonia parryi* ssp. *parryi* (*Centromadia*)

*Hemizonia pungens* ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

*Hemizonia pungens* ssp. *pungens* (*Centromadia*)

*Heracleum lanatum* (*H. maximum*)

*Herniaria hirsuta* ssp. *cinerea* (var. *cinerea*)

*Holodiscus discolor* (var. *discolor*)

*Hordeum jubatum* (subsp. *jubatum*)

*Horkelia californica* ssp. *californica* (var. *californica*)

*Horkelia californica* ssp. *dissita* (var. *elata*)

*Horkelia californica* ssp. *frondosa* (var. *frondosa*)

*Isopyrum occidentale* (*Enemion*)

*Iva axillaris* ssp. *robustior* (no subsp.)

*Juglans californica* var. *hindsii* (*J. hindsii*)

*Juncus balticus* (subsp. *ater*)

*Juncus effusus* var. *pacificus* (subsp. *pacificus*)

*Juncus lesueurii* (*J. lescurii*)

*Koeleria phleoides* (*K. gerardii*)

*Lagophylla ramosissima* ssp. *ramosissima* (no subsp.)

*Lasthenia californica* (subsp. *californica*)

*Lavatera cretica* (*Malva pseudolavatera*)

*Lepidium dictyotum* var. *dictyotum* (no var.)

*Lepidium latipes* var. *latipes* (no var.)

*Lepidium nitidum* var. *nitidum* (no var.)

*Lepidium nitidum* var. *oreganum* (no var.)

*Leptochloa fascicularis* (*L. fusca* subsp. *fascicularis*)

*Lessingia filaginifolia* var. *californica* (*Corethrogyne filaginifolia*)

*Lessingia filaginifolia* var. *filaginifolia*  
(*Corethrogyne filaginifolia*)

*Lewisia rediviva* (var. *rediviva*)

*Leymus condensatus* (*Elymus*)

*Leymus triticoides* (*Elymus*)

*Lilaea scilloides* (*Triglochin*)

*Limosella subulata* (*Limosella australis*)

*Linanthus acicularis* (*Leptosiphon*)

*Linanthus ambiguus* (*Leptosiphon*)

*Linanthus androsaceus* (*Leptosiphon*)

*Linanthus bicolor* (*Leptosiphon*)

*Linanthus ciliatus* (*Leptosiphon*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus parviflorus* (*Leptosiphon*)

*Linaria canadensis* (*Nuttallanthus texanus*)

*Lolium multiflorum* (*Festuca perennis*)

*Lolium perenne* (*Festuca perennis*)

*Lolium temulentum* (*Festuca temulenta*)

*Lonicera hispidula* var. *vacillans* (no var.)

*Lotus humistratus* (*Acmispon brachycarpus*)

*Lotus micranthus* (*Acmispon parviflorus*)

*Lotus purshianus* var. *purshianus* (*Acmispon americanus* var. *americanus*)

*Lotus scoparius* var. *scoparius* (*Acmispon glaber* var. *glaber*)

*Lotus strigosus* (*Acmispon*)

*Lotus wrangelianus* (*Acmispon*)

*Luzula comosa* (var. *comosa*)

*Madia madioides* (*Anisocarpus*)

*Malacothamnus fasciculatus* (var. *nuttallii*)

*Marah fabaceus* (*M. fabacea*)

*Marah oreganus* (*M. oregana*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Melilotus alba* (*M. albus*)

*Melilotus indica* (*M. indicus*)

*Mentha spicata* var. *spicata* (no var.)

*Mimulus aurantiacus* (var. *aurantiacus*)

*Mirabilis californica* (*M. laevis* var. *crassifolia*)

*Monardella antonina* ssp. *antonina* (*M. villosa* subsp. *villosa*)

*Monardella douglasii* ssp. *douglasii* (no subsp.)

*Monardella villosa* ssp. *globosa* (subsp. *villosa*)

*Montia fontana* ssp. *chondrosperma* (no subsp.)

*Myrica californica* (*Morella*)

*Nassella cernua* (*Stipa*)

*Nassella lepida* (*Stipa*)

*Nassella pulchra* (*Stipa*)

*Osmorhiza chilensis* (*O. berteroi*)

*Oxalis albicans* ssp. *pilosa* (*O. pilosa*)

*Parvisedum pentandrum* (*Sedella pentandra*)

*Phacelia ramosissima* var. *latifolia* (no var.)

*Phacelia ramosissima* var. *ramosissima* (no var.)

*Phlox gracilis* (*Microsteris*)

*Phoradendron macrophyllum* (*P. serotinum* subsp. *macrophyllum*)

*Phoradendron villosum* (*P. serotinum* subsp. *tomentosum*)

*Phyla nodiflora* var. *incisa* (no var.)

*Phyla nodiflora* var. *nodiflora* (no var.)

*Picris echioides* (*Helminthotheca*)

*Piperia elegans* (subsp. *elegans*)

*Piptatherum miliaceum* (*Stipa miliacea* var. *miliacea*)

*Plagiobothrys canescens* (var. *canescens*)

*Plagiobothrys fulvus* (var. *campestris*)

*Plectritis brachystemon* (*P. congesta* subsp. *brachystemon*)

*Plectritis ciliosa* ssp. *ciliosa* (no subsp.)

*Plectritis ciliosa* ssp. *insignis* (no subsp.)

*Plectritis congesta* (subsp. *congesta*)

*Pleuropogon californicus* (var. *californicus*)

*Pluchea odorata* (var. *odorata*)

*Poa bulbosa* (subsp. *vivipara*)

*Polygonum amphibium* var. *emersum* (*Persicaria* *amphibia*)

*Polygonum amphibium* var. *stipulaceum* (*Persicaria* *amphibia*)

*Polygonum arenastrum* (*P. aviculare* subsp. *depressum*)

*Polygonum hydropiperoides* (*Persicaria*)

*Polygonum lapathifolium* (*Persicaria lapathifolia*)

*Polygonum persicaria* (*Persicaria maculosa*)

*Polygonum punctatum* (*Persicaria punctata*)

*Populus balsamifera* ssp. *trichocarpa* (*P. trichocarpa*)

*Potamogeton pectinatus* (*Stuckenia pectinata*)

*Potentilla glandulosa* ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

*Psilocarphus tenellus* var. *tenellus* (no var.)

*Ranunculus aquatilis* var. *capillaceus* (var. *diffusus*)

*Ranunculus californicus* (var. *californicus*)

*Ranunculus canus* (var. *canus*)

*Ranunculus occidentalis* (var. *occidentalis*)

*Ranunculus sceleratus* (var. *sceleratus*)

*Rhamnus californica* ssp. *californica* (*Frangula*)

*Rhamnus tomentella* ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

*Rhus trilobata* (*R. aromatica*)

*Ribes menziesii* (var. *menziesii*)

*Rorippa nasturtium-aquaticum* (*Nasturtium officinale*)

*Rorippa palustris* var. *occidentalis* (subsp. *palustris*)

*Rosa gymnocarpa* (var. *gymnocarpa*)

*Rubus discolor* (*R. armeniacus*)

*Rumex maritimus* (*R. fueginus*)

*Rumex salicifolius* var. *crassus* (*R. crassus*)

*Rumex salicifolius* var. *salicifolius* (no var.)

*Rumex salicifolius* var. *transitorius* (*R. transitorius*)

*Salicornia europaea* (*S. depressa*)

*Salicornia subterminalis* (*Arthrocnemum subterminale*)

*Salicornia virginica* (*S. pacifica*)

*Salix exigua* (var. *hindsiana*)

*Salix lucida* ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

*Sambucus mexicana* (*S. nigra* subsp. *caerulea*)

*Satureja douglasii* (*Clinopodium*)

*Saxifraga californica* (*Micranthes*)

*Scirpus acutus* var. *occidentalis* (*Schoenoplectus*)

*Scirpus americanus* (*Schoenoplectus*)

*Scirpus californicus* (*Schoenoplectus*)

*Scirpus cernuus* (*Isolepis*)

*Scirpus maritimus* (*Bolboschoenus maritimus* subsp. *paludosus*)

*Scirpus robustus* (*Bolboschoenus*)

*Scrophularia californica* ssp. *californica* (no subsp.)

*Scrophularia californica* ssp. *floribunda* (no subsp.)

*Senecio breweri* (*Packera*)

*Senecio mikanioides* (*Delairea odorata*)

*Silene californica* (*S. laciniata* subsp. *californica*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Smilacina racemosa* (*Maianthemum racemosum*)

*Smilacina stellata* (*Maianthemum stellatum*)

*Solidago californica* (*S. velutina* subsp. *californica*)

*Solidago canadensis* ssp. *elongata* (*S. elongata*)

*Sparganium erectum* ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

*Sparganium eurycarpum* ssp. *eurycarpum* (var. *eurycarpum*)

*Spergula arvensis* ssp. *arvensis* (no subsp.)

*Spergularia media* ( var. *media*)

*Stachys ajugoides* var. *ajugoides* (no var.)

*Stachys ajugoides* var. *rigida* (*Stachys rigida* var. *quercetorum*)

*Streptanthus albidus* ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

*Stylomecon heterophylla* (*Papaver heterophyllum*)

*Taeniatherum caput-medusae* (*Elymus*)

*Thysanocarpus laciniatus* (var. *laciniatus*)

*Tolmiea menziesii* (*T. diplomenziesii*)

*Trifolium albopurpureum* var. *albopurpureum* (no var.)

*Trifolium albopurpureum* var. *dichotomum* (*T. dichotomum*)

*Trifolium albopurpureum* var. *olivaceum* (*T. olivaceum*)

*Trifolium barbigerum* var. *andrewsii* (*T. grayi*)

*Trifolium barbigerum* var. *barbigerum* (no var.)

*Trifolium gracilentum* var. *gracilentum* (no var.)

*Vicia americana* var. *americana* (subsp. *americana*)

*Vicia ludoviciana* var. *ludoviciana* (subsp. *ludoviciana*)

*Viola adunca* (var. *adunca*)

*Vulpia bromoides* (*Festuca*)

*Vulpia microstachys* var. *ciliata* (*Festuca microstachys*)

*Vulpia microstachys* var. *confusa* (*Festuca microstachys*)

*Vulpia microstachys* var. *microstachys* (*Festuca microstachys*)

*Vulpia microstachys* var. *pauciflora* (*Festuca microstachys*)

*Vulpia myuros* var. *hirsuta* (*Festuca myuros*)

*Vulpia myuros* var. *myuros* (*Festuca myuros*)

*Vulpia octoflora* var. *hirtella* (*Festuca octoflora*)

*Vulpia octoflora* var. *octoflora* (*Festuca octoflora*)

*Zigadenus fremontii* (*Toxicoscordion*)