

Anthony Chabot Regional Park

Checklist of Wild Plants

Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Anthony Chabot Regional Park. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Family	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

Anthony Chabot Regional Park

Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form				Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name				<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Adiantum aleuticum</i>	Five-finger Fern	Pterid-	- - A2	- NP	Shady, moist banks, streamsides, serpentine -
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	- - -	- NP	Shaded hillsides, moist woodland -
<input type="radio"/> <i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	Western Lady Fern	Woods-	- - -	- NP	Woodland, along streams, seepage area -
<input type="radio"/> <i>Azolla filiculoides</i>	Mosquito Fern	Azoll-	- - -	- NP	Common. Ponds, slow streams -
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	- - -	- NP	Locally common. Open, wooded slopes, caves -
<input type="radio"/> <i>Equisetum arvense</i>	Common Horsetail	Equiset-	- - -	- NP	Streambanks, wet meadows, springs, other wet, shaded places -
<input type="radio"/> <i>Equisetum hyemale</i> subsp. <i>affine</i>	Common Scouring Rush	Equiset-	- - -	- NP	Streams, moist, sandy, gravelly areas -
<input type="radio"/> <i>Equisetum</i> sp.	Scouring Rush	Equiset-	- - -	- NP	- -
<input type="radio"/> <i>Equisetum telmateia</i> subsp. <i>braunii</i>	Giant Horsetail	Equiset-	- - -	- NP	Streambanks, roadside ditches, seepage areas -
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	- - -	- NP	Generally rocky or dry areas -
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Pterid-	- - C	- NP	Rocky or dry areas -
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Pterid-	- - -	- NP	Gen shaded, sometimes rocky or wooded areas -
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	- - -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Dryopterid-	- - -	- NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -
<input type="radio"/> <i>Pteridium aquilinum</i> var. <i>pubescens</i>	Bracken Fern	Dennstaedti-	- - -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	- - -	- NP	Near streams, springs, seeps -

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like	
<i>Agrostis</i>	Sorted Alphabetically by Scientific Name				<i>Agrostis</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis avenacea</i>	Pacific Bent Grass	Po-	- - -	L	ZP	Open, often disturbed place (Jun-Jul)
<input type="radio"/> <i>Agrostis capillaris</i>	Colonial Bent Grass	Po-	- - -	-	ZP	Roadsides, open, disturbed places (Jul-Sep)
<input type="radio"/> <i>Agrostis exarata</i>	Spike Bent Grass	Po-	- - -	-	NP	Common. Moist or disturbed areas, open woodland, conifer forest (Jun-Aug)
<input type="radio"/> <i>Agrostis hallii</i>	Hall's Bent Grass	Po-	- - A2	-	NP	Open oak woodland, conifer forest (May-Jul)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	- - -	-	NP	Common. Open meadows, woodland, forest, subalpine (Jun-Aug)
<input type="radio"/> <i>Agrostis stolonifera</i>	Creeping Bent Grass	Po-	- - -	L	ZP	Ditches, lake margins, marshes (Jun-Sep)
<input type="radio"/> <i>Aira caryophyllea</i>	Silver Hair Grass	Po-	- - -	-	ZA	Sandy soils, open or disturbed sites (Apr-Jun)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	- - -	M	ZA	Disturbed sites (Mar-Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	- - -	M	ZA	Disturbed sites (Apr-Jun)
<input type="radio"/> <i>Brachypodium distachyon</i>	Purple False-brome	Po-	- - -	M	ZA	Disturbed areas, dry slopes (Apr-Jul)
<input type="radio"/> <i>Briza maxima</i>	Rattlesnake Grass	Po-	- - -	L	ZA	Shaded sites, roadsides, pastures, weedy on coastal dunes (Apr-Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	- - -	-	ZA	Shaded or moist, open sites (Apr-Jul)
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	- - -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)
<input type="radio"/> <i>Bromus catharticus</i> var. <i>catharticus</i> (JM93: no var.)	Rescue Grass	Po-	- - -	-	ZAP	Open, disturbed places (Apr-Nov)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	- - -	M	ZA	Open, gen disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	- - -	L	ZA	Fields, disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	- - -	-	NP	Shrubland, conifer forest, shaded streambanks, roadsides (May-Jul)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>madritensis</i>	Foxtail Chess	Po-	- - -	-	ZA	Disturbed areas, roadsides (Apr-Jan)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	- - -	H	ZA	Disturbed areas, roadsides (Mar-Jun)
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	- - B	-	NP	Seasonally wet places (May-Aug)
<input type="radio"/> <i>Carex densa</i>	Dense Sedge	Cyper-	- - A2	-	NP	
<input type="radio"/> <i>Carex</i> sp.	Sedge	Cyper-	- - -	-	-	
<input type="radio"/> <i>Cortaderia jubata</i>	Hairy Pampas Grass	Po-	- - -	H	ZP	Disturbed sites, many habitats, esp coastal (Sep-Feb)
<input type="radio"/> <i>Cortaderia selloana</i>	Smooth Pampas Grass	Po-	- - -	H	ZP	Disturbed sites (Sep-Mar)
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Po-	- - -	M	ZP	Disturbed sites (Jun-Aug)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	- - -	M	ZA	Open, disturbed sites (May-Jul)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	- - -	-	NP	Vernal pools, streambanks (May-Nov)
<input type="radio"/> <i>Cyperus erythrorhizos</i>	Redroot Nutsedge	Cyper-	- - B	-	NA	Ditches, riverbanks, shores (Jul-Oct)

Grasses & Grass-like <i>Dactylis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Dactylis</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	- - -	L	ZP	Disturbed, often moist sites (May–Aug)
<input type="radio"/> <i>Danthonia californica</i> (JM93: var. <i>californica</i>)	California Oat Grass	Po-	- - C	-	NP	Gen moist meadows, open woodland (Apr–Aug)
<input type="radio"/> <i>Deschampsia danthonioides</i>	Annual Hair Grass	Po-	- - C	-	NA	Moist to drying, open sites, meadows, streambanks, vernal pools, occ alkali soil (Mar–Aug)
<input type="radio"/> <i>Deschampsia elongata</i>	Slender Hair Grass	Po-	- - C	-	NP	Wet sites, meadows, lakeshores, shaded slopes (May–Sep)
<input type="radio"/> <i>Digitaria ischaemum</i>	Smooth Crab Grass	Po-	- - -	-	ZA	Disturbed areas (Sep–Nov)
<input type="radio"/> <i>Digitaria sanguinalis</i>	Hairy Crab Grass	Po-	- - -	-	ZA	Disturbed areas (Jun–Sep)
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Po-	- - -	-	NP	Salt marshes, coastal dunes, moist, alkaline areas (Apr–Sep)
<input type="radio"/> <i>Echinochloa crus-galli</i>	Barnyard Grass	Po-	- - -	-	ZA	Gen wet, disturbed sites, fields, roadsides (Jun–Oct)
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Po-	- - -	-	NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	- - C	-	NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus trachycaulus</i> subsp. <i>trachycaulus</i>	Slender Wheat Grass	Po-	- - B	-	NP	Dry to moist, open areas, forest, woodland (Jun–Aug)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i>)	Beardless Wild Rye	Po-	- - -	-	NP	Dry to moist, often saline, meadows (Jun–Jul)
<input type="radio"/> <i>Eragrostis curvula</i> (JM93: var. <i>curvula</i>)	Weeping Love Grass	Po-	- - -	-	NP	Roadsides (Aug–Oct)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	- - -	M	ZP	Disturbed places (May–Jun)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i>)	Brome Fescue	Po-	- - -	-	ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	- - C	-	NP	Dry, open forest, moist streambanks, chaparral (May–Jun)
<input type="radio"/> <i>Festuca elmeri</i>	Elmer Fescue	Po-	- - A2	-	NP	Moist, wooded slopes, under trees in rich soil (May–Jul)
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i>)	Hairy Fescue	Po-	- - -	-	NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i>)	Rattail Sixweeks Grass	Po-	- - -	-	ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca octoflora</i> (JM93: <i>Vulpia</i> , var. <i>hirtella</i> , <i>octoflora</i>)	Slender Six-weeks Fescue	Po-	- - A2	-	NA	Sandy to rocky soils, open sites (Mar–Jun)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i>)	Rye Grass	Po-	- - -	-	ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Po-	- - B	-	NP	Sand dunes, grassland, subalpine forest (May–Jul)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	- - -	-	NA	- -

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca sp. (perennial)</i>	Perennial Fescue	Po-	- - -	-	-	-
<input type="radio"/> <i>Gastridium phleoides (JM93: G. ventricosum)</i>	Nit Grass	Po-	- - -	-	ZA	Open, generally dry, disturbed sites (Apr–Nov)
<input type="radio"/> <i>Holcus lanatus</i>	Common Velvet Grass	Po-	- - -	M	ZP	Moist sites, roadbanks, cult fields, meadows (Jun–Aug)
<input type="radio"/> <i>Hordeum brachyantherum subsp. brachyantherum</i>	Northern Barley	Po-	- - -	-	NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum jubatum subsp. jubatum (JM93: no subsp.)</i>	Foxtail Barley	Po-	- - B	-	NAP	Roadsides, disturbed areas, meadows, marshes (May–Jul)
<input type="radio"/> <i>Hordeum marinum subsp. gussoneanum</i>	Mediterranean Barley	Po-	- - -	-	ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum subsp. leporinum</i>	Hare Barley	Po-	- - -	-	ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Hordeum murinum subsp. murinum</i>	Wall Barley	Po-	- - -	-	ZA	Moist, gen disturbed sites (Feb–May)
<input type="radio"/> <i>Juncus balticus subsp. ater (JM93: no subsp.)</i>	Baltic Rush	Junc-	- - -	-	NP	Moist to ± dry sites (Jul–Nov)
<input type="radio"/> <i>Juncus bufonius var. ?</i>	Toad Rush	Junc-	- - -	-	NA	-
<input type="radio"/> <i>Juncus bufonius var. bufonius</i>	Toad Rush	Junc-	- - -	-	NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus effusus subsp. pacificus (JM93: var.)</i>	Pacific Rush	Junc-	- - -	-	NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus occidentalis</i>	Western Rush	Junc-	- - -	-	NP	Moist gen sunny areas (May–Sep)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	- - -	-	NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus phaeocephalus var. paniculatus</i>	Panicled Brownhead Rush	Junc-	- - C	-	NP	Wet places, coastal, inland (Jun–Sep)
<input type="radio"/> <i>Juncus sp.</i>	Rush	Junc-	- - -	-	-	-
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	- - -	-	NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	- - C	-	NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Luzula comosa var. comosa (JM93: no var.)</i>	Common Woodrush	Junc-	- - -	-	NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	- - C	-	NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	- - -	-	NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	- - -	-	NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Panicum capillare</i>	Witch Grass	Po-	- - B	-	NA	Open places, fields, roadsides (Feb–Dec)
<input type="radio"/> <i>Paspalum distichum</i>	Knot Grass	Po-	- - -	-	ZP	Edges of lakes, ponds, rice fields, wet roadside ditches (Jun–Oct)
<input type="radio"/> <i>Pennisetum clandestinum</i>	Kikuyu Grass	Po-	- - -	L	ZP	Disturbed areas (All year)
<input type="radio"/> <i>Phalaris aquatica</i>	Harding Grass	Po-	- - -	M	ZP	Disturbed areas, roadsides (Apr–Aug)
<input type="radio"/> <i>Phalaris minor</i>	Little-seeded Canary Grass	Po-	- - -	-	ZA	Disturbed areas, cultivated fields (Apr–Jul)

Grasses & Grass-like <i>Phalaris</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Phalaris</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Phalaris paradoxa</i>	Hood Canary Grass	Po-	- - -	- ZA	Disturbed areas, cultivated field (May–Aug)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	- - -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa bulbosa</i> subsp. <i>vivipara</i> (JM93: no subsp.)	Bulbous Blue Grass	Po-	- - -	- ZP	Disturbed places (Mar–Jul)
<input type="radio"/> <i>Poa howellii</i>	Howell Blue Grass	Po-	- - B	- NA	Rocky banks, shaded slopes of woodland, chaparral, disturbed places (Apr–Jun)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	- - -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	- - -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Polypogon viridis</i> (JM93: <i>Agrostis</i>)	Water Beard Grass	Po-	- - -	- ZP	Common. Disturbed areas, wet areas, ponds, streambanks (May–Jun)
<input type="radio"/> <i>Schoenoplectus acutus</i> var. <i>occidentalis</i> (JM93: <i>Scirpus</i>)	Common Tule	Cyper-	- - -	- NP	Common. Marshes, shores, fens, shallow lakes, often emergent (Summer)
<input type="radio"/> <i>Scirpus microcarpus</i>	Smallfruit Bulrush	Cyper-	- - -	- NP	Marshes, wet meadows, streambanks, pond margins, sometimes weedy (Jun–Jul)
<input type="radio"/> <i>Sorghum halepense</i>	Johnson Grass	Po-	- - -	- ZP	Disturbed areas, ditch banks, roadsides (May–Aug)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i>)	Foothill Needle Grass	Po-	- - C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa miliacea</i> var. <i>miliacea</i> (JM93: <i>Piptatherum miliaceum</i>)	Smilo Grass	Po-	- - -	L ZP	Salt marshes, streambanks, chaparral, open woodland, disturbed (Mar–Sep)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Po-	- - C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Triglochin maritima</i>	Common Arrow-grass	Juncagin-	- - C	- NP	Coastal salt marshes, interior saline, brackish, alkaline marshes (Apr–Aug)
<input type="radio"/> <i>Trisetum canescens</i>	Tall False Oat	Po-	- - B	- NP	Open to shaded sites, meadows, chaparral, conifer forest (May–Aug)
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	- - -	- NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)

Herbaceous <i>Acaena</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Acaena</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acaena pinnatifida</i> var. <i>californica</i>	California Acaena	Ros-	- - C	- NP	Coastal grassland, open, rocky slopes (Mar–May)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	- - -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i>)	Spanish Clover	Fab-	- - -	- NA		
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Fab-	- - -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Fab-	- - -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon</i> or <i>Hosackia</i> sp. (JM93: <i>Lotus</i> sp.)	Lotus	Fab-	- - -	- -		
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i>)	Small-flower Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Actaea rubra</i>	Baneberry	Ranuncul-	- - B	- NP	Deep soils, moist, open to shaded sites, mixed-evergreen or conifer forests (May–Sep)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	- - -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	- - -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Allium triquetrum</i>	Three Cornered Leek	Alli-	- - -	- ZP	Locally common. Shady ± disturbed places (Mar–Apr)	
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Alli-	- - C	- NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)	
<input type="radio"/> <i>Amaranthus californicus</i>	California Amaranth	Amaranth-	- - A2	- NA	Seasonally moist flats, lake margins, disturbed areas (Jul–Oct)	
<input type="radio"/> <i>Amaranthus deflexus</i>	Low Amaranth	Amaranth-	- - -	- ZA	Railroad right-of-ways, disturbed areas (May–Nov)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Boragin-	- - -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Anaphalis margaritacea</i>	Pearly Everlasting	Aster-	- - -	- NP	Woodland, disturbed places (Jul–Oct)	
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	- - -	- NP	Generally wooded areas (Jun–Aug)	
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	- - -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	- - -	- ZA	Generally shady places (Apr–Jun)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Antirrhinum</i>	Sorted Alphabetically by Scientific Name					<i>Antirrhinum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Antirrhinum vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i> (JM93: <i>A. vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i>)	Wiry Snapdragon	Plantagin-	- - B	-	NA	Disturbed areas (Jun–Aug)
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	- - -	-	NA	Seasonally moist grassland, chaparral, woodland (Mar–May)
<input type="radio"/> <i>Apocynum androsaemifolium</i>	Bitter Dogbane	Apocyn-	- - B	-	NP	Open slopes, rocky places, with conifers, chaparral (May–Oct)
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	- - C	-	NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)
<input type="radio"/> <i>Aralia californica</i>	Elk Clover	Arali-	- - B	-	NP	Moist shade, canyons, stream sides (Jun–Aug)
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	- - -	-	NP	Common. Open to shady areas, often in drainages (May–Nov)
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	- - -	-	NP	Dry ground, valleys, foothills (May–Oct)
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvetch	Fab-	- - -	-	NA	Open, grassy areas, scrub (Mar–Jul)
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	- - -	-	NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)
<input type="radio"/> <i>Atriplex rosea</i>	Tumbling Orach	Chenopodi-	- - -	-	ZA	Common. Open, disturbed places, fields (Jul–Oct)
<input type="radio"/> <i>Baccharis glutinosa</i> (JM93: <i>B. douglasii</i>)	Marsh Baccharis	Aster-	- - -	-	NP	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)
<input type="radio"/> <i>Balsamorhiza macrolepis</i> (JM93: var. <i>macrolepis</i>)	Bigscale Balsamroot	Aster-	- 1B.2 *A1	-	NP	Open grassy or rocky slopes, valleys (Mar–Jul)
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	- - -	-	NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	- - -	L	ZA	Disturbed grassland. (Apr–Jun)
<input type="radio"/> <i>Bellis perennis</i>	English Daisy	Aster-	- - -	-	ZP	Damp, grassy areas (Dec–Sep)
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	- - -	M	ZA	Common. Disturbed areas, fields (Apr–Sep)
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	- - -	L	ZA	Disturbed areas (Jan–May)
<input type="radio"/> <i>Brodiaea elegans</i> subsp. <i>elegans</i>	Harvest Brodiaea	Themid-	- - -	-	NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)
<input type="radio"/> <i>Brodiaea terrestris</i> subsp. <i>terrestris</i>	Dwarf Brodiaea	Themid-	- - A2	-	NP	Coastal prairie, foothill woodland (Apr–Jul)
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	- - -	-	NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	- - -	-	NP	Common. Shady to open woodland, scrub (Apr–Jun)
<input type="radio"/> <i>Calochortus argillosus</i>	Clay Mariposa Lily	Lili-	- - -	-	NP	Hard clay from volcanic or metamorphic rocks (Apr–Jun)

Herbaceous <i>Calochortus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Calochortus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	- - C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	- - -	- NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calochortus umbellatus</i>	Oakland Star-tulip	Lili-	- 4.2 *A2	- NP	Open chaparral or woodland, gen on serpentine (Mar–May)	
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	- - B	- NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	- - -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	- - -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	- - -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	- - -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	- - -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	- - -	- NP	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Orobanch-	- - -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja densiflora subsp. densiflora</i>	Common Owl's-clover	Orobanch-	- - -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	- - -	- NA	Open fields, grassland (Mar–May)	
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	- - -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)	
<input type="radio"/> <i>Castilleja wightii</i>	Seaside / Coastal Paintbrush	Orobanch-	- - C	- NP	Coastal scrub (Mar–Aug)	
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i>)	California Mustard	Brassic-	- - -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun	
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	- - -	M ZAB	Pastures, disturbed places (Apr–Nov)	
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	- - -	M ZA	Disturbed fields, open woodland (Apr–Jul)	
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	- - -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)	
<input type="radio"/> <i>Centranthus ruber</i>	Red Valerian	Valerian-	- - -	- ZP	Disturbed places, rock or wall crevices, roadsides (Apr–Jul)	
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	- - -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)	
<input type="radio"/> <i>Chenopodium sp.</i>	Goosefoot	Chenopodi-	- - -	- -		
<input type="radio"/> <i>Chlorogalum pomeridianum var. pomeridianum</i>	Wavyleaf Soap Plant	Agav-	- - -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)	
<input type="radio"/> <i>Cichorium intybus</i>	Chicory	Aster-	- - -	- ZP	Common. Roadsides, disturbed places (Apr–Oct)	

Herbaceous <i>Cicuta</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Cicuta</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Cicuta douglasii</i>	Western Water-hemlock	Api-	- - B	- NP	Wet places, gen aquatic (Jun–Sep)	
<input type="radio"/> <i>Cirsium brevistylum</i>	Indian Thistle	Aster-	- - B	- NA	Moist places (Mar–Aug)	
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Aster-	- - -	- NB	Disturbed areas, grassland, woodland (May–Jul)	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	- - -	M ZB	Common. Disturbed areas (May–Oct)	
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>concinna</i>	Red Ribbons	Onagr-	- - C	- NA	Mixed-evergreen forest, woodland, coastal scrub (Apr–Jul)	
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	- - -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)	
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	- - -	- NA	Openings in woodland, forest, chaparral near coast (May–Aug)	
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	- - -	- NA	Common. Woodland (Apr–Sep)	
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i>)	Small-flower Claytonia	Monti-	- - -	- NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	- - -	- NA	Vernally moist, often shady or disturbed sites (Jan–May)	
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i>)	Yerba Buena	Lami-	- - -	- NP	Shady places, chaparral, woodland (Apr–Sep)	
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	- - -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)	
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	- - -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)	
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	- - -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)	
<input type="radio"/> <i>Corallorhiza maculata</i> var. <i>maculata</i> (JM93: no var.)	Spotted Coralroot	Orchid-	- - A2	- NP	Shaded mixed-evergreen or conifer forest, in decomposing lf litter (May–Aug)	
<input type="radio"/> <i>Corallorhiza striata</i>	Striped Coralroot	Orchid-	- - A1	- NP	Open to shaded mixed-evergreen or conifer forest, in decomposing leaf litter (Feb–Jul)	
<input type="radio"/> <i>Corethrogyne filaginifolia</i> (JM93: <i>Lessingia</i> , var. <i>filaginifolia</i>)	Common California-aster	Aster-	- - C	- NP	Coastal scrub, chaparral, grassland, foothill woodland, forest (Jul–Nov)	
<input type="radio"/> <i>Cotula australis</i>	Australian Brass Buttons	Aster-	- - -	- ZP	Common. Disturbed areas (Jan–May)	
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	- - -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)	
<input type="radio"/> <i>Crassula aquatica</i>	Water Pygmy-weed	Crassul-	- - C	- NA	Salt marshes, vernal pools, margins of lakes, ponds (Mar–Jun(Aug))	
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	- - -	- NA	Open areas (Feb–May)	
<input type="radio"/> <i>Crassula tillaea</i>	European Pygmy Weed	Crassul-	- - -	- ZA	Open, gravelly sites (Feb–May)	
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i>)	Turkey-mullein	Euphorbi-	- - -	- NA	Dry, open, often disturbed areas (May–Oct)	

Herbaceous <i>Cryptantha</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Cryptantha</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Cryptantha micromeres</i>	Minute-flowered Cryptantha	Boragin-	- - A1	- NA	Open sites, disturbed, coarse soils, chaparral, woodland, burns (Mar–Jul)	
<input type="radio"/> <i>Cryptantha torreyana</i> var. <i>pumila</i> (JM93: no var.)	Dwarf Cryptantha	Boragin-	- - A2	- NA	Generally chaparral, foothill woodland (Apr–Jun)	
<input type="radio"/> <i>Cuscuta subinclusa</i>	Canyon Dodder	Convolvul-	- - B	- NP	Generally on herbs, shrubs, in forests near streams, river canyon bottoms, salt marshes (Mar–Oct(Dec?))	
<input type="radio"/> <i>Cymbalaria muralis</i>	Kenilworth Ivy	Plantagin-	- - -	- ZP	Rock walls, shady, disturbed areas (May–Sep)	
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	- - -	M ZP	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	- - -	- NP	Chaparral, woodland (Feb–May)	
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	- - -	- NA	Rocky or sandy places (Apr–Jun)	
<input type="radio"/> <i>Deinandra corymbosa</i> (JM93: <i>Hemizonia</i> , subsp. <i>corymbosa</i>)	Coast Spikeweed	Aster-	- - A2	- NA	Grassland, openings in scrub or woodland, dunes, disturbed sites (Mar–Nov)	
<input type="radio"/> <i>Delairea odorata</i> (JM93: <i>Senecio mikanioides</i>)	German Ivy	Aster-	- - -	H ZP	Shady, ± disturbed places, riparian woodland, coastal scrub (Nov–Mar)	
<input type="radio"/> <i>Delphinium nudicaule</i>	Red Or Orange Larkspur	Ranuncul-	- - C	- NP	Moist talus, wooded, rocky slopes (Mar–Jun)	
<input type="radio"/> <i>Delphinium patens</i> subsp. <i>patens</i>	Spreading Larkspur	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Dicentra formosa</i>	Pacific Bleeding Heart	Papaver-	- - A2	- NP	Damp, shaded areas (Mar–Jul)	
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	- - -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	- - -	- NP	Open woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Digitalis purpurea</i>	Foxglove	Plantagin-	- - -	L ZB	Acid soil in open woodland, disturbed areas (May–Jul)	
<input type="radio"/> <i>Dipsacus sativus</i>	Fuller's Teasel	Dipsac-	- - -	M ZB	Disturbed areas, fields, vacant lots, pastures (May–Jul)	
<input type="radio"/> <i>Dodecatheon clevelandii</i> subsp. <i>patulum</i>	Padre Shooting Star	Primul-	- - C	- NP	Moist places, often on serpentine or in ± alkaline sites (Mar–May)	
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	- - -	- NP	Gen in shady sites (Mar–Jul)	
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i>)	Sticky Cinquefoil	Ros-	- - -	- NP	Gen ± shady or moist areas (May–Jul)	
<input type="radio"/> <i>Echinodorus berteroi</i>	Upright Burhead	Alismat-	- - A2	- NA	Ponds, ditches (Mid-summer–fall)	
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	- - -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)	
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	- - -	- NP	Dry slopes, ridges (Jun–Dec)	
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Onagr-	- - -	- NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)	

Herbaceous <i>Epilobium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Epilobium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>watsonii</i>	Watson Willowherb	Onagr-	- - B	- NP	Moist coastal bluffs, streamsides, ± disturbed sites (May–Oct)
<input type="radio"/> <i>Epipactis helleborine</i>	Broad-leaved Helleborine Orchid	Orchid-	- - -	- ZP	Generally dry slopes, roadcuts, mixed-conifer forest (Apr–Dec)
<input type="radio"/> <i>Erigeron bonariensis</i> (JM93: <i>Conyza</i>)	Flax-leaved Horseweed	Aster-	- - -	- ZA	Disturbed sites (All year)
<input type="radio"/> <i>Erigeron canadensis</i> (JM93: <i>Conyza</i>)	Horseweed	Aster-	- - -	- ZA	Disturbed places (All year)
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	- - -	- NP	--
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	- - A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Eriophyllum lanatum</i> var. <i>achilleoides</i> (JM93: var. <i>achillaeoides</i>)	Common Woolly Sunflower	Aster-	- - C	- NP	Dry, often rocky sites, chaparral, forest (May–Jul)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	- - -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	- - -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	- - -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Euphorbia lathyris</i>	Caper Spurge	Euphorbi-	- - -	- ZAB	Disturbed areas (All year)
<input type="radio"/> <i>Euphorbia peplus</i>	Petty Spurge	Euphorbi-	- - -	- ZA	Common. Disturbed areas (Feb–Aug)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	- - -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i>)	Broadleaf Aster	Aster-	- - -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	- - -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Fragaria vesca</i>	Wood Strawberry	Ros-	- - -	- NP	Gen partial shade in forest (Jan–Jul)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i>)	Checker Lily	Lili-	- - -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Fritillaria liliacea</i>	Fragrant Fritillary	Lili-	- 1B.2 *A1	- NP	Heavy soil, open hills, fields near coast (Feb–Apr)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	- - -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	- - -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	- - -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	- - -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Galium sp.</i>	Bedstraw	Rubi-	- - -	- --	
<input type="radio"/> <i>Galium triflorum</i>	Sweet-scent Bedstraw	Rubi-	- - B	- NP	Damp, shady forest (May–Jul)
<input type="radio"/> <i>Gamochaeta ustulata</i> (JM93: <i>Gnaphalium purpureum</i>)	Purple Cudweed	Aster-	- - -	- NA	Dunes, bluffs, fields, disturbed sites (Apr–Jul)
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	- - -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	- - -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)

Herbaceous <i>Geranium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Geranium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Geranium robertianum</i>	Herb Robert	Gerani-	- - -	- ZAB	Open to shaded sites (Apr–Sep)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>achilleifolia</i>	California Gilia	Polemoni-	- - B	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)
<input type="radio"/> <i>Gilia capitata</i> subsp. ?	Globe Gilia	Polemoni-	- - -	- NA	--
<input type="radio"/> <i>Gnaphalium</i> or <i>Pseudognaphalium</i> sp. (JM93: <i>Gnaphalium</i> sp.)	Everlasting Cudweed	Aster-	- - -	-	--
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i>)	Great Valley Gumplant	Aster-	- - -	- NP	Sandy or saline bottomland, roadsides (May–Nov)
<input type="radio"/> <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i>)	Hairy Gumplant	Aster-	- - C	- NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	- - -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)
<input type="radio"/> <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	- 1B.2 *A2	- NP	Open, grassy sites (Apr–Jun)
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i>)	Bristly Ox-tongue	Aster-	- - -	L ZAB	Common. Disturbed areas (All year)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Aster-	- - -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i>)	Cow Parsnip	Api-	- - -	- NP	Moist places, wooded or open (Apr–Jul)
<input type="radio"/> <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Aster-	- - -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	- - -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)
<input type="radio"/> <i>Hesperolinon californicum</i>	California Flax	Lin-	- - A2	- NA	Rocky areas, chaparral, grassland, occasionally on serpentine (Apr–Jul)
<input type="radio"/> <i>Heterotheca grandiflora</i>	Telegraph Weed	Aster-	- - -	- NA	Disturbed areas, dry streambeds, sand dunes (Jun–Oct(± all year))
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. ?	Sessileflower Goldenaster	Aster-	- - -	- NP	--
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. <i>echioides</i>	Bristly Goldenaster	Aster-	- - -	- NP	Grassland, scrub, woodland, open forest, disturbed sites (Jul–Oct)
<input type="radio"/> <i>Heuchera micrantha</i>	Small-flower Alumroot	Saxifrag-	- - C	- NP	Moist, rocky banks and cliffs (Apr–Jul)
<input type="radio"/> <i>Hieracium albiflorum</i>	White Hawkweed	Aster-	- - B	- NP	Forest (May–Sep)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	- - -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Hoita macrostachya</i>	Leather Root	Fab-	- - B	- NP	Streamsides, marshes, spring-moist places (May–Aug)
<input type="radio"/> <i>Hypericum perforatum</i>	Klamathweed	Hyperic-	- - -	M ZP	Open, disturbed areas in many plant communities (May–Aug)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Hypochaeris</i>	Sorted Alphabetically by Scientific Name					<i>Hypochaeris</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	- - -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	- - -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)	
<input type="radio"/> <i>Iris douglasiana</i>	Douglas Iris	Irid-	- - A2	- NP	Common. Grassy places, esp near coast (May–Jul)	
<input type="radio"/> <i>Iris longipetala</i>	Coast Iris	Irid-	- 4.2 *A1	- NP	Moist, coastal prairie or open coastal forest (Mar–Jun)	
<input type="radio"/> <i>Kickxia elatine</i>	Sharp Point Fluvellin	Plantagin-	- - -	- ZA	Disturbed, open places (Apr–Oct)	
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	- - -	- ZA	Roadsides, grassland (Jul–Nov)	
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	- - -	- ZA	Abundant. Disturbed places (May–Oct)	
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	- - -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)	
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: <i>subsp. ramosissima</i>)	Common Hare-leaf	Aster-	- - -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)	
<input type="radio"/> <i>Lamium amplexicaule</i>	Clasping Henbit	Lami-	- - -	- ZAB	Disturbed sites, cult or abandoned fields (Apr–Sep)	
<input type="radio"/> <i>Lasthenia californica subsp. californica</i> (JM93: <i>no subsp.</i>)	California Goldfields	Aster-	- - -	- NA	Many habitats (Feb–Jun)	
<input type="radio"/> <i>Lathyrus latifolius</i>	Perennial Sweet Pea	Fab-	- - -	- ZP	Disturbed areas, esp roadsides (Apr–May)	
<input type="radio"/> <i>Lathyrus vestitus var. vestitus</i>	Pale Purple Pacific Pea	Fab-	- - -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)	
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	- - C	- NA	Many habitats (Feb–Jul)	
<input type="radio"/> <i>Lemna gibba</i>	Swollen Duckweed	Lemn-	- - -	- NP	Common. Fresh or brackish water -	
<input type="radio"/> <i>Lepidium didymum</i> (JM93: <i>Coronopus didymus</i>)	Lesser Swine Cress	Brassic-	- - -	- ZA	Common. Disturbed areas, fields, pastures (Mar–Jul)	
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: <i>var. nitidum, oreganum</i>)	Threadleaf Peppergrass	Brassic-	- - -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
<input type="radio"/> <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	- - -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)	
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i>)	Pinklobe Leptosiphon	Polemoni-	- - C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)	
<input type="radio"/> <i>Lilium pardalinum subsp. pardalinum</i>	Leopard Lily	Lili-	- - A2	- NP	Moist places (drier along coast) (May–Aug)	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	- - -	- NP	Open, grassy slopes (Mar–Apr)	
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	- - -	- NP	Shaded slopes (Feb–Jun)	
<input type="radio"/> <i>Lobularia maritima</i>	Sweet Alyssum	Brassic-	- - -	L ZP	Disturbed areas, fields (Mar–Oct)	
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i>)	California Fluffweed	Aster-	- - -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Aster-	- - -	- ZA	Bare or grassy openings, burns (Mar–Jul)	

Herbaceous <i>Lomatium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lomatium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	- - -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium</i> sp.	Biscuit Root	Api-	- - -	- NP	--	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	- - -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lotus corniculatus</i>	Bird's-foot Trefoil	Fab-	- - -	- ZP	Open, disturbed areas (Jun–Sep)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	- - -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	
<input type="radio"/> <i>Lupinus latifolius</i> var. <i>latifolius</i>	Broadleaf Lupine	Fab-	- - -	- NP	Moist areas, open woodland (Apr–Jul)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lupinus variicolor</i>	Varied Lupine	Fab-	- - A1	- NP	Coastal terraces, beaches (Apr–Jul)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	- - -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Madia elegans</i> (JM93: subsp. <i>elegans</i> , <i>densiflora</i> , <i>vernalis</i>)	Elegant Tarweed	Aster-	- - C	- NA		
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	- - -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i>)	Fat False Solomon's Seal	Rusc-	- - -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i>)	Starry False Solomon's Seal	Rusc-	- - -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	- - -	- ZA	Disturbed places (Mar–Jun)	
<input type="radio"/> <i>Malva parviflora</i>	Cheeseweed	Malv-	- - -	- ZA	Common. Disturbed places (Mar–May)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i>)	California Man-root	Cucurbit-	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marah oregana</i> (JM93: <i>M. oreganus</i>)	Coast Man-root	Cucurbit-	- - C	- NP	Shrubby or open areas, forest edges (Mar–May)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i>)	Pineapple Weed	Aster-	- - -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	

Herbaceous <i>Medicago</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Medicago</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Medicago arabica</i>	Spotted Bur Clover	Fab-	- - -	- ZA	Disturbed and agricultural areas, fields, woodland (Mar–Jun)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Fab-	- - -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Mentha pulegium</i>	Pennyroyal	Lami-	- - -	M ZP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i>)	California Saxifrage	Saxifrag-	- - -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i>)	Slender Annual Phlox	Polemoni-	- - -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus cardinalis</i>	Scarlet Monkeyflower	Phrym-	- - C	- NP	Moist to wet places along streams, seepage areas (May–Sep)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	- - -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i>)	Coyote-mint	Lami-	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Muilla maritima</i>	Common Muilla	Themid-	- - B	- NP	Coastal scrub, grassland, woodland, valleys (Mar–Jun)	
<input type="radio"/> <i>Myosotis latifolia</i>	Broadleaved Forget-me-not	Boragin-	- - -	L ZP	Moist, disturbed, shady places (Feb–Jul)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i>)	Water Cress	Brassic-	- - -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	- - -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	- - -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Boragin-	- - -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Nuttallanthus texanus</i> (JM93: <i>Linaria canadensis</i>)	Blue Toadflax	Plantagin-	- - B	- NA	Sand or gravel (Mar–May)	
<input type="radio"/> <i>Oenanthe sarmentosa</i>	Water Parsley	Api-	- - -	- NP	Streams, marshes, ponds, gen aquatic (Jun–Oct)	
<input type="radio"/> <i>Orobanche fasciculata</i>	Clustered Broomrape	Orobanch-	- - B	- NP	Dry, gen bare places. Root parasite on various shrubs (Apr–Jul)	
<input type="radio"/> <i>Orobanche vallicola</i>	Sharp-lobe Broom-rape	Orobanch-	- - A2	- NP	Riparian woodland, on <i>Sambucus nigra</i> (Jul–Sep)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Api-	- - -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Oxalis corniculata</i>	Creeping Wood Sorrel	Oxalid-	- - -	- ZP	Disturbed areas (Most of Year)	
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	- - -	M ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Oxalis</i>	Sorted Alphabetically by Scientific Name					<i>Oxalis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Oxalis pilosa</i> (JM93: <i>O. albicans</i> subsp. <i>pilosa</i>)	Hairy Wood Sorrel	Oxalid-	- - A2	- NP	Coastal grassland, scrub, chaparral (Feb–Sep)	
<input type="radio"/> <i>Oxalis</i> sp.	Wood Sorrel	Oxalid-	- - -	- -		
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i>)	Wind Poppy	Papaver-	- - -	- NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	- - -	- NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	- - -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Persicaria amphibia</i> (JM93: <i>Polygonum amphibium</i> var. <i>emersum</i> , <i>stipulaceum</i>)	Water Smartweed	Polygon-	- - -	- NP	Shallow lakes, streams, shores (Jun–Nov)	
<input type="radio"/> <i>Petunia parviflora</i>	Wild Petunia	Solan-	- - A1	- NA	Open washes, dry streambeds (Apr–Aug)	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	- - -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	- - -	- NA	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	- - -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Pholistoma auritum</i> var. <i>auritum</i>	Fiesta Flower	Boragin-	- - B	- NA	Ocean bluffs, talus slopes, woodland, streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>macrophyllum</i> (JM93: <i>P. macrophyllum</i>)	Big-leaf Mistletoe	Visc-	- - C	- NP	On trees other than <i>Quercus</i> (especially <i>Alnus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Platanus</i> , <i>Populus</i> , <i>Robinia</i> , <i>Salix</i>) (Dec–Mar)	
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i>)	Lemon Verbena	Verben-	- - -	- NP	Wet places, pond margins (May–Nov)	
<input type="radio"/> <i>Piperia elegans</i> subsp. <i>elegans</i> (JM93: no subsp.)	Elegant Rein Orchid	Orchid-	- - B	- NP	Generally dry, open sites, scrub, conifer forest (May–Sep)	
<input type="radio"/> <i>Piperia elongata</i>	Wood Rein-orchid	Orchid-	- - A2	- NP	Generally dry sites, scrub, chaparral, mixed-evergreen or conifer forest (May–Jul)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	- - -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys trachycarpus</i>	Rough-nutlet Popcornflower	Boragin-	- - B	- NA	Shallow vernal pools, wet places in grassland, scrub, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Plantago coronopus</i>	Cut-leaf Plantain	Plantagin-	- - -	- NA	Coastal bluffs, salt marshes, trampled places, chaparral, grassy flats (Apr–Jul)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	- - -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	- - -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Plantagin-	- - -	- ZAP	Disturbed areas (Apr–Sep)	
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	- - -	- NA	Open grassland, sandy soil, burns (Mar–May)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	- - -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Polycarpon tetraphyllum</i>	Four-leaved Allseed	Caryophyll-	- - -	- ZA	Disturbed shaded areas, roadsides (Spring–fall)	

Herbaceous <i>Polygonum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Polygonum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i>)	Knotweed	Polygon-	- - -	- ZA	Disturbed places (May–Nov)	
<input type="radio"/> <i>Polygonum</i> or <i>Persicaria</i> sp. (JM93: <i>Polygonum</i> sp.)	Knotweed / Smartweed	Polygon-	- - -	- --		
<input type="radio"/> <i>Potamogeton illinoensis</i>	Shining Pondweed	Potamogeton-	- - -	- NP	Lakes, ponds, streams (Jun–Aug)	
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i>)	Hooker Fairy Bells	Lili-	- - C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)	
<input type="radio"/> <i>Prunella vulgaris</i> var. ?	European Selfheal	Lami-	- - -	- ZP	--	
<input type="radio"/> <i>Prunella vulgaris</i> var. <i>lanceolata</i>	Narrow-leaved Selfheal	Lami-	- - A1	- NP	Moist areas, gen conifer forest, woodland (May–Sep)	
<input type="radio"/> <i>Prunella vulgaris</i> var. <i>vulgaris</i>	European Selfheal	Lami-	- - -	- ZP	Moist, disturbed sites (Jun–Sep)	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i>)	California Everlasting	Aster-	- - -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i>)	Weedy Cudweed	Aster-	- - -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)	
<input type="radio"/> <i>Pseudognaphalium microcephalum</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>microcephalum</i>)	White Everlasting	Aster-	- - B	- NB	Grassy hillsides, gravelly canyon bottoms, chaparral, coastal-sage scrub (Jun–Aug)	
<input type="radio"/> <i>Pseudognaphalium ramosissimum</i> (JM93: <i>Gnaphalium</i>)	Pink Everlasting	Aster-	- - -	- NB	Dry, open slopes, woodland, sandy fields, dunes (Jul–Sep)	
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i>)	Cotton-batting Plant	Aster-	- - -	- NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	- - -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	- - B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	- - -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	- - -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)	
<input type="radio"/> <i>Ranunculus orthorhynchus</i> var. <i>bloomeri</i>	Bloomer Buttercup	Ranuncul-	- - A1	- NP	Meadows, marshy areas (Mar–May)	
<input type="radio"/> <i>Raphanus raphanistrum</i>	Jointed Charlock	Brassic-	- - -	- ZA	Disturbed areas, fields (Apr–Jul)	
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	- - -	L ZA	Disturbed areas, fields (May–Jul)	
<input type="radio"/> <i>Rorippa palustris</i> subsp. <i>palustris</i> (JM93: var. <i>occidentalis</i>)	Marsh Water Cress	Brassic-	- - A2	- NA	Generally wet places (Mar–Sep)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	- - -	M ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex conglomeratus</i>	Green Dock	Polygon-	- - -	- ZP	Common. Moist places (May–Aug)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Rumex</i>	Sorted Alphabetically by Scientific Name					<i>Rumex</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	- - -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex fueginus</i> (JM93: <i>R. maritimus</i>)	Golden Dock	Polygon-	- - A2	- NA	Riparian, disturbed places, shores, marshes, bogs, wet meadows (May–Aug)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	- - -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i>)	Thin-leaf Willow Dock	Polygon-	- - B	- NP	Wet places, margins, rocky slopes (May–Jul)	
<input type="radio"/> <i>Rumex sp.</i>	Dock	Polygon-	- - -	- -		
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	- - -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Caryophyll-	- - -	- NA	Sandy disturbed areas, river bars, streamsides (Spring–early summer)	
<input type="radio"/> <i>Sagittaria latifolia</i>	Wappato / Tule Potato / Arrowhead	Alismat-	- - A1	- NP	Ponds, slow streams (Jul–Aug)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	- - -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	- - -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	- - -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula laciniata</i>	Fringe-leaf / Coast Sanicle	Api-	- - A2	- NP	Coastal, open or shrubby slopes, woodland (Mar–May)	
<input type="radio"/> <i>Scabiosa atropurpurea</i>	Pincushion Flower	Dipsac-	- - -	- ZA	Disturbed areas (Mar–Nov)	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	- - -	- ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica</i> , <i>floribunda</i>)	California Figwort	Scrophulari-	- - -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Sedum spathulifolium</i>	Broadleaf Stonecrop	Crassul-	- - B	- NP	Outcrops, often in shade (Apr–Aug)	
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	- - C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)	
<input type="radio"/> <i>Senecio glomeratus</i> (JM93: <i>Erechtites glomerata</i>)	Cutleaf Burnweed	Aster-	- - -	M ZA	Disturbed sites, mostly coastal (Apr–Sep)	
<input type="radio"/> <i>Senecio minimus</i> (JM93: <i>Erechtites minima</i>)	Coastal Burnweed	Aster-	- - -	M ZA	Disturbed coastal sites (Jun–Sep)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	- - -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	- - -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. ?	Common Checkerbloom	Malv-	- - -	- NP	-	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>laciniata</i>	Geranium-leaved Checkerbloom	Malv-	- - C	- NP	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>malviflora</i>	Common Checkerbloom	Malv-	- - B	- NP	Coastal prairie, scrub, open forest (Mar–Jul)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	- - -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Sisymbrium</i>	Sorted Alphabetically by Scientific Name					<i>Sisymbrium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	- - -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	- - -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solanum americanum</i>	Small-flowered Nightshade	Solan-	- - -	- NAP	Open, often disturbed places (Apr–Nov)	
<input type="radio"/> <i>Solidago sp.</i>	Goldenrod	Aster-	- - -	- NP	--	
<input type="radio"/> <i>Solidago velutina subsp. californica (JM93: S. californica)</i>	California Goldenrod	Aster-	- - -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	- - -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper subsp. asper</i>	Prickly Sow Thistle	Aster-	- - -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	- - -	- ZA	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Spergula arvensis (JM93: subsp. arvensis)</i>	Stickwort	Caryophyll-	- - -	- ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Spergularia bocconi</i>	Boccone's Sand-spurry	Caryophyll-	- - -	- ZA	Salt marshes, alkaline areas, sandy soils (Spring)	
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	- - -	- ZAP	Forest, meadows, mud flats, disturbed	
<input type="radio"/> <i>Spiranthes porrifolia</i>	Western Ladies Tresses	Orchid-	- - A1	- NP	Wet meadows, freshwater marshes, seeps, grassland, oak woodland (Jun–Sep)	
<input type="radio"/> <i>Stachys rigida var. quercetorum (JM93: S. ajugoides var. rigida)</i>	Common Rigid Hedge-nettle	Lami-	- - -	- NP	Moist to ± dry places (Mar–Oct)	
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	- - -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Stephanomeria elata</i>	Tall Wreath Plant	Aster-	- - A2	- NA	Chaparral openings, grassy meadows, roadside embankments (Jul–Nov)	
<input type="radio"/> <i>Stephanomeria virgata subsp. pleurocarpa</i>	Twiggy Wreath Plant	Aster-	- - C	- NA	Chaparral openings, grassland (Jun–Nov)	
<input type="radio"/> <i>Symphotrichum chilense (JM93: Aster chilensis)</i>	Common California Wild Aster	Aster-	- - -	- NP	Grassland, salt marshes, disturbed places (Jun–Oct)	
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	- - -	- ZBP	Abundant. Esp disturbed areas (All year)	
<input type="radio"/> <i>Taraxia ovata (JM93: Camissonia)</i>	Golden Eggs Suncup	Onagr-	- - C	- NP	Grassy fields, gen clay soil (Mar–Jun)	
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	- - C	- NP	Chaparral, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Tellima grandiflora</i>	Fringe Cups	Saxifrag-	- - B	- NP	Moist slopes (Apr–Jul)	
<input type="radio"/> <i>Thalictrum fendleri var. polycarpum</i>	Foothill Meadow-rue	Ranuncul-	- - -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	- - -	- NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)	
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	- - -	M ZA	Disturbed places (Apr–Jul)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Torilis</i>	Sorted Alphabetically by Scientific Name					<i>Torilis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	- - -	- ZA	Disturbed places (Apr–Jun)	
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i>)	Common Star Lily	Melanthi-	- - -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)	
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	- - -	- ZBP	Common. Disturbed places (Mar–Nov)	
<input type="radio"/> <i>Trientalis latifolia</i>	Starflower	Myrsin-	- - C	- NP	Shaded places, esp woodland (Apr–Jul)	
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i>)	Rancheria Clover	Fab-	- - C	- NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)	
<input type="radio"/> <i>Trifolium angustifolium</i>	Narrow-leaved Clover	Fab-	- - -	- ZA	Disturbed areas (Late spring)	
<input type="radio"/> <i>Trifolium barbigerum</i> (JM93: var. <i>barbigerum</i>)	Bearded Clover	Fab-	- - B	- NP	Wet meadows, open, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Fab-	- - -	- NA	Open, grassy areas, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium campestre</i>	Hop Clover	Fab-	- - -	- ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	- - -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Fab-	- - -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	- - -	- ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium fucatum</i>	Bull Clover	Fab-	- - C	- NA	Locally abundant. Moist, open grassland, marshes, roadsides, occasionally saline or serpentine soils (Apr–Jun)	
<input type="radio"/> <i>Trifolium glomeratum</i>	Clustered Clover	Fab-	- - -	- ZA	Uncommon. Disturbed areas (Mar–May)	
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i>)	Pinpoint Clover	Fab-	- - -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	- - -	M ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium incarnatum</i>	Crimson Clover	Fab-	- - -	- ZA	Uncommon. Disturbed areas (May–Aug)	
<input type="radio"/> <i>Trifolium macraei</i>	Macrae's Clover	Fab-	- - A2	- NA	Disturbed areas, dunes (Mar–May)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	- - -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	- - -	- NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Fab-	- - C	- NA	Woody or shrubby slopes, roadsides (Mar–Jun)	
<input type="radio"/> <i>Trifolium pratense</i>	Red Clover	Fab-	- - -	- ZP	Disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Trifolium repens</i>	White Clover	Fab-	- - -	- ZP	Disturbed areas (Feb–Dec)	
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Fab-	- - -	- ZA	Meadows, roadsides, disturbed areas (Mar–Apr)	
<input type="radio"/> <i>Trifolium variegatum</i> var. ? (JM93: no var.)	White-tip Clover	Fab-	- - -	- NA	- -	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	- - -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	

Herbaceous <i>Trillium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Trillium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	- - -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)	
<input type="radio"/> <i>Triodanis biflora</i>	Small Venus-looking-glass	Campanul-	- - A2	- NA	Disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Triphysaria eriantha subsp. eriantha</i>	Yellow Johnny-tuck	Orobanch-	- - -	- NA	Grassland, foothills (Mar–May)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	- - -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triphysaria versicolor subsp. faucibarbata</i>	Smooth Owl's Clover	Orobanch-	- - A2	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	- - -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	- - -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urospermum picroides</i>	False Hawkbit	Aster-	- - -	- ZAP	Uncommon. Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Urtica dioica subsp. holosericea</i>	Hoary Nettle	Urtic-	- - -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	- - -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbascum thapsus</i>	Woolly Mullein	Scrophulari-	- - -	L ZB	Roadsides, streambanks, disturbed areas (May–Sep)	
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	- - -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	- - -	- ZA	Wet, disturbed areas, fields (Feb–May)	
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	- - -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia benghalensis</i>	One-sided Purple Vetch	Fab-	- - -	- ZP	Grassland, roadsides, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia gigantea</i>	Giant Vetch	Fab-	- - C	- NP	Coastal shrub, coastal forest, chaparral (Mar–Aug)	
<input type="radio"/> <i>Vicia sativa subsp. ?</i>	Narrowleaf / Spring Vetch	Fab-	- - -	- ZA	--	
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sp.</i>	Vetch	Fab-	- - -	-	--	
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Apocyn-	- - -	M ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	- - C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	- - -	- NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	- - C	- NP	Gen shady sites (Mar–Jun)	

Herbaceous	Grouped by Growth Form					Herbaceous
Wyethia	Sorted Alphabetically by Scientific Name					Wyethia
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="checkbox"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	- - -	-	NP	Open grassland, woodland, scrub (Mar–May(Aug))
<input type="checkbox"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	- - -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)
<input type="checkbox"/> <i>Xanthium strumarium</i>	Cocklebur	Aster-	- - -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)
<input type="checkbox"/> <i>Zeltnera sp. (JM93: Centaurium)</i>	Centaury	Gentian-	- - -	-	NA	--

Woody Acacia	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Woody Acacia
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acacia dealbata</i>	Silver Wattle	Fab-	- - -	M ZP	Locally common. Disturbed areas, often roadsides (Feb–Apr)	
<input type="radio"/> <i>Acacia melanoxylon</i>	Blackwood Acacia	Fab-	- - -	L ZP	Uncommon. Disturbed areas (Feb–Mar)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	- - -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Acer negundo</i> (JM93: var. <i>californicum</i>)	Box Elder	Sapind-	- - -	- NP	Streamsides, bottomland. (Mar–Apr)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	- - -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	- - -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Ailanthus altissima</i>	Tree Of Heaven	Simaroub-	- - -	M ZP	Disturbed areas, grassland, oak woodland, riparian areas (Jun)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	- - -	- NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Ros-	- - C	- NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	- - C	- NP	Conifer, oak forests (Mar–May)	
<input type="radio"/> <i>Arctostaphylos crustacea</i> subsp. <i>crustacea</i> (JM93: <i>A. tomentosa</i> subsp. <i>crustacea</i>)	Brittle-leaf Manzanita	Eric-	- - C	- NP	Chaparral, conifer forest (Feb–Apr)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	- - -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	- - -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Berberis pinnata</i> subsp. <i>pinnata</i>	Shinyleaf Oregon-grape	Berberid-	- - C	- NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
<input type="radio"/> <i>Carpobrotus edulis</i>	Freeway Iceplant	Aizo-	- - -	H ZP	Common. Many coastal habitats, esp. sand (All year)	
<input type="radio"/> <i>Ceanothus oliganthus</i> var. <i>sorediatus</i>	Jim Brush	Rhamn-	- - -	- NP	Slopes, ridges, chaparral, conifer forest (Jan–May)	
<input type="radio"/> <i>Ceanothus thyrsiflorus</i> var. <i>thyrsiflorus</i> (JM93: no var.)	Blue Blossom	Rhamn-	- - -	- NP	Bluffs, slopes, canyons, chaparral, coastal scrub, closed-cone-pine forest (Mar–Jun)	
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	- - C	- NP	Dry, rocky slopes, chaparral (Mar–May)	
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	- - C	- NP	Along streams, wet places (Jun–Sep)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Corn-	- - C	- NP	Generally moist places (May–Jul)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Corn-	- - C	- NP	Generally moist habitats (May–Jul)	
<input type="radio"/> <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i>)	California Hazelnut	Betul-	- - -	- NP	Common. Many habitats, esp moist, shady places (Jan–Mar)	
<input type="radio"/> <i>Cotoneaster pannosus</i> (JM93: <i>C. pannosa</i>)	Silverleaf Cotoneaster	Ros-	- - -	M ZP	Disturbed places, mixed-evergreen forest (May–Jul)	
<input type="radio"/> <i>Cytisus scoparius</i>	Scotch Broom	Fab-	- - -	H ZP	Common. Disturbed places (Apr–Jul)	

Woody	Grouped by Growth Form						Woody
Dirca	Sorted Alphabetically by Scientific Name						Dirca
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Dirca occidentalis</i>	Western Leatherwood	Thymelae-	- 1B.2 *A2	-	NP	Gen n facing slopes, mixed-evergreen forest to chaparral, gen fog belt (Nov–Mar)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>foliolosum</i>	Leafy California Buckwheat	Polygon-	- - A2	-	NP	Gravel (All year)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	- - -	-	NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Eucalyptus camaldulensis</i>	River Red Gum	Myrt-	- - -	L	ZP	Common. Disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	- - -	M	ZP	Common. Disturbed areas (Oct–Jan)	
<input type="radio"/> <i>Eucalyptus</i> sp.	Gum	Myrt-	- - -	-	ZP	- -	
<input type="radio"/> <i>Eucalyptus viminalis</i>	Manna Gum	Myrt-	- - -	-	ZP	Uncommon. Disturbed areas (Jul–Sep)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i>)	California Coffee Berry	Rhamn-	- - -	-	NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Genista monspessulana</i>	French Broom	Fab-	- - -	H	ZP	Common. Disturbed places. (Mar–Jun)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	- - C	-	NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Hedera helix</i> (JM93: subsp. <i>helix</i>)	English Ivy	Arali-	- - -	H	ZP	Woodland, open, disturbed areas (Aug–Nov)	
<input type="radio"/> <i>Hesperocyparis macrocarpa</i> (JM93: <i>Cupressus</i>)	Monterey Cypress	Cupress-	- - -	N	NP	Closed-cone-pine/cypress forests -	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	- - -	-	NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	- - -	-	NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i>)	N. California Black Walnut	Jugland-	- 1B.1 *A2	-	NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Caprifoli-	- - -	-	NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera involucrata</i> var. <i>ledebourii</i>	Black Twinberry Honeysuckle	Caprifoli-	- - B	-	NP	Moist places (May–Jul)	
<input type="radio"/> <i>Lupinus albigrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	- - -	-	NP	- -	
<input type="radio"/> <i>Lupinus albigrons</i> var. <i>albigrons</i>	Bush Lupine	Fab-	- - -	-	NP	Common. Chaparral, foothill woodland (Mar–Jun)	
<input type="radio"/> <i>Lupinus arboreus</i>	Yellow Bush Lupine	Fab-	- - A2	N	NP	Coastal bluffs, dunes, or more inland (Apr–Jul)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	- - -	-	NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Morella californica</i> (JM93: <i>Myrica</i>)	California Bayberry	Myric-	- - A2	-	NP	Coastal dunes and scrub, closed-cone pine, redwood forest (Mar–Apr)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	- - -	-	NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Phoenix canariensis</i>	Canary Island Date Palm	Arec-	- - -	L	ZP	Uncommon. Near development, disturbed areas (Oct–Apr)	

Woody	Grouped by Growth Form					Woody
<i>Physocarpus</i>	Sorted Alphabetically by Scientific Name					<i>Physocarpus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Physocarpus capitatus</i>	Pacific Ninebark	Ros-	- - C	- NP	Moist banks, n-facing slopes, mixed-conifer forest (May–Jul)	
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	- - -	- NP	Closed-cone-pine forest, oak woodland -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	- - B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Populus trichocarpa</i> (JM93: <i>P. balsamifera</i> subsp. <i>trichocarpa</i>)	Black Cottonwood	Salic-	- - C	- NP	Alluvial bottomland, streamsides (Feb–Apr)	
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Ros-	- - C	- NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Ros-	- - C	- NP	Mixed-evergreen or conifer forest (Mar–May)	
<input type="radio"/> <i>Pyracantha angustifolia</i>	Woolly Firethorn	Ros-	- - -	L ZP	Disturbed areas, fencerows, abandoned fields, roadsides (Feb–Jun)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	- - A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	- - -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>frutescens</i>	Dwarf Interior Live Oak	Fag-	- - C	- NP	Valleys, chaparral (Mar–May)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	- - -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Rhamn-	- - C	- NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	- - C	- NP	Chaparral, montane forest (Mar–Jun)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	- - -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes divaricatum</i> var. <i>pubiflorum</i>	Straggle Gooseberry	Grossulari-	- - B	- NP	Uncommon. Coastal bluffs, forest edges (Mar–May)	
<input type="radio"/> <i>Ribes malvaceum</i> var. <i>malvaceum</i>	Chaparral Currant	Grossulari-	- - B	- NP	Chaparral, oak woodland (Oct–Apr)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	- - -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes sanguineum</i> var. <i>glutinosum</i>	Red-flowering Currant	Grossulari-	- - C	- NP	Many habitats (Feb–Apr)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Fab-	- - -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	- - -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	- - -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa spithamea</i>	Coast Ground Rose	Ros-	- - C	- NP	Open forest, chaparral, especially after fire (Apr–Aug)	
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i>)	Himalayan Blackberry	Ros-	- - -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	

Woody	Grouped by Growth Form					Woody
Rubus	Sorted Alphabetically by Scientific Name					Rubus
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="checkbox"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	- - -	-	NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)
<input type="checkbox"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	- - -	-	NP	Open, disturbed areas (Mar–Jul)
<input type="checkbox"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Salic-	- - -	-	NP	Common. Floodplains, sandy gravel (Apr–May)
<input type="checkbox"/> <i>Salix laevigata</i>	Red Willow	Salic-	- - -	-	NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)
<input type="checkbox"/> <i>Salix lasiandra</i> var. <i>lasiandra</i> (JM93: <i>S. lucida</i> subsp. <i>Lasiandra</i>)	Pacific Willow	Salic-	- - -	-	NP	Wet meadows, lakeshores, riverbanks (May–Jun)
<input type="checkbox"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	- - -	-	NP	Common. Shores, marshes, meadows, etc (Jan–Jun)
<input type="checkbox"/> <i>Salix</i> sp.	Willow	Salic-	- - -	-	?P	- -
<input type="checkbox"/> <i>Salvia mellifera</i>	Black Sage	Lami-	- - -	-	NP	Coastal-sage scrub, lower chaparral (Mar–Jun)
<input type="checkbox"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i>)	Blue Elderberry	Adox-	- - -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)
<input type="checkbox"/> <i>Schinus molle</i>	Pepper Tree	Anacardi-	- - -	L	ZP	Washes, slopes, abandoned fields (Jun–Aug)
<input type="checkbox"/> <i>Sequoia sempervirens</i>	Redwood	Cupress-	- - B	-	NP	Redwood forest -
<input type="checkbox"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	- - -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)
<input type="checkbox"/> <i>Spartium junceum</i>	Spanish Broom	Fab-	- - -	H	ZP	Disturbed areas (Apr–Jun)
<input type="checkbox"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	- - -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)
<input type="checkbox"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	- - -	-	NP	Ridges, slopes, open places in woodland (Apr–May)
<input type="checkbox"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	- - -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)
<input type="checkbox"/> <i>Umbellularia californica</i>	California Bay	Laur-	- - -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)
<input type="checkbox"/> <i>Vaccinium ovatum</i>	California Huckleberry	Eric-	- - B	-	NP	Edges, clearings in conifer forest (Mar–May)
<input type="checkbox"/> <i>Vitis californica</i>	California Wild Grape	Vit-	- - -	-	NP	Streamsides, springs, canyons (May–Jun)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex deweyana ssp. *letopoda* (*C. letopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thrysiflorus (var. *thrysiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria amphibia*)

Polygonum amphibium var. *stipulaceum* (*Persicaria amphibia*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)