

Briones Regional Park

Checklist of Wild Plants

Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Briones Regional Park. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Family	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

Briones Regional Park Checklist of Wild Plants

Ferns & Horsetails		Grouped by Growth Form				Ferns & Horsetails
<i>Adiantum</i>		Sorted Alphabetically by Scientific Name				<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	- - -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Cystopteris fragilis</i>	Brittle Fern	Woodsi-	- - C	- NP	Shady, moist rock crevices, meadows, streamsides -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	- - -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum laevigatum</i>	Smooth Scouring Rush	Equiset-	- - B	- NP	Moist, sandy or gravelly areas -	
<input type="radio"/> <i>Equisetum telmateia subsp. braunii</i>	Giant Horsetail	Equiset-	- - -	- NP	Streambanks, roadside ditches, seepage areas -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	- - -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata var. mucronata</i>	Bird's-foot Fern	Pterid-	- - C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis subsp. triangularis</i>	Goldenback Fern	Pterid-	- - -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	- - -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Dryopterid-	- - -	- NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -	
<input type="radio"/> <i>Pteridium aquilinum var. pubescens</i>	Bracken Fern	Dennstaedti-	- - -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	- - -	- NP	Near streams, springs, seeps -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	- - -	- NP	Common. Open meadows, woodland, forest, subalpine (Jun–Aug)
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	- - -	- ZA	Sandy soils, open or disturbed sites (Apr–Jun)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	- - -	M ZA	Disturbed sites (Mar–Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	- - -	M ZA	Disturbed sites (Apr–Jun)
<input type="radio"/> <i>Briza maxima</i>	Rattlesnake Grass	Po-	- - -	L ZA	Shaded sites, roadsides, pastures, weedy on coastal dunes (Apr–Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	- - -	- ZA	Shaded or moist, open sites (Apr–Jul)
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	- - -	- NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr–Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	- - -	M ZA	Open, gen disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	- - -	L ZA	Fields, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	- - -	- NP	Shrubland, conifer forest, shaded streambanks, roadsides (May–Jul)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>madritensis</i>	Foxtail Chess	Po-	- - -	- ZA	Disturbed areas, roadsides (Apr–Jan)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	- - -	H ZA	Disturbed areas, roadsides (Mar–Jun)
<input type="radio"/> <i>Calamagrostis rubescens</i>	Pine Reed Grass	Po-	- - B	- NP	Wooded slopes, montane forest, chaparral, meadows (Jun–Sep)
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	- - B	- NP	Seasonally wet places (May–Aug)
<input type="radio"/> <i>Crypsis schoenoides</i>	Swamp Prickle Grass	Po-	- - -	- ZA	Wet places (Jun–Oct)
<input type="radio"/> <i>Crypsis vaginiflora</i>	Prickle Grass	Po-	- - -	- ZA	Wet soils, lake margins, vernal pools (Jun–Sep)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	- - -	M ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	- - -	- NP	Vernal pools, streambanks (May–Nov)
<input type="radio"/> <i>Echinochloa crus-galli</i>	Barnyard Grass	Po-	- - -	- ZA	Gen wet, disturbed sites, fields, roadsides (Jun–Oct)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	- - -	- NP	Common. Fresh to brackish wetland (Spring–summer)
<input type="radio"/> <i>Elymus condensatus</i> (JM93: <i>Leymus</i>)	Giant Wild-rye	Po-	- - -	- NP	Dry slopes, open woodland (Jun–Aug)
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Po-	- - -	- NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	- - C	- NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i>)	Beardless Wild Rye	Po-	- - -	- NP	Dry to moist, often saline, meadows (Jun–Jul)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i>)	Brome Fescue	Po-	- - -	- ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	- - C	- NP	Dry, open forest, moist streambanks, chaparral (May–Jun)

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i>)	Rattail Sixweeks Grass	Po-	- - -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i>)	Rye Grass	Po-	- - -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	- - -	- NA	--
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	- - -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	- - -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	- - -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Juncus balticus</i> subsp. <i>ater</i> (JM93: no subsp.)	Baltic Rush	Junc-	- - -	- NP	Moist to ± dry sites (Jul–Nov)
<input type="radio"/> <i>Juncus bufonius</i> var. ?	Toad Rush	Junc-	- - -	- NA	--
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Junc-	- - -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	- - -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	- - -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	- - C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	- - -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	- - C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	- - -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	- - -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	- - -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	- - -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Schoenoplectus californicus</i> (JM93: <i>Scirpus</i>)	Southern Bulrush	Cyper-	- - -	- NP	Common. Brackish to fresh marshes, shores (Spring–summer)
<input type="radio"/> <i>Stipa cernua</i> (JM93: <i>Nassella</i>)	Nodding Needle Grass	Po-	- - C	- NP	Grassland, chaparral, juniper woodland (Feb–Jul)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i>)	Foothill Needle Grass	Po-	- - C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Po-	- - C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like	
<i>Typha</i>	Sorted Alphabetically by Scientific Name				<i>Typha</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
○ <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	- - -	-	NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Achillea</i>	Sorted Alphabetically by Scientific Name					<i>Achillea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	- - -	- NP	Many habitats (Apr-Sep)	
<input type="radio"/> <i>Achyrachaena mollis</i>	Blow Wives	Aster-	- - -	- NA	Common. Grassy sites, often clay soils (Mar-Jun)	
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i>)	Spanish Clover	Fab-	- - -	- NA		
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Fab-	- - -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar-Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Fab-	- - -	- NP	Chaparral, roadsides, coastal sands; common (Mar-Aug)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar-Jun)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: <i>no</i> var.)	Giant Native Dandelion	Aster-	- - -	- NP	Grassland, scrub, woodland (Apr-Jul)	
<input type="radio"/> <i>Alisma triviale</i> (JM93: <i>A. plantago-aquatica</i>)	Northern Water Plantain	Alisimat-	- - B	- NP	Ponds (Spring-fall)	
<input type="radio"/> <i>Amaranthus</i> sp.	Amaranth	Amaranth-	- - -	- -		
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Boragin-	- - -	- NA	Abundant. Open, generally disturbed places (Mar-Jun)	
<input type="radio"/> <i>Amsinckia lunaris</i>	Bent-flowered Fiddleneck	Boragin-	- 1B.2 *A2	- NA	Gravelly slopes, grassland, openings in woodland (Mar-Jun)	
<input type="radio"/> <i>Amsinckia lycopsoides</i>	Bugloss-flowered Fiddleneck	Boragin-	- - B	- NA	Common. Open, grassland, foothill woodland, generally disturbed areas (Mar-Jun)	
<input type="radio"/> <i>Amsinckia tessellata</i> var. ?	Devil's Lettuce	Boragin-	- - -	- NA	- -	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar-May)	
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	- - -	- NP	Generally wooded areas (Jun-Aug)	
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	- - -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr-Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	- - -	- ZA	Generally shady places (Apr-Jun)	
<input type="radio"/> <i>Antirrhinum vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i> (JM93: <i>A. vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i>)	Wiry Snapdragon	Plantagin-	- - B	- NA	Disturbed areas (Jun-Aug)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	- - -	- NA	Seasonally moist grassland, chaparral, woodland (Mar-May)	
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	- - C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr-Sep)	
<input type="radio"/> <i>Aristolochia californica</i>	Dutchman's Pipe	Aristolochi-	- - C	- NP	Streamsides, forest, chaparral (Jan-Apr)	

Herbaceous	Grouped by Growth Form					Herbaceous
Artemisia	Sorted Alphabetically by Scientific Name					Artemisia
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	- - -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	- - -	- NP	Dry ground, valleys, foothills (May–Oct)	
<input type="radio"/> <i>Asclepias speciosa</i>	Showy Milkweed	Apocyn-	- - A2	- NP	Many habitats incl fields, roadsides (May–Sep)	
<input type="radio"/> <i>Atriplex sp.</i>	Saltbush / Orach	Chenopodi-	- - -	- -	- -	
<input type="radio"/> <i>Baccharis glutinosa (JM93: B. douglasii)</i>	Marsh Baccharis	Aster-	- - -	- NP	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	- - -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	- - -	L ZA	Disturbed grassland. (Apr–Jun)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	- - -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	- - -	M ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	- - -	L ZA	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Themid-	- - -	- NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	- - -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	- - C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus pulchellus</i>	Mount Diablo Fairy-lantern	Lili-	- 1B.2 *A2	- NP	Wooded slopes, rarely chaparral, gen n aspect (Apr–Jun)	
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	- - -	- NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	- - -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Camissoniopsis intermedia (JM93: Camissonia)</i>	Intermediate Suncup	Onagr-	- - A2	- NA	Shrubby slopes, especially burns (Mar–May)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	- - -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	- - -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	- - -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus (JM93: no subsp.)</i>	Italian Thistle	Aster-	- - -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	- - -	- NP	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Orobanch-	- - -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	- - -	- NA	Open fields, grassland (Mar–May)	

Herbaceous <i>Castilleja</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Castilleja</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	- - -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	- - -	M ZAB	Pastures, disturbed places (Apr–Nov)
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	- - -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	- - -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chenopodium album</i>	Lamb's Quarters	Chenopodi-	- - -	- ZA	Common. Disturbed areas, fields (Jun–Oct)
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	- - -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	- - -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia biloba</i> subsp. <i>biloba</i>	Mule-ear Clarkia	Onagr-	- - A2	- NA	Foothill woodland, serpentine or not (May–Aug)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	- - -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>viminea</i>	Large Clarkia	Onagr-	- - A2	- NA	Open, grassy or shrubby places (May–Jul)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	- - -	- NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	- - -	- NA	Common. Woodland (Apr–Sep)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	- - -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	- - -	- NA	Vernally moist, often shady or disturbed sites (Jan–May)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i>)	Yerba Buena	Lami-	- - -	- NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	- - -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	- - C	- NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	- - -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	- - -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)
<input type="radio"/> <i>Corallorhiza maculata</i> var. <i>maculata</i> (JM93: no var.)	Spotted Coralroot	Orchid-	- - A2	- NP	Shaded mixed-evergreen or conifer forest, in decomposing lf litter (May–Aug)
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	- - -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	- - -	- NA	Open areas (Feb–May)
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i>)	Turkey-mullein	Euphorbi-	- - -	- NA	Dry, open, often disturbed areas (May–Oct)

Herbaceous <i>Cuscuta</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Cuscuta</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Cuscuta subinclusa</i>	Canyon Dodder	Convolvul-	- - B	- NP	Generally on herbs, shrubs, in forests near streams, river canyon bottoms, salt marshes (Mar–Oct(Dec?))
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	- - -	M ZP	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	- - -	- NP	Chaparral, woodland (Feb–May)
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Ranuncul-	- - C	- NP	Open coastal grassland, chaparral (Mar–May)
<input type="radio"/> <i>Delphinium nudicaule</i>	Red Or Orange Larkspur	Ranuncul-	- - C	- NP	Moist talus, wooded, rocky slopes (Mar–Jun)
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	- - -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	- - -	- NP	Open woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	- - -	- NP	Gen in shady sites (Mar–Jul)
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i>)	Sticky Cinquefoil	Ros-	- - -	- NP	Gen ± shady or moist areas (May–Jul)
<input type="radio"/> <i>Dudleya cymosa</i> subsp. <i>cymosa</i>	Spreading Dudleya	Crassul-	- - A1	- NP	Rocky outcrops, talus slopes, less often shaded canyon slopes (May–Jul)
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	- - -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)
<input type="radio"/> <i>Epilobium campestre</i> (JM93: <i>E. pygmaeum</i>)	Smooth Willowherb	Onagr-	- - B	- NA	Vernal pools, clay mud flats (May–Sep)
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	- - -	- NP	Dry slopes, ridges (Jun–Dec)
<input type="radio"/> <i>Epipactis helleborine</i>	Broad-leaved Helleborine Orchid	Orchid-	- - -	- ZP	Generally dry slopes, roadcuts, mixed-conifer forest (Apr–Dec)
<input type="radio"/> <i>Eremalche parryi</i> subsp. <i>parryi</i>	Parry's Mallow	Malv-	- - A1x	- NA	Grassland, scrub, foothill woodland (Mar–May)
<input type="radio"/> <i>Erigeron canadensis</i> (JM93: <i>Conyza</i>)	Horseweed	Aster-	- - -	- ZA	Disturbed places (All year)
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Aster-	- - -	- NP	Grassy dunes, chaparral, oak woodland (May–Oct)
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	- - A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Eriogonum roseum</i>	Wand Wild Buckwheat	Polygon-	- - B	- NA	Common. Sand or gravel (May–Nov)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	- - -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	- - -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Erysimum capitatum</i> var. <i>capitatum</i> (JM93: var. <i>capitatum</i> , var. <i>angustatum</i>)	Western Wallflower	Brassic-	- - C	- NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)
<input type="radio"/> <i>Eschscholzia caespitosa</i>	Tufted / Foothill Poppy	Papaver-	- - A2	- NA	
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	- - -	- NP	Grassy, open areas (Feb–Sep)

Herbaceous <i>Eucrypta</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Eucrypta</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Eucrypta chrysanthemifolia</i> var. <i>chrysanthemifolia</i>	Common Eucrypta	Boragin-	- - B	- NA	Roadsides, burns, coastal bluffs, ravines (Mar–Jun)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	- - -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i>)	Checker Lily	Lili-	- - -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	- - -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	- - -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium parisiense</i>	Wall Bedstraw	Rubi-	- - -	- ZA	Warm, dry, generally rocky soil to moist areas, coastal-sage scrub, chaparral, grassy hillsides with oaks, roadsides (Apr–Aug)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	- - -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Gamochaeta ustulata</i> (JM93: <i>Gnaphalium purpureum</i>)	Purple Cudweed	Aster-	- - -	- NA	Dunes, bluffs, fields, disturbed sites (Apr–Jul)
<input type="radio"/> <i>Geranium carolinianum</i>	Carolina Geranium	Gerani-	- - -	- NA	Open to shaded sites, grassland, scrub, forest (Feb–Aug)
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	- - -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	- - -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>multicaulis</i>	Many-stem California Gilia	Polemoni-	- - -	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Feb–Jun)
<input type="radio"/> <i>Gnaphalium palustre</i>	Lowland Cudweed	Aster-	- - -	- NA	Arroyos, sandy streambeds, pond edges, potholes (May–Oct)
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i>)	Great Valley Gumplant	Aster-	- - -	- NP	Sandy or saline bottomland, roadsides (May–Nov)
<input type="radio"/> <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i>)	Hairy Gumplant	Aster-	- - C	- NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	- - -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)
<input type="radio"/> <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	- 1B.2 *A2	- NP	Open, grassy sites (Apr–Jun)
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i>)	Bristly Ox-tongue	Aster-	- - -	L ZAB	Common. Disturbed areas (All year)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>lutescens</i> (JM93: subsp. <i>congesta</i>)	Yellow Hayfield Tarweed	Aster-	- - C	- NA	Grassland, barrens, openings in chaparral and woodland, often on serpentine (Apr–Dec)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Aster-	- - -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i>)	Cow Parsnip	Api-	- - -	- NP	Moist places, wooded or open (Apr–Jul)
<input type="radio"/> <i>Hesperevax acaulis</i> var. <i>ambusticola</i>	Dwarf Fire Evax	Aster-	- - A2	- NA	Clearings, barren slopes, burns (Apr–Jun)
<input type="radio"/> <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Aster-	- - -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)

Herbaceous <i>Hesperocnide</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Hesperocnide</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	- - -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. ?	Sessileflower Goldenaster	Aster-	- - -	- NP	--
<input type="radio"/> <i>Hieracium albiflorum</i>	White Hawkweed	Aster-	- - B	- NP	Forest (May–Sep)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	- - -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Holocarpha heermannii</i>	Heermann Tarplant	Aster-	- - -	- NA	Grassland (May–Nov)
<input type="radio"/> <i>Holocarpha virgata</i> subsp. <i>virgata</i>	Sticky Tarplant	Aster-	- - -	- NA	Grassland (May–Nov)
<input type="radio"/> <i>Horkelia californica</i> var. <i>frondosa</i> (JM93: subsp. <i>frondosa</i>)	Unlobed California Horkelia	Ros-	- - C	- NP	Coastal scrub, canyons, poison-oak thickets (May–Oct)
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	- - -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	- - -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	- - -	- ZA	Roadsides, grassland (Jul–Nov)
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	- - -	- ZA	Abundant. Disturbed places (May–Oct)
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	- - -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: subsp. <i>ramosissima</i>)	Common Hare-leaf	Aster-	- - -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lasthenia californica</i> subsp. <i>californica</i> (JM93: no subsp.)	California Goldfields	Aster-	- - -	- NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lathyrus jepsonii</i> var. <i>californicus</i>	Bluff Pea	Fab-	- - B	- NP	Forest, open areas (Apr–Aug)
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	- - -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i>)	Threadleaf Peppergrass	Brassic-	- - -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)
<input type="radio"/> <i>Lepidium oxycarpum</i>	Sharp-pod Peppergrass	Brassic-	- - B	- NA	Fields, vernal pool margins, alkaline flats (Mar–May)
<input type="radio"/> <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	- - -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	- - -	- NP	Open, grassy slopes (Mar–Apr)
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	- - -	- NP	Shaded slopes (Feb–Jun)
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i>)	California Fluffweed	Aster-	- - -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Aster-	- - -	- ZA	Bare or grassy openings, burns (Mar–Jul)
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Api-	- - B	- NP	Woodland, brushy slopes (Apr–Jun)

Herbaceous <i>Lomatium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lomatium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	- - -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Api-	- - -	- NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	- - -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	- - -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	
<input type="radio"/> <i>Lupinus latifolius</i> var. <i>latifolius</i>	Broadleaf Lupine	Fab-	- - -	- NP	Moist areas, open woodland (Apr–Jul)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus pachylobus</i>	Big Pod Lupine	Fab-	- - B	- NA	Uncommon. Open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	- - -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Madia elegans</i> (JM93: subsp. <i>elegans</i> , <i>densiflora</i> , <i>vernalis</i>)	Elegant Tarweed	Aster-	- - C	- NA		
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	- - -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i>)	Fat False Solomon's Seal	Rusc-	- - -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i>)	Starry False Solomon's Seal	Rusc-	- - -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
<input type="radio"/> <i>Malacothrix floccifera</i>	Woolly / Flocked Dandelion	Aster-	- - A2	- NA	In loose soil of open burns, slides, in chaparral, pinyon/juniper woodland, yellow-pine forest (Mar–Nov)	
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	- - -	- ZA	Disturbed places (Mar–Jun)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i>)	California Man-root	Cucurbit-	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i>)	Pineapple Weed	Aster-	- - -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago lupulina</i>	Black Medick	Fab-	- - -	- ZA	Disturbed and agricultural areas, forest, mtns (Apr–Jul)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Medicago</i>	Sorted Alphabetically by Scientific Name					<i>Medicago</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Fab-	- - -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i>)	California Saxifrage	Saxifrag-	- - -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Aster-	- - -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	- - -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris elegans</i>	Nodding Silverpuffs	Aster-	- - A2	- NA	Generally inland clay grassland, often near vernal pools (Apr–Jun)	
<input type="radio"/> <i>Mimulus cardinalis</i>	Scarlet Monkeyflower	Phrym-	- - C	- NP	Moist to wet places along streams, seepage areas (May–Sep)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	- - -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i>)	Coyote-mint	Lami-	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	- - C	- NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	- - -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	- - -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Boragin-	- - -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Orobanche vallicola</i>	Sharp-lobe Broom-rape	Orobanch-	- - A2	- NP	Riparian woodland, on <i>Sambucus nigra</i> (Jul–Sep)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Api-	- - -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Oxalis corniculata</i>	Creeping Wood Sorrel	Oxalid-	- - -	- ZP	Disturbed areas (Most of Year)	
<input type="radio"/> <i>Papaver californicum</i>	Fire Poppy	Papaver-	- - A2	- NA	Burns, disturbed areas, open woodland (Apr–May)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i>)	Wind Poppy	Papaver-	- - -	- NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	- - -	- NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	- - -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Persicaria amphibia</i> (JM93: <i>Polygonum amphibium</i> var. <i>emersum</i> , <i>stipulaceum</i>)	Water Smartweed	Polygon-	- - -	- NP	Shallow lakes, streams, shores (Jun–Nov)	

Herbaceous <i>Phacelia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Phacelia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	- - -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	- - -	- NA	Common. Clay to rocky soils, slopes (Mar–May)
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	- - -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>tomentosum</i> (JM93: <i>P. villosum</i>)	Oak Mistletoe	Visc-	- - -	- NP	Generally on Quercus, rarely on Adenostoma, Arctostaphylos, Rhus, Umbellularia (Jul–Sep)
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i>)	Lemon Verbena	Verben-	- - -	- NP	Wet places, pond margins (May–Nov)
<input type="radio"/> <i>Piperia elegans</i> subsp. <i>elegans</i> (JM93: no subsp.)	Elegant Rein Orchid	Orchid-	- - B	- NP	Generally dry, open sites, scrub, conifer forest (May–Sep)
<input type="radio"/> <i>Plagiobothrys acanthocarpus</i>	Adobe Popcornflower	Boragin-	- - B	- NA	Vernal pools, moist clay soil (Mar–May)
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	- - -	- NA	Common; open woodland, grassland (Mar–May)
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	- - -	- -	- -
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	- - -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	- - -	L ZA	Common. Disturbed areas (Apr–Aug)
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	- - -	- NA	Open grassland, sandy soil, burns (Mar–May)
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	- - -	- NA	Grassy, brushy areas (Mar–Jun)
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i>)	Knotweed	Polygon-	- - -	- ZA	Disturbed places (May–Nov)
<input type="radio"/> <i>Potamogeton</i> or <i>Stuckenia</i> sp. (JM93: <i>Potamogeton</i> sp.)	Pondweed	Potamogeton-	- - -	- NP	- -
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i>)	Hooker Fairy Bells	Lili-	- - C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)
<input type="radio"/> <i>Prunella vulgaris</i> var. ?	European Selfheal	Lami-	- - -	- ZP	- -
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i>)	California Everlasting	Aster-	- - -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i>)	Weedy Cudweed	Aster-	- - -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	- - -	- NA	Common. Sand or gravel (Mar–Jul)
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	- - B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Aug)
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	- - -	- NA	Grassland, open woodland (Mar–May)
<input type="radio"/> <i>Ranunculus lobbii</i>	Lobb's Aquatic Buttercup	Ranuncul-	- 4.2 *A2	- NA	Ponds (Mar–May)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Ranunculus</i>	Sorted Alphabetically by Scientific Name					<i>Ranunculus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	- - -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)	
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	- - -	L ZA	Disturbed areas, fields (May–Jul)	
<input type="radio"/> <i>Rorippa palustris</i> subsp. <i>palustris</i> (JM93: var. <i>occidentalis</i>)	Marsh Water Cress	Brassic-	- - A2	- NA	Generally wet places (Mar–Sep)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	- - -	M ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	- - -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex obtusifolius</i>	Bitter Dock	Polygon-	- - -	- ZP	Moist places (May–Sep)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	- - -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	- - -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	- - -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	- - -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	- - -	- ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica</i> , <i>floribunda</i>)	California Figwort	Scrophulari-	- - -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberosus Skullcap	Lami-	- - C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	- - -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	- - -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sidalcea diploscypha</i>	Fringed Checkerbloom	Malv-	- - A2	- NA	Grassland, open woodland, valleys, occasionally near vernal pools, generally serpentine (Apr–May)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. ?	Common Checkerbloom	Malv-	- - -	- NP	--	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	- - -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sinapis arvensis</i>	Charlock	Brassic-	- - -	L ZA	Disturbed areas (Mar–Sep)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	- - -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisymbrium</i> sp.	Rocket / Mustard	Brassic-	- - -	- ZA	--	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	- - -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solidago elongata</i> (JM93: <i>S. canadensis</i> subsp. <i>elongata</i>)	West Coast Canada Goldenrod	Aster-	- - -	- NP	Meadows, thickets (May–Oct)	
<input type="radio"/> <i>Solidago velutina</i> subsp. <i>californica</i> (JM93: <i>S. californica</i>)	California Goldenrod	Aster-	- - -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	

Herbaceous Sonchus	Grouped by Growth Form					Herbaceous Sonchus
Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	- - -	-	ZA	Common. Slightly moist disturbed sites, along streams (All year)
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	- - -	-	ZA	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Spergula arvensis</i> (JM93: subsp. <i>arvensis</i>)	Stickwort	Caryophyll-	- - -	-	ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Stachys bullata</i>	California Hedge-nettle	Lami-	- - A1?	-	NP	Dry slopes near coast (Mar–Sep)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i>)	Common Rigid Hedge-nettle	Lami-	- - -	-	NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Stebbinsoseris heterocarpa</i>	Grassland Silverpuffs	Aster-	- - B	-	NA	Open, sometimes disturbed sites, rocky to clay soils, generally inland (Apr–Jun)
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	- - -	-	ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)
<input type="radio"/> <i>Stephanomeria virgata</i> subsp. <i>pleurocarpa</i>	Twiggy Wreath Plant	Aster-	- - C	-	NA	Chaparral openings, grassland (Jun–Nov)
<input type="radio"/> <i>Symphyotrichum chilense</i> (JM93: <i>Aster chilensis</i>)	Common California Wild Aster	Aster-	- - -	-	NP	Grassland, salt marshes, disturbed places (Jun–Oct)
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	- - -	-	ZBP	Abundant. Esp disturbed areas (All year)
<input type="radio"/> <i>Taraxia ovata</i> (JM93: <i>Camissonia</i>)	Golden Eggs Suncup	Onagr-	- - C	-	NP	Grassy fields, gen clay soil (Mar–Jun)
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	- - C	-	NP	Chaparral, pine/oak woodland (Mar–May)
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	- - -	-	NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	- - -	M	ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	- - -	-	ZA	Disturbed places (Apr–Jun)
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i>)	Common Star Lily	Melanthi-	- - -	-	NP	Grassy or wooded slopes, outcrops (Feb–Jun)
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	- - -	-	ZBP	Common. Disturbed places (Mar–Nov)
<input type="radio"/> <i>Trianthema portulacastrum</i>	Horse-purslane	Aizo-	- - A1	-	NA	Uncommon. Moist or seasonally dry wetlands, disturbed areas (Jun–Nov)
<input type="radio"/> <i>Trientalis latifolia</i>	Starflower	Myrsin-	- - C	-	NP	Shaded places, esp woodland (Apr–Jul)
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i>)	Rancheria Clover	Fab-	- - C	-	NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)
<input type="radio"/> <i>Trifolium barbigerum</i> (JM93: var. <i>barbigerum</i>)	Bearded Clover	Fab-	- - B	-	NP	Wet meadows, open, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Fab-	- - -	-	NA	Open, grassy areas, forest (Apr–Jun)
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	- - -	-	NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Fab-	- - -	-	NA	Grassland, coastal woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	- - -	-	ZA	Agricultural, disturbed areas, lawns (Spring)
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	- - -	M	ZA	Disturbed areas, roadsides (Apr–May)

Herbaceous <i>Trifolium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Trifolium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Trifolium obtusiflorum</i>	Clammy Clover	Fab-	- - B	- NA	Moist disturbed areas, gravel bars, marshes (Apr–Jul)
<input type="radio"/> <i>Trifolium pratense</i>	Red Clover	Fab-	- - -	- ZP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Fab-	- - -	- ZA	Meadows, roadsides, disturbed areas (Mar–Apr)
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	- - -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Trifolium wormskioldii</i>	Cow Clover	Fab-	- - A2	- NA	Beaches to mtn meadows, ridges, generally open moist or marshy places (May–Oct)
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	- - -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)
<input type="radio"/> <i>Triphysaria eriantha subsp. eriantha</i>	Yellow Johnny-tuck	Orobanch-	- - -	- NA	Grassland, foothills (Mar–May)
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	- - -	- NA	Grassland (Apr–Jun)
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	- - -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	- - -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)
<input type="radio"/> <i>Turritis glabra (JM93: Arabis, var. glabra)</i>	Tower Mustard	Brassic-	- - C	- NP	Open fields, meadows, slopes (Apr–Jul)
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	- - -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)
<input type="radio"/> <i>Urtica dioica subsp. holosericea</i>	Hoary Nettle	Urtic-	- - -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	- - -	- NP	Open, dry to wet places (May–Sep)
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	- - -	- ZA	Wet, disturbed areas, fields (Feb–May)
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	- - -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	- - -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	- - -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	- - -	- NP	Grassland (Apr–Aug)
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	- - C	- NP	Gen shady sites (Mar–Jun)
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	- - -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	- - -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Xanthium</i>	Sorted Alphabetically by Scientific Name					<i>Xanthium</i>
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="checkbox"/> <i>Xanthium strumarium</i>	Cocklebur	Aster-	- - -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)

Woody Acacia	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Woody Acacia
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acacia longifolia</i>	Sydney Golden Wattle	Fab-	- - -	- ZP	Uncommon. Disturbed places, especially sandy coastal areas (Jan-Apr)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	- - -	- NP	Common. Streambanks, canyons (Mar-Jun)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	- - -	- NP	Dry slopes, ridges, chaparral (May-Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	- - -	- NP	Dry slopes, canyons, borders of streams (May-Jun)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	- - -	- NP	Along permanent streams (Apr-Jun)	
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Ros-	- - C	- NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr-Jun)	
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	- - C	- NP	Conifer, oak forests (Mar-May)	
<input type="radio"/> <i>Arctostaphylos crustacea</i> subsp. <i>crustacea</i> (JM93: <i>A. tomentosa</i> subsp. <i>crustacea</i>)	Brittle-leaf Manzanita	Eric-	- - C	- NP	Chaparral, conifer forest (Feb-Apr)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	- - -	- NP	Coastal scrub, chaparral, open woodland (Aug-Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	- - -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul-Dec)	
<input type="radio"/> <i>Brickellia californica</i>	California Brickellbush	Aster-	- - B	- NP	Dry, rocky hillsides, canyons, sea bluffs (Jul-Dec)	
<input type="radio"/> <i>Ceanothus cuneatus</i> var. <i>cuneatus</i>	Buckbrush	Rhamn-	- - C	- NP	Sandy to rocky flats, slopes, ridges (Feb-May)	
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	- - C	- NP	Dry, rocky slopes, chaparral (Mar-May)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	- - -	- NP	Hillsides, chaparral, open woodland (Jan-Jun)	
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	- - C	- NP	Along streams, wet places (Jun-Sep)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Corn-	- - C	- NP	Generally moist places (May-Jul)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Corn-	- - C	- NP	Generally moist habitats (May-Jul)	
<input type="radio"/> <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i>)	California Hazelnut	Betul-	- - -	- NP	Common. Many habitats, esp moist, shady places (Jan-Mar)	
<input type="radio"/> <i>Dendromecon rigida</i>	Bush Poppy	Papaver-	- - A2	- NP	Dry slopes, washes, esp recent burns (Apr-Jun)	
<input type="radio"/> <i>Eriodictyon californicum</i>	California Yerba Santa	Boragin-	- - -	- NP	Slopes, fields, roadsides, woodland, chaparral (Apr-Jul)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. ?	California Buckwheat	Polygon-	- - -	- NP	--	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	- - -	- NP	Many dry habitats (Apr-Aug)	
<input type="radio"/> <i>Eucalyptus camaldulensis</i>	River Red Gum	Myrt-	- - -	L ZP	Common. Disturbed areas (Apr-Jul)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	- - -	M ZP	Common. Disturbed areas (Oct-Jan)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i>)	California Coffee Berry	Rhamn-	- - -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May-Jul)	

Woody Garrya	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Woody Garrya
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Garrya elliptica</i>	Coast Silk Tassel	Garry-	- - B	- NP	Seacliffs, sand dunes, chaparral, foothill-pine woodland (Jan–Mar)	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	- - -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	- - -	- NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i>)	N. California Black Walnut	Jugland-	- 1B.1 *A2	- NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Caprifoli-	- - -	- NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera involucrata</i> var. <i>ledebourii</i>	Black Twinberry Honeysuckle	Caprifoli-	- - B	- NP	Moist places (May–Jul)	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>albifrons</i>	Bush Lupine	Fab-	- - -	- NP	Common. Chaparral, foothill woodland (Mar–Jun)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	- - -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Morella californica</i> (JM93: <i>Myrica</i>)	California Bayberry	Myric-	- - A2	- NP	Coastal dunes and scrub, closed-cone pine, redwood forest (Mar–Apr)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	- - -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Olea europaea</i>	Olive	Ole-	- - -	L ZP	Generally waif, persisting from cultivation (Feb–Jun)	
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	- - -	- NP	Closed-cone-pine forest, oak woodland -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	- - B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Prunus dulcis</i>	Almond	Ros-	- - -	- ZP	Canyons, roadsides, grassland (as waif) (Feb–Mar)	
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Ros-	- - C	- NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Ros-	- - C	- NP	Mixed-evergreen or conifer forest (Mar–May)	
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Ros-	- - -	- NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Ptelea crenulata</i>	Western Hop Tree	Rut-	- - B	- NP	Scrub, woodland (Apr–May)	
<input type="radio"/> <i>Pyracantha angustifolia</i>	Woolly Firethorn	Ros-	- - -	L ZP	Disturbed areas, fencerows, abandoned fields, roadsides (Feb–Jun)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	- - A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	- - C	- NP	Dry slopes, chaparral (Feb–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	- - -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	- - -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	

Woody Quercus	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Woody Quercus
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	- - -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	- - B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>frutescens</i>	Dwarf Interior Live Oak	Fag-	- - C	- NP	Valleys, chaparral (Mar–May)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	- - -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Rhus aromatica</i> (JM93: <i>R. trilobata</i>)	Skunk Bush	Anacardi-	- - C	- NP	Slopes, washes, scrub (Mar–May)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	- - -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes divaricatum</i> var. <i>pubiflorum</i>	Straggle Gooseberry	Grossulari-	- - B	- NP	Uncommon. Coastal bluffs, forest edges (Mar–May)	
<input type="radio"/> <i>Ribes malvaceum</i> var. <i>malvaceum</i>	Chaparral Currant	Grossulari-	- - B	- NP	Chaparral, oak woodland (Oct–Apr)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	- - -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	- - -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	- - -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i>)	Himalayan Blackberry	Ros-	- - -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	- - -	- NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	- - -	- NP	Open, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Salic-	- - -	- NP	Common. Floodplains, sandy gravel (Apr–May)	
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	- - -	- NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="radio"/> <i>Salvia mellifera</i>	Black Sage	Lami-	- - -	- NP	Coastal-sage scrub, lower chaparral (Mar–Jun)	
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i>)	Blue Elderberry	Adox-	- - -	- NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	- - -	- NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="radio"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	- - -	- NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="radio"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	- - -	- NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	- - -	- NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="radio"/> <i>Umbellularia californica</i>	California Bay	Laur-	- - -	- NP	Common. Canyons, valleys, chaparral (Nov–May)	
<input type="radio"/> <i>Viburnum ellipticum</i>	Oval-leaved Viburnum	Adox-	- 2.3 *A2	- NP	Chaparral, yellow-pine forest, generally n-facing slopes (Jun–Aug)	

Woody	Grouped by Growth Form					Woody
<i>Vitis</i>	Sorted Alphabetically by Scientific Name					<i>Vitis</i>
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="checkbox"/> <i>Vitis californica</i>	California Wild Grape	Vit-	- - -	-	NP	Streamsides, springs, canyons (May–Jun)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex deweyana ssp. *leptopoda* (*C. leptopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thyrsoiflorus (var. *thyrsoiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria amphibia*)

Polygonum amphibium var. *stipulaceum* (*Persicaria amphibia*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)