

# Carquinez Strait Regional Shoreline

## Checklist of Wild Plants

### Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Carquinez Strait Regional Shoreline. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
<b>Scientific Name</b>	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> ( <a href="http://ucjeps.berkeley.edu/IJM.html">ucjeps.berkeley.edu/IJM.html</a> )
<b>(JM93 if different)</b>	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
<b>Common Name</b>	According to <i>JM2</i> and other references (not standardized)
<b>Family</b>	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
<b>Status</b>	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): <b>Federal/California (Fed./Calif.)   California Native Plant Society (CNPS)   East Bay chapter of the CNPS (EBCNPS)</b>  <b>Fed./Calif.:</b> FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare  <b>CNPS</b> (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California  <b>EBCNPS</b> (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
<b>Inv</b>	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
<b>OL</b>	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
<b>Habitat (Bloom)</b>	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, [wlegard@ebparks.org](mailto:wlegard@ebparks.org)). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

# Carquinez Strait Regional Shoreline Checklist of Wild Plants

Ferns & Horsetails		Grouped by Growth Form				Ferns & Horsetails
<i>Adiantum</i>		Sorted Alphabetically by Scientific Name				<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	-   -   -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	-   -   -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum laevigatum</i>	Smooth Scouring Rush	Equiset-	-   -   B	- NP	Moist, sandy or gravelly areas -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	-   -   -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Pterid-	-   -   C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Pterid-	-   -   -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	-   -   -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Polypodium scolieri</i>	Leather-leaf Fern	Polypodi-	-   -   A1	- NP	Coast, generally in heavy fog-drip or salt-spray zones, on plants (especially Douglas fir, Eucalyptus), dunes, rocky cliffs, bluffs, mtn ridges, granitic or volcanic rocks, mossy logs, or in soil -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis avenacea</i>	Pacific Bent Grass	Po-	-   -   -	L	ZP	Open, often disturbed place (Jun-Jul)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	-   -   -	-	NP	Common. Open meadows, woodland, forest, subalpine (Jun-Aug)
<input type="radio"/> <i>Aira caryophyllea</i>	Silver Hair Grass	Po-	-   -   -	-	ZA	Sandy soils, open or disturbed sites (Apr-Jun)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	-   -   -	M	ZA	Disturbed sites (Mar-Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	-   -   -	M	ZA	Disturbed sites (Apr-Jun)
<input type="radio"/> <i>Bolboschoenus robustus (JM93: Scirpus)</i>	Seacoast Bulrush	Cyper-	-   -   -	-	NP	Brackish to saline coastal marshes (Summer)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	-   -   -	-	ZA	Shaded or moist, open sites (Apr-Jul)
<input type="radio"/> <i>Bromus carinatus var. carinatus</i>	California Brome	Po-	-   -   -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	-   -   -	M	ZA	Open, gen disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	-   -   -	L	ZA	Fields, disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus madritensis subsp. madritensis</i>	Foxtail Chess	Po-	-   -   -	-	ZA	Disturbed areas, roadsides (Apr-Jan)
<input type="radio"/> <i>Bromus madritensis subsp. rubens</i>	Red Brome	Po-	-   -   -	H	ZA	Disturbed areas, roadsides (Mar-Jun)
<input type="radio"/> <i>Bromus tectorum</i>	Cheat Grass	Po-	-   -   -	H	ZA	Open, disturbed areas (May-Aug)
<input type="radio"/> <i>Carex sp.</i>	Sedge	Cyper-	-   -   -	-	-	-
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	-   -   -	M	ZA	Open, disturbed sites (May-Jul)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	-   -   -	-	NP	Vernal pools, streambanks (May-Nov)
<input type="radio"/> <i>Deschampsia danthonioides</i>	Annual Hair Grass	Po-	-   -   C	-	NA	Moist to drying, open sites, meadows, streambanks, vernal pools, occ alkali soil (Mar-Aug)
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Po-	-   -   -	-	NP	Salt marshes, coastal dunes, moist, alkaline areas (Apr-Sep)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	-   -   -	-	NP	Common. Fresh to brackish wetland (Spring-summer)
<input type="radio"/> <i>Elymus condensatus (JM93: Leymus)</i>	Giant Wild-rye	Po-	-   -   -	-	NP	Dry slopes, open woodland (Jun-Aug)
<input type="radio"/> <i>Elymus glaucus subsp. glaucus</i>	Western Wild-rye	Po-	-   -   -	-	NP	Open areas, chaparral, woodland, forest (Jun-Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	-   -   C	-	NP	Open, sandy to rocky areas (May-Jul)
<input type="radio"/> <i>Elymus triticoides (JM93: Leymus)</i>	Beardless Wild Rye	Po-	-   -   -	-	NP	Dry to moist, often saline, meadows (Jun-Jul)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	-   -   -	M	ZP	Disturbed places (May-Jun)
<input type="radio"/> <i>Festuca bromoides (JM93: Vulpia)</i>	Brome Fescue	Po-	-   -   -	-	ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May-Jun)
<input type="radio"/> <i>Festuca microstachys (JM93: Vulpia, var. ciliata, confusa, microstachys, pauciflora)</i>	Hairy Fescue	Po-	-   -   -	-	NA	Disturbed, open, gen sandy soils (Apr-Jun)

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i> )	Rattail Sixweeks Grass	Po-	-   -   -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i> )	Rye Grass	Po-	-   -   -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Po-	-   -   B	- NP	Sand dunes, grassland, subalpine forest (May–Jul)
<input type="radio"/> <i>Gastridium phleoides</i> (JM93: <i>G. ventricosum</i> )	Nit Grass	Po-	-   -   -	- ZA	Open, generally dry, disturbed sites (Apr–Nov)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	-   -   -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>californicum</i>	California Barley	Po-	-   -   B	- NP	Meadows, pastures, streambanks (Mar–Jul)
<input type="radio"/> <i>Hordeum jubatum</i> subsp. <i>jubatum</i> (JM93: no subsp.)	Foxtail Barley	Po-	-   -   B	- NAP	Roadsides, disturbed areas, meadows, marshes (May–Jul)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	-   -   -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Juncus bufonius</i> var. ?	Toad Rush	Junc-	-   -   -	- NA	- -
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Junc-	-   -   -	- NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	-   -   -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus</i> sp.	Rush	Junc-	-   -   -	-	- -
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	-   -   -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	-   -   C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	-   -   -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	-   -   C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	-   -   -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	-   -   -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Phalaris canariensis</i>	Canary Grass	Po-	-   -   -	- ZA	Disturbed areas (Apr–Jul)
<input type="radio"/> <i>Phalaris minor</i>	Little-seeded Canary Grass	Po-	-   -   -	- ZA	Disturbed areas, cultivated fields (Apr–Jul)
<input type="radio"/> <i>Phalaris paradoxa</i>	Hood Canary Grass	Po-	-   -   -	- ZA	Disturbed areas, cultivated field (May–Aug)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	-   -   -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa bulbosa</i> subsp. <i>vivipara</i> (JM93: no subsp.)	Bulbous Blue Grass	Po-	-   -   -	- ZP	Disturbed places (Mar–Jul)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	-   -   -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Po-	-   -   -	- ZP	Common. Streambanks (May–Aug)

<b>Grasses &amp; Grass-like</b>	<b>Grouped by Growth Form</b>				<b>Grasses &amp; Grass-like</b>
<b><i>Polypogon</i></b>	<b>Sorted Alphabetically by Scientific Name</b>				<b><i>Polypogon</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	-   -   -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i> )	Foothill Needle Grass	Po-	-   -   C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i> )	Purple Needle Grass	Po-	-   -   C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	-   -   -	- NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)

Herbaceous <i>Achillea</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Achillea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	-   -   -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Aster-	-   -   -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)	
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i> )	Colchita	Fab-	-   -   -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i> )	Deerweed	Fab-	-   -   -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i> )	California Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	-   -   -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Aster-	-   -   -	- NA	Many open habitats (May–Jun)	
<input type="radio"/> <i>Agoseris retrorsa</i>	Spearleaf Native Dandelion	Aster-	-   -   B	- NP	Scrub, oak woodland, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	-   -   -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Allium triquetrum</i>	Three Cornered Leek	Alli-	-   -   -	- ZP	Locally common. Shady ± disturbed places (Mar–Apr)	
<input type="radio"/> <i>Amaranthus</i> sp.	Amaranth	Amaranth-	-   -   -	- -		
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i> )	Common Fiddleneck	Boragin-	-   -   -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	-   -   -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	-   -   -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	-   -   -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	-   -   -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Aristolochia californica</i>	Dutchman's Pipe	Aristolochi-	-   -   C	- NP	Streamsides, forest, chaparral (Jan–Apr)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	-   -   -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	-   -   B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)	
<input type="radio"/> <i>Asclepias cordifolia</i>	Purple Milkweed	Apocyn-	-   -   A1	- NP	Rocky slopes, talus, woodland, chaparral, lava flows (May–Jul)	
<input type="radio"/> <i>Astragalus asymmetricus</i>	San Juaquin Milkvetch	Fab-	-   -   B	- NP	Grassy areas, open woodland, disturbed sites (Apr–Jul)	
<input type="radio"/> <i>Atriplex prostrata</i> (JM93: <i>A. triangularis</i> )	Fat-hen	Chenopodi-	-   -   -	- NA	Wet places, marshes (Apr–Oct)	

Herbaceous <i>Barbarea</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Barbarea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	-   -   -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Bassia hyssopifolia</i>	Hyssop-leaf Bassia	Chenopodi-	-   -   -	L ZA	Disturbed sites, fields, roadsides (May–Nov)	
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	-   -   -	L ZA	Disturbed grassland. (Apr–Jun)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	-   -   -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	-   -   -	M ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	-   -   -	L ZA	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Calandrinia breweri</i>	Brewer Red Maids	Monti-	-   4.2   *A2	- NA	Sandy to loamy soil, disturbed sites, burns (Feb–May)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	-   -   -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Calochortus pulchellus</i>	Mount Diablo Fairy-lantern	Lili-	-   1B.2   *A2	- NP	Wooded slopes, rarely chaparral, gen n aspect (Apr–Jun)	
<input type="radio"/> <i>Calystegia purpurata subsp. purpurata</i>	Climbing Morning-glory	Convolvul-	-   -   -	- NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	-   -   -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	-   -   -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	-   -   -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	-   -   -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	-   -   -	- NP	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Orobanch-	-   -   -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja densiflora subsp. densiflora</i>	Common Owl's-clover	Orobanch-	-   -   -	- NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	-   -   -	- NA	Open fields, grassland (Mar–May)	
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	-   -   -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)	
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	-   -   -	M ZAB	Pastures, disturbed places (Apr–Nov)	
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	-   -   -	M ZA	Disturbed fields, open woodland (Apr–Jul)	
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	-   -   -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)	
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	-   -   -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)	
<input type="radio"/> <i>Chenopodium album</i>	Lamb's Quarters	Chenopodi-	-   -   -	- ZA	Common. Disturbed areas, fields (Jun–Oct)	
<input type="radio"/> <i>Chlorogalum pomeridianum var. pomeridianum</i>	Wavyleaf Soap Plant	Agav-	-   -   -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	-   -   -	M ZB	Common. Disturbed areas (May–Oct)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Clarkia</i>	Sorted Alphabetically by Scientific Name					<i>Clarkia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Clarkia affinis</i>	Small Clarkia	Onagr-	-   -   C	- NA	Openings in woodland, chaparral (May–Jun)	
<input type="radio"/> <i>Clarkia biloba</i> subsp. <i>biloba</i>	Mule-ear Clarkia	Onagr-	-   -   A2	- NA	Foothill woodland, serpentine or not (May–Aug)	
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	-   -   -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)	
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>viminea</i>	Large Clarkia	Onagr-	-   -   A2	- NA	Open, grassy or shrubby places (May–Jul)	
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	-   -   -	- NA	Common. Woodland (Apr–Sep)	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	-   -   -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	-   -   -	- NA	Vernally moist, often shady or disturbed sites (Jan–May)	
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i> )	Yerba Buena	Lami-	-   -   -	- NP	Shady places, chaparral, woodland (Apr–Sep)	
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	-   -   -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)	
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	-   -   -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)	
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	-   -   -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)	
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	-   -   -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)	
<input type="radio"/> <i>Crassula aquatica</i>	Water Pygmy-weed	Crassul-	-   -   C	- NA	Salt marshes, vernal pools, margins of lakes, ponds (Mar–Jun(Aug))	
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	-   -   -	- NA	Open areas (Feb–May)	
<input type="radio"/> <i>Cressa truxillensis</i>	Alkali Weed	Convolvul-	-   -   -	- NP	Saline and alkaline substrates (May–Oct)	
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	-   -   -	M ZP	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	-   -   -	- NP	Chaparral, woodland (Feb–May)	
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	-   -   -	- NA	Rocky or sandy places (Apr–Jun)	
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	-   -   -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	-   -   -	- NP	Open woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Dipsacus sativus</i>	Fuller's Teasel	Dipsac-	-   -   -	M ZB	Disturbed areas, fields, vacant lots, pastures (May–Jul)	
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	-   -   -	- NP	Gen in shady sites (Mar–Jul)	
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i> )	Sticky Cinquefoil	Ros-	-   -   -	- NP	Gen ± shady or moist areas (May–Jul)	
<input type="radio"/> <i>Dysphania ambrosioides</i> (JM93: <i>Chenopodium</i> )	Mexican Tea	Chenopodi-	-   -   -	- ZAP	Disturbed places (Jul–Sep)	



Herbaceous <i>Epilobium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Epilobium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	-   -   -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)
<input type="radio"/> <i>Epilobium minutum</i>	Chaparral Willowherb	Onagr-	-   -   -	- NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)
<input type="radio"/> <i>Erigeron sp. (JM93: Conyza)</i>	Fleabane Daisy	Aster-	-   -   -	- -	
<input type="radio"/> <i>Eriogonum nudum var. auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	-   -   A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	-   -   -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	-   -   -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Erysimum capitatum var. capitatum (JM93: var. capitatum, var. angustatum)</i>	Western Wallflower	Brassic-	-   -   C	- NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	-   -   -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Eucrypta chrysanthemifolia var. chrysanthemifolia</i>	Common Eucrypta	Boragin-	-   -   B	- NA	Roadsides, burns, coastal bluffs, ravines (Mar–Jun)
<input type="radio"/> <i>Euphorbia lathyris</i>	Caper Spurge	Euphorbi-	-   -   -	- ZAB	Disturbed areas (All year)
<input type="radio"/> <i>Euphorbia oblongata</i>	European Spurge	Euphorbi-	-   -   -	L ZP	Disturbed areas (Jun–Aug)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	-   -   -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina (JM93: Aster radulinus)</i>	Broadleaf Aster	Aster-	-   -   -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	-   -   -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Frankenia salina</i>	Alkali Heath	Frankeni-	-   -   -	- NP	Salt marshes, alkali flats (Apr–Sep)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	-   -   -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum subsp. californicum</i>	California Bedstraw	Rubi-	-   -   -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	-   -   -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)
<input type="radio"/> <i>Galium porrigens var. porrigens</i>	Climbing Bedstraw	Rubi-	-   -   -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	-   -   -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	-   -   -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Gilia tricolor subsp. ?</i>	Bird's-eye Gilia	Polemoni-	-   -   -	- NA	--
<input type="radio"/> <i>Gnaphalium palustre</i>	Lowland Cudweed	Aster-	-   -   -	- NA	Arroyos, sandy streambeds, pond edges, potholes (May–Oct)
<input type="radio"/> <i>Grindelia camporum (JM93: var. camporum, G. hirsutula var. davyi)</i>	Great Valley Gumplant	Aster-	-   -   -	- NP	Sandy or saline bottomland, roadsides (May–Nov)
<input type="radio"/> <i>Grindelia sp.</i>	Gumplant	Aster-	-   -   -	- NP	--

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Helianthella</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Helianthella</i></b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	-   1B.2   *A2	- NP	Open, grassy sites (Apr–Jun)	
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i> )	Bristly Ox-tongue	Aster-	-   -   -	L ZAB	Common. Disturbed areas (All year)	
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Aster-	-   -   -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)	
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i> )	Cow Parsnip	Api-	-   -   -	- NP	Moist places, wooded or open (Apr–Jul)	
<input type="radio"/> <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Aster-	-   -   -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)	
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. <i>echioides</i>	Bristly Goldenaster	Aster-	-   -   -	- NP	Grassland, scrub, woodland, open forest, disturbed sites (Jul–Oct)	
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	-   -   -	M ZBP	Disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Holocarpha virgata</i> subsp. <i>virgata</i>	Sticky Tarplant	Aster-	-   -   -	- NA	Grassland (May–Nov)	
<input type="radio"/> <i>Horkelia californica</i> var. <i>frondosa</i> (JM93: subsp. <i>frondosa</i> )	Unlobed California Horkelia	Ros-	-   -   C	- NP	Coastal scrub, canyons, poison-oak thickets (May–Oct)	
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	-   -   -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	-   -   -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)	
<input type="radio"/> <i>Iris pseudacorus</i>	Yellow Flag	Irid-	-   -   -	L ZP	Common. Pond margins, estuaries (Apr–Jun)	
<input type="radio"/> <i>Iva axillaris</i> (JM93: subsp. <i>robustior</i> )	Poverty Weed	Aster-	-   -   -	- NP	Seasonally wet, saline habitats, roadsides (Apr–Oct)	
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	-   -   -	- ZA	Abundant. Disturbed places (May–Oct)	
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	-   -   -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)	
<input type="radio"/> <i>Lathyrus latifolius</i>	Perennial Sweet Pea	Fab-	-   -   -	- ZP	Disturbed areas, esp roadsides (Apr–May)	
<input type="radio"/> <i>Lathyrus</i> sp.	Pea	Fab-	-   -   -	- - -		
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	-   -   -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)	
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	-   -   C	- NA	Many habitats (Feb–Jul)	
<input type="radio"/> <i>Lepidium draba</i> (JM93: <i>Cardaria</i> )	Heart-podded Hoary Cress	Brassic-	-   -   -	M ZP	Disturbed areas, saline soils, pastures, fields (Apr–Aug)	
<input type="radio"/> <i>Lepidium latifolium</i>	Perennial Peppergrass	Brassic-	-   -   -	H ZP	Pastures, disturbed areas, fields, grassland, saline meadows, streambanks, sagebrush scrub, edge of marshes (Jun–Sep)	
<input type="radio"/> <i>Lepidium latipes</i> (JM93: var. <i>latipes</i> )	Dwarf Peppergrass	Brassic-	-   -   B	- NA	Alkaline soils, vernal pool margins, salt marsh edges, pastures (Mar–Jun)	
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i> )	Threadleaf Peppergrass	Brassic-	-   -   -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	

Herbaceous <i>Lepidium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Lepidium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Lepidium oxycarpum</i>	Sharp-pod Peppergrass	Brassic-	-   -   B	- NA	Fields, vernal pool margins, alkaline flats (Mar–May)
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i> )	Bicolor Leptosiphon	Polemoni-	-   -   -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	-   -   -	- NP	Open, grassy slopes (Mar–Apr)
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	-   -   -	- NP	Shaded slopes (Feb–Jun)
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i> )	California Fluffweed	Aster-	-   -   -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i> )	Daggerleaf Cottonrose	Aster-	-   -   -	- ZA	Bare or grassy openings, burns (Mar–Jul)
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	-   -   -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	-   -   -	- NP	Open grassy slopes, meadows, woodland (Feb–May)
<input type="radio"/> <i>Lotus corniculatus</i>	Bird's-foot Trefoil	Fab-	-   -   -	- ZP	Open, disturbed areas (Jun–Sep)
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	-   -   -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	-   -   -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)
<input type="radio"/> <i>Lythrum</i> sp.	Loosestrife	Lythr-	-   -   -	- -	- -
<input type="radio"/> <i>Madia exigua</i>	Thread-stem Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites, in sandy or clayey soils, including serpentine (Apr–Jul)
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	-   -   -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites (May–Oct)
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i> )	Fat False Solomon's Seal	Rusc-	-   -   -	- NP	Moist, open woodland, streambanks (Mar–Jul)
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	-   -   -	- ZA	Disturbed places (Mar–Jun)
<input type="radio"/> <i>Malvella leprosa</i>	Alkal-mallow	Malv-	-   -   -	- NP	Valleys, gen saline (Apr–Nov)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Marah</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Marah</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i> )	California Man-root	Cucurbit-	-   -   -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	-   -   -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i> )	Pineapple Weed	Aster-	-   -   -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	-   -   -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus albus</i> (JM93: <i>M. alba</i> )	White Sweetclover	Fab-	-   -   -	- ZAB	Locally abundant. Pastures, open disturbed sites (May–Sep)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i> )	Sourclover	Fab-	-   -   -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Mentha spicata</i> (JM93: <i>var. spicata</i> )	Spearmint	Lami-	-   -   -	- ZP	Moist places, marshes, lakeshores, fields (Jul–Oct)	
<input type="radio"/> <i>Micropus californicus var. californicus</i>	Cottonweed	Aster-	-   -   -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris douglasii subsp. ?</i>	Douglas Silverpuffs	Aster-	-   -   -	- NA	--	
<input type="radio"/> <i>Microseris douglasii subsp. douglasii</i>	Douglas Silverpuffs	Aster-	-   -   -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris elegans</i>	Nodding Silverpuffs	Aster-	-   -   A2	- NA	Generally inland clay grassland, often near vernal pools (Apr–Jun)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	-   -   -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Monardella villosa subsp. villosa</i> (JM93: <i>M. villosa subsp. villosa, globosa, M. antonina subsp. antonina</i> )	Coyote-mint	Lami-	-   -   -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i> )	Water Cress	Brassic-	-   -   -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	-   -   C	- NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	-   -   -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	-   -   -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii var. menziesii</i>	Baby Blue-eyes	Boragin-	-   -   -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Nemophila parviflora var. parviflora</i>	Small-flower Nemophila	Boragin-	-   -   C	- NA	Woodland, forest, roadsides, slopes (Mar–Jul)	
<input type="radio"/> <i>Nuttallanthus texanus</i> (JM93: <i>Linaria canadensis</i> )	Blue Toadflax	Plantagin-	-   -   B	- NA	Sand or gravel (Mar–May)	

Herbaceous <i>Osmorhiza</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Osmorhiza</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i> )	Sweet-cicely	Api-	-   -   -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Oxalis corniculata</i>	Creeping Wood Sorrel	Oxalid-	-   -   -	- ZP	Disturbed areas (Most of Year)	
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	-   -   -	M ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i> )	Wind Poppy	Papaver-	-   -   -	- NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	-   -   -	- NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	-   -   -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	-   -   -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	-   -   -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Phacelia tanacetifolia</i>	Tansy-leaf Phacelia	Boragin-	-   -   B	- NA	Sandy to gravelly slopes, open areas (Mar–May)	
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i> )	Lemon Verbena	Verben-	-   -   -	- NP	Wet places, pond margins (May–Nov)	
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Boragin-	-   -   -	- NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	-   -   -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	-   -   -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Plantagin-	-   -   -	- ZAP	Disturbed areas (Apr–Sep)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: subsp. <i>ciliosa</i> , <i>insignis</i> )	Longspur Plectritis	Valerian-	-   -   B	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	-   -   -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i> )	Knotweed	Polygon-	-   -   -	- ZA	Disturbed places (May–Nov)	
<input type="radio"/> <i>Pseudognaphalium beneolens</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>beneolens</i> )	Scented White Everlasting	Aster-	-   -   C	- NB	Dry, open slopes, disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i> )	California Everlasting	Aster-	-   -   -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i> )	Weedy Cudweed	Aster-	-   -   -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	-   -   -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	-   -   -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus canus</i> var. <i>canus</i> (JM93: no var.)	Sacramento Valley Buttercup	Ranuncul-	-   -   B	- NP	Grassland, open oak woodlan (Mar–Jul)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Ranunculus</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Ranunculus</i></b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	-   -   -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	-   -   -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)	
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	-   -   -	L ZA	Disturbed areas, fields (May–Jul)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	-   -   -	M ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	-   -   -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex obtusifolius</i>	Bitter Dock	Polygon-	-   -   -	- ZP	Moist places (May–Sep)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	-   -   -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	-   -   -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Sagina decumbens subsp. occidentalis</i>	Western Pearlwort	Caryophyll-	-   -   C	- NA	Dry streams, chaparral, grassy areas, rock outcrops, vernal pools (Spring–early summer)	
<input type="radio"/> <i>Salicornia pacifica (JM93: S. virginica)</i>	Pacific Pickleweed	Chenopodi-	-   -   -	- NP	Salt marshes, alkaline flats (Jul–Nov)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	-   -   -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	-   -   -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	-   -   -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	-   -   -	- ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Scrophularia californica (JM93: subsp. californica, floribunda)</i>	California Figwort	Scrophulari-	-   -   -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberous Skullcap	Lami-	-   -   C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Senecio sylvaticus</i>	Woodland Ragwort	Aster-	-   -   -	- ZA	Disturbed woodland, open, sandy sites (Mar–Sep)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	-   -   -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	-   -   -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sidalcea malviflora subsp. malviflora</i>	Common Checkerbloom	Malv-	-   -   B	- NP	Coastal prairie, scrub, open forest (Mar–Jul)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	-   -   -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	-   -   -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sinapis arvensis</i>	Charlock	Brassic-	-   -   -	L ZA	Disturbed areas (Mar–Sep)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	-   -   -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	-   -   -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solanum xanti</i>	Purple Nightshade	Solan-	-   -   A1?	- NP	Shrubland, oak/pine woodland, conifer forest (Feb–Jun)	
<input type="radio"/> <i>Solidago velutina subsp. californica (JM93: S. californica)</i>	California Goldenrod	Aster-	-   -   -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	

Herbaceous <i>Soliva</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Soliva</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	-   -   -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper subsp. asper</i>	Prickly Sow Thistle	Aster-	-   -   -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	-   -   -	- ZA	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Spergularia bocconi</i>	Boccone's Sand-spurry	Caryophyll-	-   -   -	- ZA	Salt marshes, alkaline areas, sandy soils (Spring)	
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	-   -   -	- ZAP	Forest, meadows, mud flats, disturbed	
<input type="radio"/> <i>Stachys rigida var. quercetorum (JM93: S. ajugoides var. rigida)</i>	Common Rigid Hedge-nettle	Lami-	-   -   -	- NP	Moist to ± dry places (Mar–Oct)	
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	-   -   -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	-   -   C	- NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)	
<input type="radio"/> <i>Stephanomeria virgata subsp. pleurocarpa</i>	Twiggy Wreath Plant	Aster-	-   -   C	- NA	Chaparral openings, grassland (Jun–Nov)	
<input type="radio"/> <i>Symphotrichum chilense (JM93: Aster chilensis)</i>	Common California Wild Aster	Aster-	-   -   -	- NP	Grassland, salt marshes, disturbed places (Jun–Oct)	
<input type="radio"/> <i>Thalictrum fendleri var. polycarpum</i>	Foothill Meadow-rue	Ranuncul-	-   -   -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	-   -   -	M ZA	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	-   -   -	- ZA	Disturbed places (Apr–Jun)	
<input type="radio"/> <i>Toxicoscordion fremontii (JM93: Zigadenus)</i>	Common Star Lily	Melanthi-	-   -   -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)	
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	-   -   -	- ZBP	Common. Disturbed places (Mar–Nov)	
<input type="radio"/> <i>Tribulus terrestris</i>	Puncture Vine	Zygophyll-	-   -   -	- ZA	Dry, disturbed areas incl roadsides, railways, vacant lots (Apr–Oct)	
<input type="radio"/> <i>Trifolium bifidum var. decipiens</i>	Deceiving Clover	Fab-	-   -   -	- NA	Open, grassy areas, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	-   -   -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium depauperatum var. amplexans</i>	Fused-bract Pale Sack Clover	Fab-	-   -   -	- NA	Grassland, coastal woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium depauperatum var. truncatum</i>	Truncate Sack Clover	Fab-	-   -   -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	-   -   -	- ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium gracilentum (JM93: var. gracilentum)</i>	Pinpoint Clover	Fab-	-   -   -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	-   -   -	M ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium incarnatum</i>	Crimson Clover	Fab-	-   -   -	- ZA	Uncommon. Disturbed areas (May–Aug)	

Herbaceous <i>Trifolium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Trifolium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	-   -   -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	-   -   -	- NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium repens</i>	White Clover	Fab-	-   -   -	- ZP	Disturbed areas (Feb–Dec)	
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Fab-	-   -   -	- ZA	Meadows, roadsides, disturbed areas (Mar–Apr)	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	-   -   -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Triphysaria eriantha subsp. eriantha</i>	Yellow Johnny-tuck	Orobanch-	-   -   -	- NA	Grassland, foothills (Mar–May)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	-   -   -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	-   -   -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	-   -   -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Urtica dioica subsp. holosericea</i>	Hoary Nettle	Urtic-	-   -   -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	-   -   -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	-   -   -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	-   -   -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)	
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Apocyn-	-   -   -	M ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	-   -   -	- NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	-   -   C	- NP	Gen shady sites (Mar–Jun)	
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	-   -   -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Xanthium strumarium</i>	Cocklebur	Aster-	-   -   -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	



<b>Herbaceous</b>	<b>Grouped by Growth Form</b>				<b>Herbaceous</b>
<b><i>Zantedeschia</i></b>	<b>Sorted Alphabetically by Scientific Name</b>				<b><i>Zantedeschia</i></b>
<input checked="" type="checkbox"/> <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>
<input type="checkbox"/> <i>Zantedeschia aethiopica</i>	Calla-lily	Ar-	-   -   -	L ZP	Disturbed areas, near former habitations (Mar-Jun)

<b>Woody Acacia</b>	<b>Grouped by Growth Form Sorted Alphabetically by Scientific Name</b>					<b>Woody Acacia</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acacia melanoxylon</i>	Blackwood Acacia	Fab-	-   -   -	L ZP	Uncommon. Disturbed areas (Feb-Mar)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	-   -   -	- NP	Common. Streambanks, canyons (Mar-Jun)	
<input type="radio"/> <i>Acer negundo</i> (JM93: var. <i>californicum</i> )	Box Elder	Sapind-	-   -   -	- NP	Streamsides, bottomland. (Mar-Apr)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	-   -   -	- NP	Dry slopes, ridges, chaparral (May-Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	-   -   -	- NP	Dry slopes, canyons, borders of streams (May-Jun)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	-   -   -	- NP	Coastal scrub, chaparral, open woodland (Aug-Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	-   -   -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul-Dec)	
<input type="radio"/> <i>Ceanothus thyrsiflorus</i> var. <i>thyrsiflorus</i> (JM93: no var.)	Blue Blossom	Rhamn-	-   -   -	- NP	Bluffs, slopes, canyons, chaparral, coastal scrub, closed-cone-pine forest (Mar-Jun)	
<input type="radio"/> <i>Cephalanthus occidentalis</i> (JM93: var. <i>californicus</i> )	California Button Willow	Rubi-	-   -   B	- NP	Lake, stream edges (May-Sep)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	-   -   -	- NP	Hillsides, chaparral, open woodland (Jan-Jun)	
<input type="radio"/> <i>Cotoneaster pannosus</i> (JM93: <i>C. pannosa</i> )	Silverleaf Cotoneaster	Ros-	-   -   -	M ZP	Disturbed places, mixed-evergreen forest (May-Jul)	
<input type="radio"/> <i>Dirca occidentalis</i>	Western Leatherwood	Thymelae-	-   1B.2   *A2	- NP	Gen n facing slopes, mixed-evergreen forest to chaparral, gen fog belt (Nov-Mar)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	-   -   -	- NP	Many dry habitats (Apr-Aug)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	-   -   -	M ZP	Common. Disturbed areas (Oct-Jan)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i> )	California Coffee Berry	Rhamn-	-   -   -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May-Jul)	
<input type="radio"/> <i>Hedera helix</i> (JM93: subsp. <i>helix</i> )	English Ivy	Arali-	-   -   -	H ZP	Woodland, open, disturbed areas (Aug-Nov)	
<input type="radio"/> <i>Helianthemum scoparium</i>	Peak Rush-rose	Cist-	-   -   B	- NP	Dry sandy or rocky soil of hills, slopes, ridges (Feb-Sep)	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	-   -   -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun-Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	-   -   -	- NP	Moist woodland edges, rocky slopes (May-Aug)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i> )	N. California Black Walnut	Jugland-	-   1B.1   *A2	- NP	Along streams, disturbed slopes (Apr-May)	
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	-   -   -	- NP	- -	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	-   -   -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar-Jun)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Oemleria</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Oemleria</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	-   -   -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Olea europaea</i>	Olive	Ole-	-   -   -	L ZP	Generally waif, persisting from cultivation (Feb–Jun)	
<input type="radio"/> <i>Phoenix canariensis</i>	Canary Island Date Palm	Arec-	-   -   -	L ZP	Uncommon. Near development, disturbed areas (Oct–Apr)	
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	-   -   -	- NP	Closed-cone-pine forest, oak woodland -	
<input type="radio"/> <i>Populus fremontii subsp. fremontii</i>	Fremont Cottonwood	Salic-	-   -   -	- NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)	
<input type="radio"/> <i>Prunus dulcis</i>	Almond	Ros-	-   -   -	- ZP	Canyons, roadsides, grassland (as waif) (Feb–Mar)	
<input type="radio"/> <i>Prunus virginiana var. demissa</i>	Western Choke Cherry	Ros-	-   -   -	- NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Ptelea crenulata</i>	Western Hop Tree	Rut-	-   -   B	- NP	Scrub, woodland (Apr–May)	
<input type="radio"/> <i>Pyracantha angustifolia</i>	Woolly Firethorn	Ros-	-   -   -	L ZP	Disturbed areas, fencerows, abandoned fields, roadsides (Feb–Jun)	
<input type="radio"/> <i>Quercus agrifolia var. agrifolia</i>	Coast Live Oak	Fag-	-   -   A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	-   -   -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	-   -   -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	-   -   B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Ribes californicum var. californicum</i>	Hillside Gooseberry	Grossulari-	-   -   -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Fab-	-   -   -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	-   -   -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa var. gymnocarpa (JM93: no var.)</i>	Wood Rose	Ros-	-   -   -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa sp.</i>	Rose	Ros-	-   -   -	- NP	- -	
<input type="radio"/> <i>Rosa spithamea</i>	Coast Ground Rose	Ros-	-   -   C	- NP	Open forest, chaparral, especially after fire (Apr–Aug)	
<input type="radio"/> <i>Rubus armeniacus (JM93: R. discolor)</i>	Himalayan Blackberry	Ros-	-   -   -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	-   -   -	- NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	-   -   -	- NP	Open, disturbed areas (Mar–Jul)	

<b>Woody</b>	<b>Grouped by Growth Form</b>						<b>Woody</b>
<b>Salix</b>	<b>Sorted Alphabetically by Scientific Name</b>						<b>Salix</b>
<input checked="" type="checkbox"/> <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>	
<input type="checkbox"/> <i>Salix laevigata</i>	Red Willow	Salic-	-   -   -	-	NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)	
<input type="checkbox"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	-   -   -	-	NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="checkbox"/> <i>Sambucus nigra subsp. caerulea (JM93: S. mexicana)</i>	Blue Elderberry	Adox-	-   -   -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="checkbox"/> <i>Schinus molle</i>	Pepper Tree	Anacardi-	-   -   -	L	ZP	Washes, slopes, abandoned fields (Jun–Aug)	
<input type="checkbox"/> <i>Symphoricarpos albus var. laevigatus</i>	Snowberry	Caprifoli-	-   -   -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="checkbox"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	-   -   -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="checkbox"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	-   -   -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="checkbox"/> <i>Umbellularia californica</i>	California Bay	Laur-	-   -   -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)	

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Acer negundo* var. *californicum* (no var.)

*Adenostoma fasciculatum* (var. *fasciculatum*)

*Agoseris grandiflora* (var. *grandiflora*)

*Agoseris heterophylla* (var. *cryptopleura*)

*Agrostis viridis* (*Polypogon*)

*Alisma plantago-aquatica* (*A. triviale*)

*Amsinckia menziesii* var. *intermedia* (*A. intermedia*)

*Amsinckia menziesii* var. *menziesii* (no var.)

*Antirrhinum vexillo-calyculatum* ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

*Arabis breweri* var. *breweri* (*Boechera breweri* subsp. *breweri*)

*Arabis glabra* var. *glabra* (*Turritis glabra*)

*Arctostaphylos tomentosa* ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

*Aster chilensis* (*Symphyotrichum chilense*)

*Aster lentus* (*Symphyotrichum lentum*)

*Aster radulinus* (*Eurybia radulina*)

*Aster subulatus* var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

*Atriplex joaquiniana* (*A. joaquinana*)

*Atriplex lentiformis* ssp. *lentiformis* (no subsp.)

*Atriplex triangularis* (*A. prostrata*)

*Azolla mexicana* (*A. microphylla*)

*Baccharis douglasii* (*B. glutinosa*)

*Baccharis pilularis* (subsp. *consanguinea*)

*Baccharis salicifolia* (subsp. *salicifolia*)

*Balsamorhiza macrolepis* var. *macrolepis* (no var.)

*Blepharizonia plumosa* ssp. *plumosa* (no subsp.)

*Blepharizonia plumosa* ssp. *viscida* (*Blepharizonia laxa*)

*Boschniakia strobilacea* (*Kopsiopsis*)

*Bromus catharticus* (var. *catharticus*)

*Camissonia boothii* ssp. *decorticans* (*Eremothera*)

*Camissonia graciliflora* (*Tetrapteron graciliflorum*)

*Camissonia intermedia* (*Camissoniopsis*)

*Camissonia micrantha* (*Camissoniopsis*)

*Camissonia ovata* (*Taraxia*)

*Cardaria draba* (*Lepidium*)

*Carduus pycnocephalus* (subsp. *pycnocephalus*)

*Carex deweyana* ssp. *leptopoda* (*C. leptopoda*)

*Carex lanuginosa* (*C. pellita*)

*Carex ovalis* (*C. leporina*)

*Ceanothus thrysiflorus* (var. *thrysiflorus*)

*Centaurium davyi* (*Zeltnera*)

*Centaurium muehlenbergii* (*Zeltnera*)

*Cephalanthus occidentalis* var. *californicus* (no var.)

*Chamomilla suaveolens* (*Matricaria discoidea*)

*Chenopodium ambrosioides* (*Dysphania*)

*Chenopodium macrospermum* var. *halophilum* (no var.)

*Chenopodium multifidum* (*Dysphania multifida*)

*Chenopodium rubrum* (var. *rubrum*)

*Chrysothamnus nauseosus* ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

*Cirsium cymosum* (var. *cymosum*)

*Claytonia parviflora* ssp. *parviflora* (subsp. *utahensis*)

*Collinsia bartsifolia* var. *davidsonii* (var. *stricta*)

*Collinsia heterophylla* (var. *heterophylla*)

*Conyza bilbaoana* (*Erigeron sumatrensis*)

*Conyza bonariensis* (*Erigeron*)

*Conyza canadensis* (*Erigeron*)

*Conyza coulteri* (*Laennecia*)

*Corallorhiza maculata* (var. *maculata*)

*Cordylanthus mollis* ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

*Coronopus didymus* (*Lepidium didymum*)

*Corylus cornuta* var. *californica* (subsp. *californica*)

*Cotoneaster pannosa* (*C. pannosus*)

*Cryptantha decipiens* (*C. rattanii*)

*Cryptantha intermedia* (var. *intermedia*)

*Cryptantha muricata* (var. *muricata*)

*Cryptantha torreyana* (var. *pumila*)

*Cupressus macrocarpa* (*Hesperocyparis*)

*Cuscuta salina* var. *major* (*C. pacifica* var. *pacifica*)

*Cynara cardunculus* (subsp. *flavescens*)

*Danthonia californica* var. *californica* (no var.)

*Dicentra chrysantha* (*Ehrendorferia*)

*Disporum hookeri* (*Prosartes*)

*Disporum smithii* (*Prosartes*)

*Draba cuneifolia* var. *integrifolia* (no var.)

*Duchesnea indica* (no var.)

*Epilobium pygmaeum* (*E. campestre*)

*Eragrostis curvula* var. *curvula* (no var.)

*Erechtites glomerata* (*Senecio glomeratus*)

*Erechtites minima* (*Senecio minimus*)

*Eremocarpus setigerus* (*Croton*)

*Erigeron philadelphicus* (var. *philadelphicus*)

*Eriogonum umbellatum* var. *bahiiforme* (var. *smallianum*)

*Eriophyllum lanatum* var. *achillaeoides* (var. *achilleoides*)

*Eryngium aristulatum* var. *aristulatum* (*E. jepsonii*)

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

*Erysimum capitatum* var. *angustatum* (var. *capitatum*)

*Filago californica* (*Logfia filaginoides*)

*Filago gallica* (*Logfia*)

*Fritillaria affinis* var. *affinis* (no var.)

*Galium trifidum* var. *pacificum* (subsp. *columbianum*)

*Gastridium ventricosum* (*G. phleoides*)

*Gnaphalium bicolor* (*Pseudognaphalium biolettii*)

*Gnaphalium californicum* (*Pseudognaphalium*)

*Gnaphalium canescens* ssp. *beneolens*  
(*Pseudognaphalium beneolens*)

*Gnaphalium canescens* ssp. *microcephalum*  
(*Pseudognaphalium microcephalum*)

*Gnaphalium luteo-album* (*Pseudognaphalium luteoalbum*)

*Gnaphalium purpureum* (*Gamochaeta ustulata*)

*Gnaphalium ramosissimum* (*Pseudognaphalium*)

*Gnaphalium stramineum* (*Pseudognaphalium*)

*Grindelia camporum* var. *camporum* (no var.)

*Grindelia hirsutula* var. *davyi* (*G. camporum*)

*Grindelia hirsutula* var. *hirsutula* (no var.)

*Guillenia flavescens* (*Caulanthus*)

*Guillenia lasiophylla* (*Caulanthus lasiophyllus*)

*Hedera helix* ssp. *helix* (no subsp.)

*Heliotropium curassavicum* (var. *oculatum*)

*Hemizonia congesta* ssp. *congesta* (subsp. *lutescens*)

*Hemizonia corymbosa* ssp. *corymbosa* (*Deinandra corymbosa*)

*Hemizonia fitchii* (*Centromadia*)

*Hemizonia lobbii* (*Deinandra*)

*Hemizonia parryi* ssp. *congdonii* (*Centromadia*)

*Hemizonia parryi* ssp. *parryi* (*Centromadia*)

*Hemizonia pungens* ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

*Hemizonia pungens* ssp. *pungens* (*Centromadia*)

*Heracleum lanatum* (*H. maximum*)

*Herniaria hirsuta* ssp. *cinerea* (var. *cinerea*)

*Holodiscus discolor* (var. *discolor*)

*Hordeum jubatum* (subsp. *jubatum*)

*Horkelia californica* ssp. *californica* (var. *californica*)

*Horkelia californica* ssp. *dissita* (var. *elata*)

*Horkelia californica* ssp. *frondosa* (var. *frondosa*)

*Isopyrum occidentale* (*Enemion*)

*Iva axillaris* ssp. *robustior* (no subsp.)

*Juglans californica* var. *hindsii* (*J. hindsii*)

*Juncus balticus* (subsp. *ater*)

*Juncus effusus* var. *pacificus* (subsp. *pacificus*)

*Juncus lesueurii* (*J. lescurii*)

*Koeleria phleoides* (*K. gerardii*)

*Lagophylla ramosissima* ssp. *ramosissima* (no subsp.)

*Lasthenia californica* (subsp. *californica*)

*Lavatera cretica* (*Malva pseudolavatera*)

*Lepidium dictyotum* var. *dictyotum* (no var.)

*Lepidium latipes* var. *latipes* (no var.)

*Lepidium nitidum* var. *nitidum* (no var.)

*Lepidium nitidum* var. *oreganum* (no var.)

*Leptochloa fascicularis* (*L. fusca* subsp. *fascicularis*)

*Lessingia filaginifolia* var. *californica* (*Corethrogyne filaginifolia*)

*Lessingia filaginifolia* var. *filaginifolia*  
(*Corethrogyne filaginifolia*)

*Lewisia rediviva* (var. *rediviva*)

*Leymus condensatus* (*Elymus*)

*Leymus triticoides* (*Elymus*)

*Lilaea scilloides* (*Triglochin*)

*Limosella subulata* (*Limosella australis*)

*Linanthus acicularis* (*Leptosiphon*)

*Linanthus ambiguus* (*Leptosiphon*)

*Linanthus androsaceus* (*Leptosiphon*)

*Linanthus bicolor* (*Leptosiphon*)

*Linanthus ciliatus* (*Leptosiphon*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus parviflorus* (*Leptosiphon*)

*Linaria canadensis* (*Nuttallanthus texanus*)

*Lolium multiflorum* (*Festuca perennis*)

*Lolium perenne* (*Festuca perennis*)

*Lolium temulentum* (*Festuca temulenta*)

*Lonicera hispidula* var. *vacillans* (no var.)

*Lotus humistratus* (*Acmispon brachycarpus*)

*Lotus micranthus* (*Acmispon parviflorus*)

*Lotus purshianus* var. *purshianus* (*Acmispon americanus* var. *americanus*)

*Lotus scoparius* var. *scoparius* (*Acmispon glaber* var. *glaber*)

*Lotus strigosus* (*Acmispon*)

*Lotus wrangelianus* (*Acmispon*)

*Luzula comosa* (var. *comosa*)

*Madia madioides* (*Anisocarpus*)

*Malacothamnus fasciculatus* (var. *nuttallii*)

*Marah fabaceus* (*M. fabacea*)

*Marah oreganus* (*M. oregana*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Melilotus alba* (*M. albus*)

*Melilotus indica* (*M. indicus*)

*Mentha spicata* var. *spicata* (no var.)

*Mimulus aurantiacus* (var. *aurantiacus*)

*Mirabilis californica* (*M. laevis* var. *crassifolia*)

*Monardella antonina* ssp. *antonina* (*M. villosa* subsp. *villosa*)

*Monardella douglasii* ssp. *douglasii* (no subsp.)

*Monardella villosa* ssp. *globosa* (subsp. *villosa*)

*Montia fontana* ssp. *chondrosperma* (no subsp.)

*Myrica californica* (*Morella*)

*Nassella cernua* (*Stipa*)

*Nassella lepida* (*Stipa*)

*Nassella pulchra* (*Stipa*)

*Osmorhiza chilensis* (*O. berteroi*)

*Oxalis albicans* ssp. *pilosa* (*O. pilosa*)

*Parvisedum pentandrum* (*Sedella pentandra*)

*Phacelia ramosissima* var. *latifolia* (no var.)

*Phacelia ramosissima* var. *ramosissima* (no var.)

*Phlox gracilis* (*Microsteris*)

*Phoradendron macrophyllum* (*P. serotinum* subsp. *macrophyllum*)

*Phoradendron villosum* (*P. serotinum* subsp. *tomentosum*)

*Phyla nodiflora* var. *incisa* (no var.)

*Phyla nodiflora* var. *nodiflora* (no var.)

*Picris echioides* (*Helminthotheca*)

*Piperia elegans* (subsp. *elegans*)

*Piptatherum miliaceum* (*Stipa miliacea* var. *miliacea*)

*Plagiobothrys canescens* (var. *canescens*)

*Plagiobothrys fulvus* (var. *campestris*)

*Plectritis brachystemon* (*P. congesta* subsp. *brachystemon*)

*Plectritis ciliosa* ssp. *ciliosa* (no subsp.)

*Plectritis ciliosa* ssp. *insignis* (no subsp.)

*Plectritis congesta* (subsp. *congesta*)

*Pleuropogon californicus* (var. *californicus*)

*Pluchea odorata* (var. *odorata*)

*Poa bulbosa* (subsp. *vivipara*)

*Polygonum amphibium* var. *emersum* (*Persicaria amphibia*)

*Polygonum amphibium* var. *stipulaceum* (*Persicaria amphibia*)

*Polygonum arenastrum* (*P. aviculare* subsp. *depressum*)

*Polygonum hydropiperoides* (*Persicaria*)

*Polygonum lapathifolium* (*Persicaria lapathifolia*)

*Polygonum persicaria* (*Persicaria maculosa*)

*Polygonum punctatum* (*Persicaria punctata*)

*Populus balsamifera* ssp. *trichocarpa* (*P. trichocarpa*)

*Potamogeton pectinatus* (*Stuckenia pectinata*)

*Potentilla glandulosa* ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

*Psilocarphus tenellus* var. *tenellus* (no var.)

*Ranunculus aquatilis* var. *capillaceus* (var. *diffusus*)

*Ranunculus californicus* (var. *californicus*)

*Ranunculus canus* (var. *canus*)

*Ranunculus occidentalis* (var. *occidentalis*)

*Ranunculus sceleratus* (var. *sceleratus*)

*Rhamnus californica* ssp. *californica* (*Frangula*)

*Rhamnus tomentella* ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

*Rhus trilobata* (*R. aromatica*)

*Ribes menziesii* (var. *menziesii*)

*Rorippa nasturtium-aquaticum* (*Nasturtium officinale*)

*Rorippa palustris* var. *occidentalis* (subsp. *palustris*)

*Rosa gymnocarpa* (var. *gymnocarpa*)

*Rubus discolor* (*R. armeniacus*)

*Rumex maritimus* (*R. fueginus*)

*Rumex salicifolius* var. *crassus* (*R. crassus*)

*Rumex salicifolius* var. *salicifolius* (no var.)

*Rumex salicifolius* var. *transitorius* (*R. transitorius*)

*Salicornia europaea* (*S. depressa*)

*Salicornia subterminalis* (*Arthrocnemum subterminale*)

*Salicornia virginica* (*S. pacifica*)

*Salix exigua* (var. *hindsiana*)

*Salix lucida* ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

*Sambucus mexicana* (*S. nigra* subsp. *caerulea*)

*Satureja douglasii* (*Clinopodium*)

*Saxifraga californica* (*Micranthes*)

*Scirpus acutus* var. *occidentalis* (*Schoenoplectus*)

*Scirpus americanus* (*Schoenoplectus*)

*Scirpus californicus* (*Schoenoplectus*)

*Scirpus cernuus* (*Isolepis*)

*Scirpus maritimus* (*Bolboschoenus maritimus* subsp. *paludosus*)

*Scirpus robustus* (*Bolboschoenus*)

*Scrophularia californica* ssp. *californica* (no subsp.)

*Scrophularia californica* ssp. *floribunda* (no subsp.)

*Senecio breweri* (*Packera*)

*Senecio mikanioides* (*Delairea odorata*)

*Silene californica* (*S. laciniata* subsp. *californica*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Smilacina racemosa* (*Maianthemum racemosum*)

*Smilacina stellata* (*Maianthemum stellatum*)

*Solidago californica* (*S. velutina* subsp. *californica*)

*Solidago canadensis* ssp. *elongata* (*S. elongata*)

*Sparganium erectum* ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

*Sparganium eurycarpum* ssp. *eurycarpum* (var. *eurycarpum*)

*Spergula arvensis* ssp. *arvensis* (no subsp.)

*Spergularia media* ( var. *media*)

*Stachys ajugoides* var. *ajugoides* (no var.)

*Stachys ajugoides* var. *rigida* (*Stachys rigida* var. *quercetorum*)

*Streptanthus albidus* ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

*Stylomecon heterophylla* (*Papaver heterophyllum*)

*Taeniatherum caput-medusae* (*Elymus*)

*Thysanocarpus laciniatus* (var. *laciniatus*)

*Tolmiea menziesii* (*T. diplomenziesii*)

*Trifolium albopurpureum* var. *albopurpureum* (no var.)

*Trifolium albopurpureum* var. *dichotomum* (*T. dichotomum*)

*Trifolium albopurpureum* var. *olivaceum* (*T. olivaceum*)

*Trifolium barbigerum* var. *andrewsii* (*T. grayi*)

*Trifolium barbigerum* var. *barbigerum* (no var.)

*Trifolium gracilentum* var. *gracilentum* (no var.)

*Vicia americana* var. *americana* (subsp. *americana*)

*Vicia ludoviciana* var. *ludoviciana* (subsp. *ludoviciana*)

*Viola adunca* (var. *adunca*)

*Vulpia bromoides* (*Festuca*)

*Vulpia microstachys* var. *ciliata* (*Festuca microstachys*)

*Vulpia microstachys* var. *confusa* (*Festuca microstachys*)

*Vulpia microstachys* var. *microstachys* (*Festuca microstachys*)

*Vulpia microstachys* var. *pauciflora* (*Festuca microstachys*)

*Vulpia myuros* var. *hirsuta* (*Festuca myuros*)

*Vulpia myuros* var. *myuros* (*Festuca myuros*)

*Vulpia octoflora* var. *hirtella* (*Festuca octoflora*)

*Vulpia octoflora* var. *octoflora* (*Festuca octoflora*)

*Zigadenus fremontii* (*Toxicoscordion*)