

Del Valle Regional Park

Checklist of Wild Plants

Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Del Valle Regional Park. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Family	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

Del Valle Regional Park Checklist of Wild Plants

Ferns & Horsetails		Grouped by Growth Form				Ferns & Horsetails
<i>Adiantum</i>		Sorted Alphabetically by Scientific Name				<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	- - -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	Western Lady Fern	Woods-	- - -	- NP	Woodland, along streams, seepage area -	
<input type="radio"/> <i>Cheilanthes intertexta</i>	Coastal Lip Fern	Pterid-	- - A2	- NP	Crevices, bases of rocks -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	- - -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum laevigatum</i>	Smooth Scouring Rush	Equiset-	- - B	- NP	Moist, sandy or gravelly areas -	
<input type="radio"/> <i>Equisetum</i> sp.	Scouring Rush	Equiset-	- - -	- NP	- -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	- - -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Pterid-	- - C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Pterid-	- - -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	- - -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Pteridium aquilinum</i> var. <i>pubescens</i>	Bracken Fern	Dennstaedti-	- - -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>		
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Agrostis avenacea</i>	Pacific Bent Grass	Po-	- - -	L	ZP	Open, often disturbed place (Jun-Jul)	
<input type="radio"/> <i>Agrostis exarata</i>	Spike Bent Grass	Po-	- - -	-	NP	Common. Moist or disturbed areas, open woodland, conifer forest (Jun-Aug)	
<input type="radio"/> <i>Agrostis stolonifera</i>	Creeping Bent Grass	Po-	- - -	L	ZP	Ditches, lake margins, marshes (Jun-Sep)	
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	- - -	-	ZA	Sandy soils, open or disturbed sites (Apr-Jun)	
<input type="radio"/> <i>Arundo donax</i>	Giant Reed	Po-	- - -	H	ZP	Moist places, seeps, ditchbanks (Mar-Sep)	
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	- - -	M	ZA	Disturbed sites (Mar-Jun)	
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	- - -	M	ZA	Disturbed sites (Apr-Jun)	
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	- - -	-	ZA	Shaded or moist, open sites (Apr-Jul)	
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	- - -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)	
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	- - -	M	ZA	Open, gen disturbed areas (Apr-Jul)	
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	- - -	L	ZA	Fields, disturbed areas (Apr-Jul)	
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>madritensis</i>	Foxtail Chess	Po-	- - -	-	ZA	Disturbed areas, roadsides (Apr-Jan)	
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	- - -	H	ZA	Disturbed areas, roadsides (Mar-Jun)	
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	- - B	-	NP	Seasonally wet places (May-Aug)	
<input type="radio"/> <i>Carex</i> sp.	Sedge	Cyper-	- - -	-	--		
<input type="radio"/> <i>Crypsis schoenoides</i>	Swamp Prickle Grass	Po-	- - -	-	ZA	Wet places (Jun-Oct)	
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Po-	- - -	M	ZP	Disturbed sites (Jun-Aug)	
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	- - -	M	ZA	Open, disturbed sites (May-Jul)	
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	- - -	-	NP	Vernal pools, streambanks (May-Nov)	
<input type="radio"/> <i>Cyperus odoratus</i>	Coarse Nutsedge	Cyper-	- - A1	-	NP	Wet disturbed soils (Jul-Oct)	
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Po-	- - -	-	NP	Salt marshes, coastal dunes, moist, alkaline areas (Apr-Sep)	
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	- - -	-	NP	Common. Fresh to brackish wetland (Spring-summer)	
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Po-	- - -	-	NP	Open areas, chaparral, woodland, forest (Jun-Aug)	
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i>)	Beardless Wild Rye	Po-	- - -	-	NP	Dry to moist, often saline, meadows (Jun-Jul)	
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	- - -	M	ZP	Disturbed places (May-Jun)	
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i>)	Brome Fescue	Po-	- - -	-	ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May-Jun)	
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	- - C	-	NP	Dry, open forest, moist streambanks, chaparral (May-Jun)	
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i>)	Hairy Fescue	Po-	- - -	-	NA	Disturbed, open, gen sandy soils (Apr-Jun)	

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like
<i>Festuca</i>	Sorted Alphabetically by Scientific Name				<i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca octoflora</i> (JM93: <i>Vulpia</i> , var. <i>hirtella</i> , <i>octoflora</i>)	Slender Six-weeks Fescue	Po-	- - A2	- NA	Sandy to rocky soils, open sites (Mar–Jun)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , L. <i>multiflorum</i>)	Rye Grass	Po-	- - -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	- - -	- NA	- -
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	- - -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	- - -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	- - -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>murinum</i>	Wall Barley	Po-	- - -	- ZA	Moist, gen disturbed sites (Feb–May)
<input type="radio"/> <i>Juncus balticus</i> subsp. <i>ater</i> (JM93: no subsp.)	Baltic Rush	Junc-	- - -	- NP	Moist to ± dry sites (Jul–Nov)
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Junc-	- - -	- NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus dubius</i>	Mariposa Rush	Junc-	- - A1	- NP	Wet places (Jul–Sep)
<input type="radio"/> <i>Juncus mexicanus</i>	Mexican Rush	Junc-	- - B	- NP	Common. Coast to montane meadows (Aug–Nov)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	- - -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	- - -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	- - C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Lamarckia aurea</i>	Goldentop	Po-	- - -	- ZA	Open ground, moist seeps, rocky hillsides, sandy soil (Feb–May)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	- - C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	- - -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	- - -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Panicum capillare</i>	Witch Grass	Po-	- - B	- NA	Open places, fields, roadsides (Feb–Dec)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	- - -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa howellii</i>	Howell Blue Grass	Po-	- - B	- NA	Rocky banks, shaded slopes of woodland, chaparral, disturbed places (Apr–Jun)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	- - -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	- - -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Schoenoplectus acutus</i> var. <i>occidentalis</i> (JM93: <i>Scirpus</i>)	Common Tule	Cyper-	- - -	- NP	Common. Marshes, shores, fens, shallow lakes, often emergent (Summer)
<input type="radio"/> <i>Scirpus</i> , <i>Isolepis</i> , or <i>Bolboschoenus</i> sp. (JM93: <i>Scirpus</i> sp.)	Bulrush	Cyper-	- - -	- NAP	

Grasses & Grass-like <i>Stipa</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Stipa</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i>)	Foothill Needle Grass	Po-	- - C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa miliacea</i> var. <i>miliacea</i> (JM93: <i>Piptatherum miliaceum</i>)	Smilo Grass	Po-	- - -	L ZP	Salt marshes, streambanks, chaparral, open woodland, disturbed (Mar–Sep)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Po-	- - C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Trisetum canescens</i>	Tall False Oat	Po-	- - B	- NP	Open to shaded sites, meadows, chaparral, conifer forest (May–Aug)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)

Herbaceous <i>Achillea</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Achillea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	- - -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Aster-	- - -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)	
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Fab-	- - -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Fab-	- - -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon</i> or <i>Hosackia</i> sp. (JM93: <i>Lotus</i> sp.)	Lotus	Fab-	- - -	- -		
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	- - -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Aster-	- - -	- NA	Many open habitats (May–Jun)	
<input type="radio"/> <i>Agoseris retrorsa</i>	Spearleaf Native Dandelion	Aster-	- - B	- NP	Scrub, oak woodland, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	- - -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Alli-	- - C	- NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Boragin-	- - -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Amsinckia menziesii</i> (JM93: var. <i>menziesii</i>)	Small-flowered Fiddleneck	Boragin-	- - -	- NA	Shade-tolerant, open, disturbed areas at forest/woodland edges (May–Jul)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	- - -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	- - -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	- - -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	- - -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Artemisia dracunculus</i>	Wild Tarragon	Aster-	- - A1	- NP	Common. Many habitats, especially disturbed sites (Aug–Oct)	
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	- - -	- NP	Dry ground, valleys, foothills (May–Oct)	
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	- - -	- NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	- - -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Bellardia</i>	Sorted Alphabetically by Scientific Name					<i>Bellardia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	- - -	L	ZA	Disturbed grassland. (Apr–Jun)
<input type="radio"/> <i>Blepharizonia laxa</i> (JM93: <i>B. plumosa</i> subsp. <i>viscida</i>)	Stickyleaf Big Tarweed	Aster-	- - A2	-	NA	Openings in woodland, chaparral, grassland (Jul–Nov)
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	- - -	-	NA	Shade of trees, rocks, shrubs (Mar–Apr)
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	- - -	M	ZA	Common. Disturbed areas, fields (Apr–Sep)
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	- - -	L	ZA	Disturbed areas (Jan–May)
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	- - -	-	NP	Common. Shady to open woodland, scrub (Apr–Jun)
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	- - C	-	NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	- - B	-	NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calystegia purpurata</i> subsp. <i>purpurata</i>	Climbing Morning-glory	Convolvul-	- - -	-	NP	Chaparral, coastal scrub (May–Jun)
<input type="radio"/> <i>Calystegia subacaulis</i> subsp. <i>subacaulis</i>	Shortstem Morning-glory	Convolvul-	- - -	-	NP	Dry, open scrub or woodland (Apr–Jun)
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	- - -	-	ZA	Disturbed areas (Jan–Oct)
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	- - -	-	NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)
<input type="radio"/> <i>Carduus pycnocephalus</i> subsp. <i>pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	- - -	M	ZA	Roadsides, pastures, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Castilleja affinis</i> subsp. <i>affinis</i>	Common Indian Paintbrush	Orobanch-	- - -	-	NP	Chaparral, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Castilleja exserta</i> subsp. <i>exserta</i>	Purple Owl's-clover	Orobanch-	- - -	-	NA	Open fields, grassland (Mar–May)
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i>)	California Mustard	Brassic-	- - -	-	NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	- - -	M	ZAB	Pastures, disturbed places (Apr–Nov)
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	- - -	H	ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	- - -	-	ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chamaesyce serpyllifolia</i> subsp. <i>serpyllifolia</i>	Thyme-leafed Spurge	Euphorbi-	- - B	-	NA	Common. Dry habitats (Aug–Oct)
<input type="radio"/> <i>Chenopodium album</i>	Lamb's Quarters	Chenopodi-	- - -	-	ZA	Common. Disturbed areas, fields (Jun–Oct)
<input type="radio"/> <i>Chenopodium californicum</i>	California Goosefoot	Chenopodi-	- - -	-	NP	Generally open sites, sandy to clay soils (Mar–Sep)
<input type="radio"/> <i>Chenopodium</i> sp.	Goosefoot	Chenopodi-	- - -	-	--	
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	- - -	-	NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Cirsium brevistylum</i>	Indian Thistle	Aster-	- - B	-	NA	Moist places (Mar–Aug)

Herbaceous <i>Cirsium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Cirsium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	- - -	M ZB	Common. Disturbed areas (May–Oct)	
<input type="radio"/> <i>Clarkia affinis</i>	Small Clarkia	Onagr-	- - C	- NA	Openings in woodland, chaparral (May–Jun)	
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	- - -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)	
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	- - -	- NA	--	
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	- - -	- NA	Common. Woodland (Apr–Sep)	
<input type="radio"/> <i>Claytonia parviflora</i> subsp. ?	Miner's Lettuce	Monti-	- - -	- NA	--	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	- - -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)	
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	- - -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)	
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>sparsiflora</i>	Few-flowered Collinsia	Plantagin-	- - C	- NA	Grassy, sometimes disturbed or rocky places, chaparral, oak woodland, dry mixed woodland (Mar–May)	
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	- - -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)	
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	- - -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)	
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	- - -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)	
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i>)	Turkey-mullein	Euphorbi-	- - -	- NA	Dry, open, often disturbed areas (May–Oct)	
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	- - -	- NP	Chaparral, woodland (Feb–May)	
<input type="radio"/> <i>Datisca glomerata</i>	Durango Root	Datisc-	- - A2	- NP	Dry streambeds or washes (May–Jul)	
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>pallidum</i>	Pale-flowered Western Larkspur	Ranuncul-	- - C	- NP	Oak woodland, e slope coast ranges (Mar–May)	
<input type="radio"/> <i>Delphinium variegatum</i> subsp. <i>variegatum</i>	Royal Larkspur	Ranuncul-	- - C	- NP	Grassland, open oak woodland (Mar–May)	
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	- - -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	- - -	- NP	Open woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Dichelostemma</i> sp.	Blue Dicks/ookow	Themid-	- - -	- NP	--	
<input type="radio"/> <i>Dipsacus sativus</i>	Fuller's Teasel	Dipsac-	- - -	M ZB	Disturbed areas, fields, vacant lots, pastures (May–Jul)	
<input type="radio"/> <i>Dittrichia graveolens</i>	Stinkwort	Aster-	- - -	M ZA	Disturbed areas (Sep–Nov)	
<input type="radio"/> <i>Dodecatheon clevelandii</i> subsp. <i>patulum</i>	Padre Shooting Star	Primul-	- - C	- NP	Moist places, often on serpentine or in ± alkaline sites (Mar–May)	
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	- - -	- NP	Gen in shady sites (Mar–Jul)	
<input type="radio"/> <i>Echinodorus berteroi</i>	Upright Burhead	Alismat-	- - A2	- NA	Ponds, ditches (Mid-summer–fall)	
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	- - -	- NP	Dry slopes, ridges (Jun–Dec)	

Herbaceous <i>Epilobium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Epilobium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Epilobium densiflorum</i>	Dense-flower Willowherb	Onagr-	- - C	- NA	Streambanks, outwashes, seasonal moist flats (May–Oct)
<input type="radio"/> <i>Epilobium minutum</i>	Chaparral Willowherb	Onagr-	- - -	- NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)
<input type="radio"/> <i>Eremothera boothii</i> subsp. <i>decorticans</i> (JM93: <i>Camissonia</i>)	Shredding Evening-primrose	Onagr-	- - A1	- NA	Open, generally steep and rocky, especially shale slopes (Mar–Jun)
<input type="radio"/> <i>Erigeron bonariensis</i> (JM93: <i>Conyza</i>)	Flax-leaved Horseweed	Aster-	- - -	- ZA	Disturbed sites (All year)
<input type="radio"/> <i>Erigeron canadensis</i> (JM93: <i>Conyza</i>)	Horseweed	Aster-	- - -	- ZA	Disturbed places (All year)
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	- - A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Eriogonum roseum</i>	Wand Wild Buckwheat	Polygon-	- - B	- NA	Common. Sand or gravel (May–Nov)
<input type="radio"/> <i>Eriogonum</i> sp.	Wild Buckwheat	Polygon-	- - -	- NAP - -	
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	- - -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	- - -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	- - -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Euphorbia lathyris</i>	Caper Spurge	Euphorbi-	- - -	- ZAB	Disturbed areas (All year)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	- - -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i>)	Broadleaf Aster	Aster-	- - -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)
<input type="radio"/> <i>Euthamia occidentalis</i>	Western Goldenrod	Aster-	- - -	- NP	Marshes, streambanks, meadows (Jul–Nov)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	- - -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	- - -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	- - -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	- - -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	- - -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Galium</i> sp.	Bedstraw	Rubi-	- - -	- - -	
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	- - -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	- - -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>achilleifolia</i>	California Gilia	Polemoni-	- - B	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)
<input type="radio"/> <i>Githopsis diffusa</i> subsp. <i>robusta</i>	Southern Bluecup	Campanul-	- - A2	- NA	Shaded or disturbed areas, burns (Apr–Jun)
<input type="radio"/> <i>Glycyrrhiza lepidota</i>	Wild Licorice	Fab-	- - C	- NP	In colonies, moist, gen open, disturbed areas incl streambanks, roadsides, alkaline soils or not (May–Jul)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Grindelia</i>	Sorted Alphabetically by Scientific Name					<i>Grindelia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i>)	Great Valley Gumplant	Aster-	- - -	- NP	Sandy or saline bottomland, roadsides (May–Nov)	
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	- - -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)	
<input type="radio"/> <i>Helianthus californicus</i>	California Sunflower	Aster-	- - C	- NP	Springs, marshes, streambanks, canyons (Jul–Oct)	
<input type="radio"/> <i>Heliotropium curassavicum</i> var. <i>oculatum</i> (JM93: no var.)	Seaside Heliotrope	Boragin-	- - -	- NP	Moist to dry, saline to alkaline soils, gen near water (Feb–Oct)	
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i>)	Bristly Ox-tongue	Aster-	- - -	L ZAB	Common. Disturbed areas (All year)	
<input type="radio"/> <i>Heterotheca oregona</i> var. <i>rudis</i>	Inland Oregon Goldenaster	Aster-	- - A1	- NP	Seasonally dry streambed (Jul–Oct)	
<input type="radio"/> <i>Heterotheca oregona</i> var. <i>scaberrima</i>	Sticky Oregon Goldenaster	Aster-	- - A1	- NP	Seasonally dry streambeds (Jul–Oct)	
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	- - -	M ZBP	Disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Hoita macrostachya</i>	Leather Root	Fab-	- - B	- NP	Streamsides, marshes, spring-moist places (May–Aug)	
<input type="radio"/> <i>Holocarpha heermannii</i>	Heermann Tarplant	Aster-	- - -	- NA	Grassland (May–Nov)	
<input type="radio"/> <i>Holocarpha obconica</i>	San Joaquin Tarplant	Aster-	- - B	- NA	Grassland (Apr–Nov)	
<input type="radio"/> <i>Holocarpha virgata</i> subsp. <i>virgata</i>	Sticky Tarplant	Aster-	- - -	- NA	Grassland (May–Nov)	
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	- - -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	- - -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)	
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	- - -	- ZA	Roadsides, grassland (Jul–Nov)	
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	- - -	- ZA	Abundant. Disturbed places (May–Oct)	
<input type="radio"/> <i>Laennecia coulteri</i> (JM93: <i>Conyza</i>)	Coulter's Horseweed	Aster-	- - A1	- NA	Disturbed sites, clayey or sandy soils, often seasonally wet, alkaline (May–Nov)	
<input type="radio"/> <i>Lamium amplexicaule</i>	Clasping Henbit	Lami-	- - -	- ZAB	Disturbed sites, cult or abandoned fields (Apr–Sep)	
<input type="radio"/> <i>Lasthenia californica</i> subsp. <i>californica</i> (JM93: no subsp.)	California Goldfields	Aster-	- - -	- NA	Many habitats (Feb–Jun)	
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	- - -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)	
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	- - C	- NA	Many habitats (Feb–Jul)	
<input type="radio"/> <i>Lemna minor</i>	Common Lesser Duckweed	Lemn-	- - -	- NP	Common. Freshwater (Aug)	
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i>)	Threadleaf Peppergrass	Brassic-	- - -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
<input type="radio"/> <i>Lepidium oxycarpum</i>	Sharp-pod Peppergrass	Brassic-	- - B	- NA	Fields, vernal pool margins, alkaline flats (Mar–May)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Leptosiphon</i>	Sorted Alphabetically by Scientific Name					<i>Leptosiphon</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i>)	Pinklobe Leptosiphon	Polemoni-	- - C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)	
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i>)	Bicolor Leptosiphon	Polemoni-	- - -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Leptosiphon ciliatus</i> (JM93: <i>Linanthus</i>)	Whisker Brush	Polemoni-	- - B	- NA	Common. Open or wooded areas (Mar–Jul)	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	- - -	- NP	Open, grassy slopes (Mar–Apr)	
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	- - -	- NP	Shaded slopes (Feb–Jun)	
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i>)	California Fluffweed	Aster-	- - -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Aster-	- - -	- ZA	Bare or grassy openings, burns (Mar–Jul)	
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Api-	- - B	- NP	Woodland, brushy slopes (Apr–Jun)	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	- - -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lotus corniculatus</i>	Bird's-foot Trefoil	Fab-	- - -	- ZP	Open, disturbed areas (Jun–Sep)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i>)	California Man-root	Cucurbit-	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marah oregana</i> (JM93: <i>M. oreganus</i>)	Coast Man-root	Cucurbit-	- - C	- NP	Shrubby or open areas, forest edges (Mar–May)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i>)	Pineapple Weed	Aster-	- - -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago arabica</i>	Spotted Bur Clover	Fab-	- - -	- ZA	Disturbed and agricultural areas, fields, woodland (Mar–Jun)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus albus</i> (JM93: <i>M. alba</i>)	White Sweetclover	Fab-	- - -	- ZAB	Locally abundant. Pastures, open disturbed sites (May–Sep)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Fab-	- - -	- ZAB	Open, disturbed areas (Apr–Oct)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Mentha</i>	Sorted Alphabetically by Scientific Name					<i>Mentha</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Mentha spicata</i> (JM93: var. <i>spicata</i>)	Spearmint	Lami-	- - -	- ZP	Moist places, marshes, lakeshores, fields (Jul–Oct)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i>)	California Saxifrage	Saxifrag-	- - -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Aster-	- - -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris acuminata</i>	Sierra Foothill Silverpuffs	Aster-	- - B	- NA	Grassy, open rocky or clay soil (Apr–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	- - -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i>)	Slender Annual Phlox	Polemoni-	- - -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	- - -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Mimulus pilosus</i>	Downy Monkeyflower	Phrym-	- - A2	- NA	Moist, sandy areas, especially by small streams, disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>M. globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i>)	Coyote-mint	Lami-	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Najas marina</i>	Holly-leaved Water-nymph	Hydrocharit-	- - -	- NA	Ponds, lakes, marshes, rivers (Jul–Aug)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i>)	Water Cress	Brassic-	- - -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	- - C	- NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)	
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Polemoni-	- - C	- NA	Open, slopes, gravel, clay (May–Jul)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	- - -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila pedunculata</i>	Meadow Nemophila	Boragin-	- - B	- NA	Common. Ocean bluffs, grassland, slopes, meadows, sandbars, fields, woodland, streambanks (Feb–Jul)	
<input type="radio"/> <i>Nicotiana attenuata</i>	Coyote Tobacco	Solan-	- - A1	- NA	Open, well-drained slopes (May–Oct)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Api-	- - -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza brachypoda</i>	California Sweet Cicely	Api-	- - B	- NP	Moist ravines, conifer forest, woodland (Mar–May)	
<input type="radio"/> <i>Packera breweri</i> (JM93: <i>Senecio</i>)	Brewer's Ragwort	Aster-	- - B	- NP	Common. Seasonally damp grassland, oak savanna, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i>)	Wind Poppy	Papaver-	- - -	- NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	- - -	- NP	Dry chaparral, oak/pine forest (Mar–May)	

Herbaceous <i>Persicaria</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Persicaria</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Persicaria lapathifolia</i> (JM93: <i>Polygonum lapathifolium</i>)	Willow Weed	Polygon-	- - C	- NA	Moist places, disturbed areas, roadsides, fields (Jun–Oct)	
<input type="radio"/> <i>Persicaria maculosa</i> (JM93: <i>Polygonum persicaria</i>)	Lady's Thumb	Polygon-	- - -	- ZA	Moist disturbed areas, field (Jun–Nov)	
<input type="radio"/> <i>Pholistoma membranaceum</i>	White Fiesta Flower	Boragin-	- - -	- NA	Beaches, bluffs, ravines, wooded slopes, desert washes (Feb–May)	
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>tomentosum</i> (JM93: <i>P. villosum</i>)	Oak Mistletoe	Visc-	- - -	- NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)	
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i>)	Lemon Verbena	Verben-	- - -	- NP	Wet places, pond margins (May–Nov)	
<input type="radio"/> <i>Piperia elegans</i> subsp. <i>elegans</i> (JM93: no subsp.)	Elegant Rein Orchid	Orchid-	- - B	- NP	Generally dry, open sites, scrub, conifer forest (May–Sep)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	- - -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	- - -	- -	- -	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	- - -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	- - -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Plantagin-	- - -	- ZAP	Disturbed areas (Apr–Sep)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: subsp. <i>ciliosa</i> , <i>insignis</i>)	Longspur Plectritis	Valerian-	- - B	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	- - -	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	- - -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i>)	Knotweed	Polygon-	- - -	- ZA	Disturbed places (May–Nov)	
<input type="radio"/> <i>Potamogeton illinoensis</i>	Shining Pondweed	Potamogeton-	- - -	- NP	Lakes, ponds, streams (Jun–Aug)	
<input type="radio"/> <i>Potamogeton nodosus</i>	Long-leaved Pondweed	Potamogeton-	- - -	- NP	Shallow water, lakes, ponds, streams (May–Aug)	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i>)	California Everlasting	Aster-	- - -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i>)	Weedy Cudweed	Aster-	- - -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)	
<input type="radio"/> <i>Pseudognaphalium ramosissimum</i> (JM93: <i>Gnaphalium</i>)	Pink Everlasting	Aster-	- - -	- NB	Dry, open slopes, woodland, sandy fields, dunes (Jul–Sep)	
<input type="radio"/> <i>Psilocarphus tenellus</i> (JM93: var. <i>tenellus</i>)	Slender Woolly-marbles	Aster-	- - -	- NA	Common. Dry, seasonally moist slopes, flats, burns, trails, rarely vernal pools (Mar–Jul)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	- - -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	- - B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Ranunculus</i>	Sorted Alphabetically by Scientific Name					<i>Ranunculus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	- - -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus occidentalis</i> var. <i>occidentalis</i> (JM93: no var.)	Western Buttercup	Ranuncul-	- - A2	- NP	Grassy slopes in meadows or open woodland (Mar–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	- - -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex transitorius</i> (JM93: <i>R. salicifolius</i> var. <i>transitorius</i>)	Ovate Willow Dock	Polygon-	- - -	- NP	Coastal dunes, wet margins of meadows (Apr–Jun)	
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	- - -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Caryophyll-	- - -	- NA	Sandy disturbed areas, river bars, streamsides (Spring–early summer)	
<input type="radio"/> <i>Sagina decumbens</i> subsp. <i>occidentalis</i>	Western Pearlwort	Caryophyll-	- - C	- NA	Dry streams, chaparral, grassy areas, rock outcrops, vernal pools (Spring–early summer)	
<input type="radio"/> <i>Salsola tragus</i>	Russian Thistle	Chenopodi-	- - -	L ZAP	Common. Disturbed places (Jul–Oct)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	- - -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	- - -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	- - -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica</i> , <i>floribunda</i>)	California Figwort	Scrophulari-	- - -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberous Skullcap	Lami-	- - C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	- - -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	- - -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	- - -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	- - -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	- - -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solidago velutina</i> subsp. <i>californica</i> (JM93: <i>S. californica</i>)	California Goldenrod	Aster-	- - -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	- - -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	- - -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	- - -	- ZA	Abundant. Disturbed places (All year)	

Herbaceous <i>Spergularia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Spergularia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	- - -	-	ZAP	Forest, meadows, mud flats, disturbed
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Lami-	- - C	-	NP	Streambanks, springs, pine/oak forest (Jun–Oct)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i>)	Common Rigid Hedge-nettle	Lami-	- - -	-	NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	- - -	-	ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	- - C	-	NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)
<input type="radio"/> <i>Stephanomeria virgata</i> subsp. <i>pleurocarpa</i>	Twiggy Wreath Plant	Aster-	- - C	-	NA	Chaparral openings, grassland (Jun–Nov)
<input type="radio"/> <i>Symphotrichum chilense</i> (JM93: <i>Aster chilensis</i>)	Common California Wild Aster	Aster-	- - -	-	NP	Grassland, salt marshes, disturbed places (Jun–Oct)
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	- - -	-	NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)
<input type="radio"/> <i>Thysanocarpus laciniatus</i> var. <i>laciniatus</i> (JM93: no var.)	Narrowleaf Fringepod	Brassic-	- - B	-	NA	Oak woodland, rocky ridges, slopes, chaparral, washes (Mar–May)
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	- - -	M	ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	- - -	-	ZA	Disturbed places (Apr–Jun)
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i>)	Common Star Lily	Melanthi-	- - -	-	NP	Grassy or wooded slopes, outcrops (Feb–Jun)
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	- - -	-	ZBP	Common. Disturbed places (Mar–Nov)
<input type="radio"/> <i>Trichostema lanceolatum</i>	Vinegar Weed	Lami-	- - -	-	NA	Dry, open, gen disturbed habitats (Jun–Nov)
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i>)	Rancheria Clover	Fab-	- - C	-	NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>bifidum</i>	Pinole Clover	Fab-	- - B	-	NA	Open, grassy areas, oak chaparral, forest (Apr–Jun)
<input type="radio"/> <i>Trifolium dichotomum</i> (JM93: <i>T. albopurpureum</i> var. <i>dichotomum</i>)	Branched Indian Clover	Fab-	- - B	-	NA	Coastal dunes, open slopes, meadows, oak woodland, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i>)	Pinpoint Clover	Fab-	- - -	-	NA	Open, disturbed places, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	- - -	M	ZA	Disturbed areas, roadsides (Apr–May)
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	- - -	-	NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Fab-	- - C	-	NA	Woody or shrubby slopes, roadsides (Mar–Jun)
<input type="radio"/> <i>Trifolium</i> sp.	Clover	Fab-	- - -	-	- -	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	- - -	-	NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	- - -	-	NA	Grassland (Apr–Jun)

Herbaceous <i>Triteleia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Triteleia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	- - -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Brassic-	- - B	- NP	Grassy banks, open fields, roadsides, pastures (Mar–May)	
<input type="radio"/> <i>Turritis glabra</i> (JM93: <i>Arabis</i> , var. <i>glabra</i>)	Tower Mustard	Brassic-	- - C	- NP	Open fields, meadows, slopes (Apr–Jul)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	- - -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urtica dioica</i> subsp. <i>holosericea</i>	Hoary Nettle	Urtic-	- - -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	- - -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbascum thapsus</i>	Woolly Mullein	Scrophulari-	- - -	L ZB	Roadsides, streambanks, disturbed areas (May–Sep)	
<input type="radio"/> <i>Verbena lasiostachys</i> var. ?	Western Vervain	Verben-	- - -	- NP	--	
<input type="radio"/> <i>Verbena lasiostachys</i> var. <i>scabrida</i>	Robust Vervain	Verben-	- - -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Veronica americana</i>	American Brooklime	Plantagin-	- - C	- NP	Common. Moist to wet soil, springs, slow streams, meadows, lakeshores (May–Aug)	
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	- - -	- ZA	Wet, disturbed areas, fields (Feb–May)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>nigra</i>	Narrow-leaved Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>sativa</i>	Spring Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa</i> subsp. <i>varia</i>	Sparsely Hairy Vetch	Fab-	- - -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	- - C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	- - -	- NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	- - C	- NP	Gen shady sites (Mar–Jun)	
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	- - -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))	
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	- - -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Xanthium strumarium</i>	Cocklebur	Aster-	- - -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Api-	- - C	- NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)	

Woody	Grouped by Growth Form					Woody
Acer	Sorted Alphabetically by Scientific Name					Acer
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	- - -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	- - -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	- - -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	- - -	- NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Ros-	- - C	- NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Eric-	- - C	- NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	- - -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	- - -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Baccharis salicifolia</i> subsp. <i>salicifolia</i> (JM93: no subsp.)	Mule Fat	Aster-	- - -	- NP	Riparian woodland, canyon bottoms, disturbed sites, often forming thickets (All year)	
<input type="radio"/> <i>Brickellia californica</i>	California Brickellbush	Aster-	- - B	- NP	Dry, rocky hillsides, canyons, sea bluffs (Jul–Dec)	
<input type="radio"/> <i>Ceanothus cuneatus</i> var. <i>cuneatus</i>	Buckbrush	Rhamn-	- - C	- NP	Sandy to rocky flats, slopes, ridges (Feb–May)	
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	- - C	- NP	Dry, rocky slopes, chaparral (Mar–May)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	- - -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
<input type="radio"/> <i>Ericameria arborescens</i>	Golden-fleece	Aster-	- - A2	- NP	Woodland, open forest, chaparral, especially after fire (Aug–Nov)	
<input type="radio"/> <i>Ericameria linearifolia</i>	Interior Goldenbush	Aster-	- - -	- NP	Dry slopes, valleys, foothill and desert woodland, saltbush and creosote-bush scrub (Mar–May)	
<input type="radio"/> <i>Ericameria nauseosa</i> var. <i>mohavensis</i> (JM93: <i>Chrysothamnus nauseosus</i> subsp. <i>mohavensis</i>)	Mojave Rabbitbrush	Aster-	- - A2	- NP	Common. Dry scrub (Aug–Oct)	
<input type="radio"/> <i>Eriodictyon californicum</i>	California Yerba Santa	Boragin-	- - -	- NP	Slopes, fields, roadsides, woodland, chaparral (Apr–Jul)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	- - -	- NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Eriophyllum jepsonii</i>	Jepson's Woolly Sunflower	Aster-	- 4.3 *A2	- NP	Dry oak woodland (Apr–Jun)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i>)	California Coffee Berry	Rhamn-	- - -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	- - C	- NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	- - -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	

Woody	Grouped by Growth Form					Woody
<i>Holodiscus</i>	Sorted Alphabetically by Scientific Name					<i>Holodiscus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	- - -	- NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Juniperus californica</i>	California Juniper	Cupress-	- - B	- NP	Dry slopes, flats, pinyon/juniper woodland -	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Caprifoli-	- - -	- NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera interrupta</i>	Chaparral Vine Honeysuckle	Caprifoli-	- - -	- NP	Dry slopes, ridges, floodplains, oak woodland, chaparral (Apr–May)	
<input type="radio"/> <i>Lonicera subspicata</i> var. <i>denudata</i>	Southern Honeysuckle	Caprifoli-	- - B	- NP	Chaparral slopes (Jun–Jul)	
<input type="radio"/> <i>Malacothamnus fremontii</i>	Fremont Bush-mallow	Malv-	- - B	- NP	Open chaparral to pine woodland (May–Jul)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	- - -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Nicotiana glauca</i>	Tree Tobacco	Solan-	- - -	M ZP	Open, disturbed flats or slopes (Apr–Aug)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	- - -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Olea europaea</i>	Olive	Ole-	- - -	L ZP	Generally waif, persisting from cultivation (Feb–Jun)	
<input type="radio"/> <i>Pinus coulteri</i>	Coulter Pine	Pin-	- - A2	- NP	Chaparral, lower mixed-conifer, mixed-hardwood forests -	
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	- - -	- NP	Closed-cone-pine forest, oak woodland -	
<input type="radio"/> <i>Pinus sabiniana</i>	Gray Pine	Pin-	- - -	- NP	Foothill woodland, n oak woodland, chaparral, infertile soils in mixed-conifer and hardwood forests -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	- - B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Populus fremontii</i> subsp. <i>fremontii</i>	Fremont Cottonwood	Salic-	- - -	- NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)	
<input type="radio"/> <i>Prunus ilicifolia</i> subsp. <i>ilicifolia</i>	Holly-leaved Cherry	Ros-	- - C	- NP	Canyons, slopes, scrubland, woodland (Apr–May)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	- - A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	- - C	- NP	Dry slopes, chaparral (Feb–Apr)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	- - -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	
<input type="radio"/> <i>Quercus garryana</i> var. <i>garryana</i>	Oregon / Garry Oak	Fag-	- - B	- NP	Slopes, mixed-evergreen or conifer forest (Apr–Jun)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	- - -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	- - B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	- - C	- NP	Chaparral, montane forest (Mar–Jun)	

Woody	Grouped by Growth Form					Woody
Ribes	Sorted Alphabetically by Scientific Name					Ribes
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	- - -	-	NP	Forest openings, woodland (Feb–Mar)
<input type="radio"/> <i>Ribes malvaceum</i> var. <i>malvaceum</i>	Chaparral Currant	Grossulari-	- - B	-	NP	Chaparral, oak woodland (Oct–Apr)
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	- - -	-	NP	Common. Forest openings, chaparral (Feb–Apr)
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	- - -	-	NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	- - -	-	NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	- - -	-	NP	Open, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Salic-	- - -	-	NP	Common. Floodplains, sandy gravel (Apr–May)
<input type="radio"/> <i>Salix laevigata</i>	Red Willow	Salic-	- - -	-	NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	- - -	-	NP	Common. Shores, marshes, meadows, etc (Jan–Jun)
<input type="radio"/> <i>Salix</i> sp.	Willow	Salic-	- - -	-	?P	- -
<input type="radio"/> <i>Salvia mellifera</i>	Black Sage	Lami-	- - -	-	NP	Coastal-sage scrub, lower chaparral (Mar–Jun)
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i>)	Blue Elderberry	Adox-	- - -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	- - -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)
<input type="radio"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	- - -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)
<input type="radio"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	- - -	-	NP	Ridges, slopes, open places in woodland (Apr–May)
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	- - -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)
<input type="radio"/> <i>Umbellularia californica</i>	California Bay	Laur-	- - -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)
<input type="radio"/> <i>Vitis californica</i>	California Wild Grape	Vit-	- - -	-	NP	Streamsides, springs, canyons (May–Jun)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex deweyana ssp. *leptopoda* (*C. leptopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thyrsoiflorus (var. *thyrsoiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria amphibia*)

Polygonum amphibium var. *stipulaceum* (*Persicaria amphibia*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)