

# Garin/Dry Creek Pioneer Regional Park

## Checklist of Wild Plants

### Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Garin/Dry Creek Pioneer Regional Park. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
<b>Scientific Name</b>	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> ( <a href="http://ucjeps.berkeley.edu/IJM.html">ucjeps.berkeley.edu/IJM.html</a> )
<b>(JM93 if different)</b>	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
<b>Common Name</b>	According to <i>JM2</i> and other references (not standardized)
<b>Family</b>	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
<b>Status</b>	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): <b>Federal/California (Fed./Calif.)   California Native Plant Society (CNPS)   East Bay chapter of the CNPS (EBCNPS)</b>  <b>Fed./Calif.:</b> FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare  <b>CNPS</b> (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California  <b>EBCNPS</b> (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
<b>Inv</b>	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
<b>OL</b>	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
<b>Habitat (Bloom)</b>	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, [wlegard@ebparks.org](mailto:wlegard@ebparks.org)). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

# Garin/Dry Creek Pioneer Regional Park

## Checklist of Wild Plants

Ferns & Horsetails		Grouped by Growth Form				Ferns & Horsetails
<i>Adiantum</i>		Sorted Alphabetically by Scientific Name				<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	-   -   -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Athyrium filix-femina var. cyclosorum</i>	Western Lady Fern	Woodsi-	-   -   -	- NP	Woodland, along streams, seepage area -	
<input type="radio"/> <i>Azolla filiculoides</i>	Mosquito Fern	Azoll-	-   -   -	- NP	Common. Ponds, slow streams -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	-   -   -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum telmateia subsp. braunii</i>	Giant Horsetail	Equiset-	-   -   -	- NP	Streambanks, roadside ditches, seepage areas -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	-   -   -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata var. mucronata</i>	Bird's-foot Fern	Pterid-	-   -   C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis subsp. triangularis</i>	Goldenback Fern	Pterid-	-   -   -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	-   -   -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Dryopterid-	-   -   -	- NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -	
<input type="radio"/> <i>Pteridium aquilinum var. pubescens</i>	Bracken Fern	Dennstaedti-	-   -   -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	-   -   -	- NP	Near streams, springs, seeps -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	-   -   -	- NP	Common. Open meadows, woodland, forest, subalpine (Jun–Aug)
<input type="radio"/> <i>Agrostis sp.</i>	Bent Grass	Po-	-   -   -	- -	
<input type="radio"/> <i>Agrostis stolonifera</i>	Creeping Bent Grass	Po-	-   -   -	L ZP	Ditches, lake margins, marshes (Jun–Sep)
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	-   -   -	- ZA	Sandy soils, open or disturbed sites (Apr–Jun)
<input type="radio"/> <i>Arundo donax</i>	Giant Reed	Po-	-   -   -	H ZP	Moist places, seeps, ditchbanks (Mar–Sep)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	-   -   -	M ZA	Disturbed sites (Mar–Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	-   -   -	M ZA	Disturbed sites (Apr–Jun)
<input type="radio"/> <i>Brachypodium distachyon</i>	Purple False-brome	Po-	-   -   -	M ZA	Disturbed areas, dry slopes (Apr–Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	-   -   -	- ZA	Shaded or moist, open sites (Apr–Jul)
<input type="radio"/> <i>Bromus carinatus var. carinatus</i>	California Brome	Po-	-   -   -	- NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr–Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	-   -   -	M ZA	Open, gen disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	-   -   -	L ZA	Fields, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus madritensis subsp. madritensis</i>	Foxtail Chess	Po-	-   -   -	- ZA	Disturbed areas, roadsides (Apr–Jan)
<input type="radio"/> <i>Bromus madritensis subsp. rubens</i>	Red Brome	Po-	-   -   -	H ZA	Disturbed areas, roadsides (Mar–Jun)
<input type="radio"/> <i>Bromus tectorum</i>	Cheat Grass	Po-	-   -   -	H ZA	Open, disturbed areas (May–Aug)
<input type="radio"/> <i>Carex praegracilis</i>	Freeway Sedge	Cyper-	-   -   C	- NP	Common. Often alkaline, ± moist places (Apr–Aug)
<input type="radio"/> <i>Carex sp.</i>	Sedge	Cyper-	-   -   -	- -	
<input type="radio"/> <i>Cortaderia jubata</i>	Hairy Pampas Grass	Po-	-   -   -	H ZP	Disturbed sites, many habitats, esp coastal (Sep–Feb)
<input type="radio"/> <i>Cortaderia selloana</i>	Smooth Pampas Grass	Po-	-   -   -	H ZP	Disturbed sites (Sep–Mar)
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Po-	-   -   -	M ZP	Disturbed sites (Jun–Aug)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	-   -   -	M ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	-   -   -	- NP	Vernal pools, streambanks (May–Nov)
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	-   -   -	L ZP	Disturbed, often moist sites (May–Aug)
<input type="radio"/> <i>Danthonia californica (JM93: var. californica)</i>	California Oat Grass	Po-	-   -   C	- NP	Gen moist meadows, open woodland (Apr–Aug)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	-   -   -	- NP	Common. Fresh to brackish wetland (Spring–summer)
<input type="radio"/> <i>Elymus caput-medusae (JM93: Taeniatherum)</i>	Medusa Head	Po-	-   -   -	H ZA	Disturbed areas (Apr–Jul)
<input type="radio"/> <i>Elymus glaucus subsp. glaucus</i>	Western Wild-rye	Po-	-   -   -	- NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	-   -   C	- NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus triticoides (JM93: Leymus)</i>	Beardless Wild Rye	Po-	-   -   -	- NP	Dry to moist, often saline, meadows ( Jun–Jul)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	-   -   -	M ZP	Disturbed places (May–Jun)

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i> )	Brome Fescue	Po-	-   -   -	- ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i> )	Hairy Fescue	Po-	-   -   -	- NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i> )	Rattail Sixweeks Grass	Po-	-   -   -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i> )	Rye Grass	Po-	-   -   -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	-   -   -	- NA	--
<input type="radio"/> <i>Glyceria leptostachya</i>	Water Manna Grass	Po-	-   -   A1	- NP	Freshwater marshes, lakes (May–Jun)
<input type="radio"/> <i>Holcus lanatus</i>	Common Velvet Grass	Po-	-   -   -	M ZP	Moist sites, roadbanks, cult fields, meadows (Jun–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	-   -   -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	-   -   -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>murinum</i>	Wall Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites (Feb–May)
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Junc-	-   -   -	- NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Junc-	-   -   -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	-   -   -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus</i> sp.	Rush	Junc-	-   -   -	-	--
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	-   -   -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	-   -   C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Lamarckia aurea</i>	Goldentop	Po-	-   -   -	- ZA	Open ground, moist seeps, rocky hillsides, sandy soil (Feb–May)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	-   -   -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	-   -   C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	-   -   -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	-   -   -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Paspalum distichum</i>	Knot Grass	Po-	-   -   -	- ZP	Edges of lakes, ponds, rice fields, wet roadside ditches (Jun–Oct)
<input type="radio"/> <i>Phalaris aquatica</i>	Harding Grass	Po-	-   -   -	M ZP	Disturbed areas, roadsides (Apr–Aug)

Grasses & Grass-like <i>Poa</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Poa</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	-   -   -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	-   -   -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Po-	-   -   -	- ZP	Common. Streambanks (May–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	-   -   -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Schoenoplectus californicus</i> (JM93: <i>Scirpus</i> )	Southern Bulrush	Cyper-	-   -   -	- NP	Common. Brackish to fresh marshes, shores (Spring–summer)
<input type="radio"/> <i>Scirpus</i> , <i>Isolepis</i> , or <i>Bolboschoenus</i> sp. (JM93: <i>Scirpus</i> sp.)	Bulrush	Cyper-	-   -   -	- NAP	
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i> )	Foothill Needle Grass	Po-	-   -   C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa miliacea</i> var. <i>miliacea</i> (JM93: <i>Piptatherum miliaceum</i> )	Smilo Grass	Po-	-   -   -	L ZP	Salt marshes, streambanks, chaparral, open woodland, disturbed (Mar–Sep)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i> )	Purple Needle Grass	Po-	-   -   C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Triglochin scilloides</i> (JM93: <i>Lilaea</i> )	Flowering-quillwort	Juncagin-	-   -   C	- NA	Vernal pools, streams, ponds, lake margins (Mar–Oct)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	-   -   -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	-   -   -	- NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)

Herbaceous <i>Achillea</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Achillea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	-   -   -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i> )	Deerweed	Fab-	-   -   -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i> )	Small-flower Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i> )	California Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Actaea rubra</i>	Baneberry	Ranuncul-	-   -   B	- NP	Deep soils, moist, open to shaded sites, mixed-evergreen or conifer forests (May–Sep)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	-   -   -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Agoseris</i> sp.	Agoseris	Aster-	-   -   -	- NP	- -	
<input type="radio"/> <i>Alisma triviale</i> (JM93: <i>A. plantago-aquatica</i> )	Northern Water Plantain	Alismat-	-   -   B	- NP	Ponds (Spring–fall)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	-   -   -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i> )	Common Fiddleneck	Boragin-	-   -   -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	-   -   -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	-   -   -	- NP	Generally wooded areas (Jun–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	-   -   -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	-   -   -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	-   -   C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)	
<input type="radio"/> <i>Aralia californica</i>	Elk Clover	Arali-	-   -   B	- NP	Moist shade, canyons, stream sides (Jun–Aug)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	-   -   -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	-   -   B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)	
<input type="radio"/> <i>Asclepias speciosa</i>	Showy Milkweed	Apocyn-	-   -   A2	- NP	Many habitats incl fields, roadsides (May–Sep)	
<input type="radio"/> <i>Baccharis glutinosa</i> (JM93: <i>B. douglasii</i> )	Marsh Baccharis	Aster-	-   -   -	- NP	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	-   -   -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	-   -   -	L ZA	Disturbed grassland. (Apr–Jun)	
<input type="radio"/> <i>Berula erecta</i>	Cutleaf Water-parsnip	Api-	-   -   B	- NP	Marshy areas, streams (Jul–Oct)	

Herbaceous	Grouped by Growth Form						Herbaceous
<i>Bowlesia</i>	Sorted Alphabetically by Scientific Name						<i>Bowlesia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	-   -   -	-	NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	-   -   -	M	ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	-   -   -	L	ZA	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Themid-	-   -   -	-	NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	-   -   -	-	NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	-   -   -	-	NP	Common. Shady to open woodland, scrub (Apr–Jun)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	-   -   C	-	NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	-   -   -	-	NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calystegia purpurata subsp. purpurata</i>	Climbing Morning-glory	Convolvul-	-   -   -	-	NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	-   -   -	-	NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	-   -   -	-	ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	-   -   -	-	NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	-   -   -	-	NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	-   -   -	M	ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	-   -   -	-	NP	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	-   -   -	-	NA	Open fields, grassland (Mar–May)	
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	-   -   -	-	NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)	
<input type="radio"/> <i>Castilleja sp.</i>	Paintbrush / Owl's Clover	Orobanch-	-   -   -	-	NAP		
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	-   -   -	M	ZAB	Pastures, disturbed places (Apr–Nov)	
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	-   -   -	M	ZA	Disturbed fields, open woodland (Apr–Jul)	
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	-   -   -	H	ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)	
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	-   -   -	-	ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)	
<input type="radio"/> <i>Chenopodium rubrum var. rubrum</i> (JM93: no var.)	Red Goosefoot	Chenopodi-	-   -   A1	-	NA	Open, saline places, drying mudflats Aug–Oct	
<input type="radio"/> <i>Chlorogalum pomeridianum var. pomeridianum</i>	Wavyleaf Soap Plant	Agav-	-   -   -	-	NP	Common. Open grassland, chaparral, woodland (May–Aug)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Cirsium</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Cirsium</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Cirsium</i> sp.	Thistle	Aster-	-   -   -	-	- -	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	-   -   -	M	ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	-   -   -	-	NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	-   -   -	-	NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	-   -   -	-	NA	- -
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	-   -   -	-	NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	-   -   -	-	NA	Vernally moist, often shady or disturbed sites (Jan–May)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i> )	Yerba Buena	Lami-	-   -   -	-	NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	-   -   C	-	NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	-   -   -	M	ZB	Common. Moist, esp disturbed places (Apr–Jul)
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	-   -   -	-	ZP	Roadsides, open areas in many pl communities (Mar–Oct)
<input type="radio"/> <i>Corethrogyne filaginifolia</i> (JM93: <i>Lessingia</i> , var. <i>filaginifolia</i> )	Common California-aster	Aster-	-   -   C	-	NP	Coastal scrub, chaparral, grassland, foothill woodland, forest (Jul–Nov)
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	-   -   -	-	NA	Open areas (Feb–May)
<input type="radio"/> <i>Crepis vesicaria</i> subsp. <i>taraxacifolia</i>	Dandelion-leaf Hawksbeard	Aster-	-   -   -	-	ZAB	Sandy clearings, hillsides, disturbed places (Feb–Oct)
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i> )	Turkey-mullein	Euphorbi-	-   -   -	-	NA	Dry, open, often disturbed areas (May–Oct)
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	-   -   -	M	ZP	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	-   -   -	-	NP	Chaparral, woodland (Feb–May)
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	-   -   -	-	NA	Rocky or sandy places (Apr–Jun)
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	-   -   -	-	NP	Open woodland, scrub, desert, grassland (Mar–Jun)
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	-   -   -	-	NP	Open woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Dichelostemma</i> sp.	Blue Dicks/ookow	Themid-	-   -   -	-	NP	- -
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i> )	Sticky Cinquefoil	Ros-	-   -   -	-	NP	Gen ± shady or moist areas (May–Jul)
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	-   -   -	-	NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)



Herbaceous <i>Epilobium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Epilobium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	-   -   -	- NP	Dry slopes, ridges (Jun–Dec)
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Onagr-	-   -   -	- NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>watsonii</i>	Watson Willowherb	Onagr-	-   -   B	- NP	Moist coastal bluffs, streamsides, ± disturbed sites (May–Oct)
<input type="radio"/> <i>Epilobium</i> sp.	Willowherb	Onagr-	-   -   -	- NA	--
<input type="radio"/> <i>Erigeron philadelphicus</i> var. <i>philadelphicus</i> (JM93: no var.)	Philadelphia Fleabane	Aster-	-   -   C	- NBP	Streamsides, other moist habitats (May–Jun)
<input type="radio"/> <i>Erigeron sumatrensis</i> (JM93: <i>Conyza bilbaoana</i> )	Tropical Horseweed	Aster-	-   -   -	- ZA	Disturbed sites (All year)
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	-   -   -	- NP	--
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	-   -   A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium brachycarpum</i>	Southern European Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	-   -   -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	-   -   -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	-   -   -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Euphorbia peplus</i>	Petty Spurge	Euphorbi-	-   -   -	- ZA	Common. Disturbed areas (Feb–Aug)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	-   -   -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i> )	Broadleaf Aster	Aster-	-   -   -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	-   -   -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Fragaria vesca</i>	Wood Strawberry	Ros-	-   -   -	- NP	Gen partial shade in forest (Jan–Jul)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i> )	Checker Lily	Lili-	-   -   -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	-   -   -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	-   -   -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	-   -   -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	-   -   -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	-   -   -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	-   -   -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i> )	Great Valley Gumplant	Aster-	-   -   -	- NP	Sandy or saline bottomland, roadsides (May–Nov)

Herbaceous <i>Grindelia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Grindelia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i> )	Hairy Gumplant	Aster-	-   -   C	- NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)
<input type="radio"/> <i>Grindelia</i> sp.	Gumplant	Aster-	-   -   -	- NP	--
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	-   -   -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)
<input type="radio"/> <i>Heliotropium curassavicum</i> var. <i>oculatum</i> (JM93: no var.)	Seaside Heliotrope	Boragin-	-   -   -	- NP	Moist to dry, saline to alkaline soils, gen near water (Feb–Oct)
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i> )	Bristly Ox-tongue	Aster-	-   -   -	L ZAB	Common. Disturbed areas (All year)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. ?	Yellow Hayfield Tarweed	Aster-	-   -   -	- NA	--
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Aster-	-   -   -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i> )	Cow Parsnip	Api-	-   -   -	- NP	Moist places, wooded or open (Apr–Jul)
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	-   -   -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)
<input type="radio"/> <i>Heterotheca grandiflora</i>	Telegraph Weed	Aster-	-   -   -	- NA	Disturbed areas, dry streambeds, sand dunes (Jun–Oct(± all year))
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. <i>bolanderi</i>	Bolander Goldenaster	Aster-	-   -   C	- NP	Dunes, headlands, grassy coastal slopes (Jun–Sep)
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. <i>echioides</i>	Bristly Goldenaster	Aster-	-   -   -	- NP	Grassland, scrub, woodland, open forest, disturbed sites (Jul–Oct)
<input type="radio"/> <i>Heuchera micrantha</i>	Small-flower Alumroot	Saxifrag-	-   -   C	- NP	Moist, rocky banks and cliffs (Apr–Jul)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	-   -   -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Horkelia californica</i> var. <i>frondosa</i> (JM93: subsp. <i>frondosa</i> )	Unlobed California Horkelia	Ros-	-   -   C	- NP	Coastal scrub, canyons, poison-oak thickets (May–Oct)
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	-   -   -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	-   -   -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	-   -   -	- ZA	Abundant. Disturbed places (May–Oct)
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	-   -   -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)
<input type="radio"/> <i>Lathyrus jepsonii</i> var. <i>californicus</i>	Bluff Pea	Fab-	-   -   B	- NP	Forest, open areas (Apr–Aug)
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	-   -   -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Lemna gibba</i>	Swollen Duckweed	Lemn-	-   -   -	- NP	Common. Fresh or brackish water -
<input type="radio"/> <i>Lemna</i> sp.	Duckweed	Lemn-	-   -   -	- NP	--

Herbaceous <i>Lepidium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lepidium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lepidium didymum</i> (JM93: <i>Coronopus didymus</i> )	Lesser Swine Cress	Brassic-	-   -   -	- ZA	Common. Disturbed areas, fields, pastures (Mar–Jul)	
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i> )	Threadleaf Peppergrass	Brassic-	-   -   -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
<input type="radio"/> <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	-   -   -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)	
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i> )	Bicolor Leptosiphon	Polemoni-	-   -   -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Linum bienne</i>	Narrow-leaf Flax	Lin-	-   -   -	- ZP	Garden escape; grassland, woodland, disturbed places, esp coastal (Mar–Jun)	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	-   -   -	- NP	Open, grassy slopes (Mar–Apr)	
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	-   -   -	- NP	Shaded slopes (Feb–Jun)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i> )	Daggerleaf Cottonrose	Aster-	-   -   -	- ZA	Bare or grassy openings, burns (Mar–Jul)	
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Api-	-   -   B	- NP	Woodland, brushy slopes (Apr–Jun)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	-   -   -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Api-	-   -   -	- NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	-   -   -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus</i> sp.	Lupine	Fab-	-   -   -	- NAP - -		
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	-   -   -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	-   -   -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i> )	Fat False Solomon's Seal	Rusc-	-   -   -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i> )	Starry False Solomon's Seal	Rusc-	-   -   -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	-   -   -	- ZA	Disturbed places (Mar–Jun)	

Herbaceous <i>Malva</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Malva</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Malva parviflora</i>	Cheeseweed	Malv-	-   -   -	- ZA	Common. Disturbed places (Mar–May)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i> )	California Man-root	Cucurbit-	-   -   -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marah oregana</i> (JM93: <i>M. oreganus</i> )	Coast Man-root	Cucurbit-	-   -   C	- NP	Shrubby or open areas, forest edges (Mar–May)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	-   -   -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i> )	Pineapple Weed	Aster-	-   -   -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	-   -   -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Medicago sativa</i>	Alfalfa	Fab-	-   -   -	- ZP	Disturbed, agricultural areas (Apr–Oct)	
<input type="radio"/> <i>Melilotus albus</i> (JM93: <i>M. alba</i> )	White Sweetclover	Fab-	-   -   -	- ZAB	Locally abundant. Pastures, open disturbed sites (May–Sep)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i> )	Sourclover	Fab-	-   -   -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Melissa officinalis</i>	Lemon Balm	Lami-	-   -   -	- ZP	Moist sites, meadows, fields (Jun–Sep)	
<input type="radio"/> <i>Mentha arvensis</i>	Field Mint	Lami-	-   -   B	- NP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha piperita</i>	Peppermint	Lami-	-   -   -	- ZP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha pulegium</i>	Pennyroyal	Lami-	-   -   -	M ZP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha sp.</i>	Mint	Lami-	-   -   -	- -		
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i> )	California Saxifrage	Saxifrag-	-   -   -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	-   -   -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i> )	Slender Annual Phlox	Polemoni-	-   -   -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	-   -   -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i> )	Coyote-mint	Lami-	-   -   -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Myosotis latifolia</i>	Broadleaved Forget-me-not	Boragin-	-   -   -	L ZP	Moist, disturbed, shady places (Feb–Jul)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i> )	Water Cress	Brassic-	-   -   -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	-   -   -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	-   -   -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila parviflora</i> var. <i>parviflora</i>	Small-flower Nemophila	Boragin-	-   -   C	- NA	Woodland, forest, roadsides, slopes (Mar–Jul)	

Herbaceous <i>Osmorhiza</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Osmorhiza</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i> )	Sweet-cicely	Api-	-   -   -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	-   -   -	M ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)
<input type="radio"/> <i>Oxalis pilosa</i> (JM93: <i>O. albicans</i> subsp. <i>pilosa</i> )	Hairy Wood Sorrel	Oxalid-	-   -   A2	- NP	Coastal grassland, scrub, chaparral (Feb–Sep)
<input type="radio"/> <i>Oxalis</i> sp.	Wood Sorrel	Oxalid-	-   -   -	- -	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	-   -   -	- NP	Dry chaparral, oak/pine forest (Mar–May)
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	-   -   -	- NP	Open grassland, serpentine outcrops (Jul–Aug)
<input type="radio"/> <i>Persicaria amphibia</i> (JM93: <i>Polygonum amphibium</i> var. <i>emersum</i> , <i>stipulaceum</i> )	Water Smartweed	Polygon-	-   -   -	- NP	Shallow lakes, streams, shores (Jun–Nov)
<input type="radio"/> <i>Persicaria lapathifolia</i> (JM93: <i>Polygonum lapathifolium</i> )	Willow Weed	Polygon-	-   -   C	- NA	Moist places, disturbed areas, roadsides, fields (Jun–Oct)
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	-   -   -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	-   -   -	- NA	Common. Clay to rocky soils, slopes (Mar–May)
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	-   -   -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)
<input type="radio"/> <i>Phacelia nemoralis</i> subsp. <i>nemoralis</i>	Bristly Phacelia	Boragin-	-   -   B	- NBP	Moist slopes, streambanks, mixed-evergreen forest (Apr–Jul)
<input type="radio"/> <i>Pholistoma auritum</i> var. <i>auritum</i>	Fiesta Flower	Boragin-	-   -   B	- NA	Ocean bluffs, talus slopes, woodland, streambanks, canyons (Mar–Jun)
<input type="radio"/> <i>Piperia</i> sp.	Rein Orchid	Orchid-	-   -   -	- NP	--
<input type="radio"/> <i>Plagiobothrys canescens</i> var. <i>canescens</i> (JM93: no var.)	Valley Popcornflower	Boragin-	-   -   C	- NA	Common. Grassland, woodland, coastal scrub, desert scrub, roadsides (Mar–Jun)
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	-   -   -	- -	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	-   -   -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	-   -   -	L ZA	Common. Disturbed areas (Apr–Aug)
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Plantagin-	-   -   -	- ZAP	Disturbed areas (Apr–Sep)
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	-   -   -	- NA	Open grassland, sandy soil, burns (Mar–May)
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	-   -   -	- NA	Grassy, brushy areas (Mar–Jun)
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i> )	Knotweed	Polygon-	-   -   -	- ZA	Disturbed places (May–Nov)
<input type="radio"/> <i>Potamogeton crispus</i>	Crisp-leaved Pondweed	Potamogeton-	-   -   -	M ZP	Uncommon. Shallow water, ponds, streams. (Jul–Sep)

Herbaceous <i>Prosartes</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Prosartes</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i> )	Hooker Fairy Bells	Lili-	-   -   C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i> )	California Everlasting	Aster-	-   -   -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i> )	Weedy Cudweed	Aster-	-   -   -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i> )	Cotton-batting Plant	Aster-	-   -   -	- NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	-   -   -	- NA	Common. Sand or gravel (Mar–Jul)
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	-   -   B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	-   -   -	- NP	Grassland, open woodland (Mar–Aug)
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	-   -   -	- NA	Grassland, open woodland (Mar–May)
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	-   -   -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)
<input type="radio"/> <i>Ranunculus repens</i>	Creeping Buttercup	Ranuncul-	-   -   -	L ZP	Meadows, road banks, disturbed places, edges of marshes, streams (All year)
<input type="radio"/> <i>Ranunculus sceleratus</i> var. <i>sceleratus</i>	Cursed Crowsfoot	Ranuncul-	-   -   -	- NA	Ponds, riverbanks (Apr–Jun, Oct)
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	-   -   -	L ZA	Disturbed areas, fields (May–Jul)
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	-   -   -	M ZP	± Disturbed, often acidic places (Apr–Jul)
<input type="radio"/> <i>Rumex conglomeratus</i>	Green Dock	Polygon-	-   -   -	- ZP	Common. Moist places (May–Aug)
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	-   -   -	L ZP	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	-   -   -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)
<input type="radio"/> <i>Rumex sp.</i>	Dock	Polygon-	-   -   -	- -	-
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	-   -   -	- NP	Woodland (May–Sep)
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Caryophyll-	-   -   -	- NA	Sandy disturbed areas, river bars, streambanks (Spring–early summer)
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	-   -   -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	-   -   -	- NP	Open slopes, ravines, woodland (Mar–May)
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	-   -   -	- ZA	Grassy slopes, roadsides (Apr–Jun)
<input type="radio"/> <i>Scrophularia californica</i> (JM93: <i>subsp. californica, floribunda</i> )	California Figwort	Scrophulari-	-   -   -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	-   -   -	- ZA	Common. Disturbed areas (Feb–Jul)

Herbaceous <i>Sherardia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Sherardia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	-   -   -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. ?	Common Checkerbloom	Malv-	-   -   -	- NP	--
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>malviflora</i>	Common Checkerbloom	Malv-	-   -   B	- NP	Coastal prairie, scrub, open forest (Mar–Jul)
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	-   -   -	- ZA	Fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	-   -   -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)
<input type="radio"/> <i>Sinapis arvensis</i>	Charlock	Brassic-	-   -   -	L ZA	Disturbed areas (Mar–Sep)
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	-   -   -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	-   -   -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)
<input type="radio"/> <i>Solanum americanum</i>	Small-flowered Nightshade	Solan-	-   -   -	- NAP	Open, often disturbed places (Apr–Nov)
<input type="radio"/> <i>Solanum nigrum</i>	Black Nightshade	Solan-	-   -   -	- ZA	Disturbed places (Mar–Oct)
<input type="radio"/> <i>Solidago velutina</i> subsp. <i>californica</i> (JM93: <i>S. californica</i> )	California Goldenrod	Aster-	-   -   -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	-   -   -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	-   -   -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	-   -   -	- ZA	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Spergula arvensis</i> (JM93: subsp. <i>arvensis</i> )	Stickwort	Caryophyll-	-   -   -	- ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	-   -   -	- ZAP	Forest, meadows, mud flats, disturbed
<input type="radio"/> <i>Stachys ajugoides</i> (JM93: var. <i>ajugoides</i> )	Bugle Hedge-nettle	Lami-	-   -   A2	- NA	Moist, open places, gen remain wet into summer (Jun–Sep)
<input type="radio"/> <i>Stachys bullata</i>	California Hedge-nettle	Lami-	-   -   A1?	- NP	Dry slopes near coast (Mar–Sep)
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Lami-	-   -   C	- NP	Streambanks, springs, pine/oak forest (Jun–Oct)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i> )	Common Rigid Hedge-nettle	Lami-	-   -   -	- NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	-   -   -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)
<input type="radio"/> <i>Streptanthus glandulosus</i> subsp. <i>glandulosus</i> (JM93: incl. <i>S. albidus</i> subsp. <i>peramoenus</i> )	Bristly Jewel Flower	Brassic-	-   -   B	- NA	Serpentine, bare slopes, chaparral & woodland openings (Apr–Jul)
<input type="radio"/> <i>Symphyotrichum chilense</i> (JM93: <i>Aster chilensis</i> )	Common California Wild Aster	Aster-	-   -   -	- NP	Grassland, salt marshes, disturbed places (Jun–Oct)
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	-   -   -	- ZBP	Abundant. Esp disturbed areas (All year)
<input type="radio"/> <i>Taraxia ovata</i> (JM93: <i>Camissonia</i> )	Golden Eggs Suncup	Onagr-	-   -   C	- NP	Grassy fields, gen clay soil (Mar–Jun)

Herbaceous <i>Tauschia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Tauschia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	-   -   C	- NP	Chaparral, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	-   -   -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	-   -   -	M ZA	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	-   -   -	- ZA	Disturbed places (Apr–Jun)	
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i> )	Common Star Lily	Melanthi-	-   -   -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)	
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	-   -   -	- ZBP	Common. Disturbed places (Mar–Nov)	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Fab-	-   -   -	- NA	Open, grassy areas, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium campestre</i>	Hop Clover	Fab-	-   -   -	- ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	-   -   -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	-   -   -	- ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium fragiferum</i>	Strawberry Clover	Fab-	-   -   -	- ZP	Roadsides, gen in saline soil (Late spring)	
<input type="radio"/> <i>Trifolium glomeratum</i>	Clustered Clover	Fab-	-   -   -	- ZA	Uncommon. Disturbed areas (Mar–May)	
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i> )	Pinpoint Clover	Fab-	-   -   -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	-   -   -	M ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium incarnatum</i>	Crimson Clover	Fab-	-   -   -	- ZA	Uncommon. Disturbed areas (May–Aug)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	-   -   -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	-   -   -	- NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium repens</i>	White Clover	Fab-	-   -   -	- ZP	Disturbed areas (Feb–Dec)	
<input type="radio"/> <i>Trifolium sp.</i>	Clover	Fab-	-   -   -	- -	- -	
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Fab-	-   -   -	- ZA	Meadows, roadsides, disturbed areas (Mar–Apr)	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	-   -   -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	-   -   -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)	
<input type="radio"/> <i>Trillium ovatum</i> subsp. <i>ovatum</i>	Western White Trillium	Melanthi-	-   -   A2	- NP	Redwood, mixed-evergreen forest on moist wooded slopes (Feb–Apr)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	-   -   -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	-   -   -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Turritis glabra</i> (JM93: <i>Arabis</i> , var. <i>glabra</i> )	Tower Mustard	Brassic-	-   -   C	- NP	Open fields, meadows, slopes (Apr–Jul)	



Herbaceous <i>Uropappus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Uropappus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	-   -   -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)
<input type="radio"/> <i>Urtica dioica subsp. holosericea</i>	Hoary Nettle	Urtic-	-   -   -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	-   -   -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	-   -   -	- NP	Open, dry to wet places (May–Sep)
<input type="radio"/> <i>Veronica americana</i>	American Brooklime	Plantagin-	-   -   C	- NP	Common. Moist to wet soil, springs, slow streams, meadows, lakeshores (May–Aug)
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	-   -   -	- ZA	Wet, disturbed areas, fields (Feb–May)
<input type="radio"/> <i>Veronica sp.</i>	Speedwell	Plantagin-	-   -   -	- -	
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	-   -   -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Vicia gigantea</i>	Giant Vetch	Fab-	-   -   C	- NP	Coastal shrub, coastal forest, chaparral (Mar–Aug)
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)
<input type="radio"/> <i>Vicia sp.</i>	Vetch	Fab-	-   -   -	- -	
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Apocyn-	-   -   -	M ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	-   -   C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	-   -   -	- NP	Grassland (Apr–Aug)
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	-   -   C	- NP	Gen shady sites (Mar–Jun)
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	-   -   -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	-   -   -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)
<input type="radio"/> <i>Zeltnera muehlenbergii (JM93: Centaurium)</i>	Monterey Centaury	Gentian-	-   -   -	- NA	Moist coastal bluffs, forest openings (Jun–Aug)

<b>Woody Acacia</b>	<b>Grouped by Growth Form Sorted Alphabetically by Scientific Name</b>					<b>Woody Acacia</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Acacia melanoxylo</i>	Blackwood Acacia	Fab-	-   -   -	L	ZP	Uncommon. Disturbed areas (Feb-Mar)
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	-   -   -	-	NP	Common. Streambanks, canyons (Mar-Jun)
<input type="radio"/> <i>Acer negundo</i> (JM93: var. <i>californicum</i> )	Box Elder	Sapind-	-   -   -	-	NP	Streamsides, bottomland. (Mar-Apr)
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	-   -   -	-	NP	Dry slopes, ridges, chaparral (May-Jun)
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	-   -   -	-	NP	Dry slopes, canyons, borders of streams (May-Jun)
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	-   -   -	-	NP	Along permanent streams (Apr-Jun)
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	-   -   C	-	NP	Conifer, oak forests (Mar-May)
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	-   -   -	-	NP	Coastal scrub, chaparral, open woodland (Aug-Nov)
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	-   -   -	-	NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul-Dec)
<input type="radio"/> <i>Baccharis salicifolia</i> subsp. <i>salicifolia</i> (JM93: no subsp.)	Mule Fat	Aster-	-   -   -	-	NP	Riparian woodland, canyon bottoms, disturbed sites, often forming thickets (All year)
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	-   -   C	-	NP	Dry, rocky slopes, chaparral (Mar-May)
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	-   -   C	-	NP	Along streams, wet places (Jun-Sep)
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Corn-	-   -   C	-	NP	Generally moist places (May-Jul)
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Corn-	-   -   C	-	NP	Generally moist habitats (May-Jul)
<input type="radio"/> <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i> )	California Hazelnut	Betul-	-   -   -	-	NP	Common. Many habitats, esp moist, shady places (Jan-Mar)
<input type="radio"/> <i>Cotoneaster pannosus</i> (JM93: <i>C. pannosa</i> )	Silverleaf Cotoneaster	Ros-	-   -   -	M	ZP	Disturbed places, mixed-evergreen forest (May-Jul)
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>foliolosum</i>	Leafy California Buckwheat	Polygon-	-   -   A2	-	NP	Gravel (All year)
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	-   -   -	-	NP	Many dry habitats (Apr-Aug)
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	-   -   -	M	ZP	Common. Disturbed areas (Oct-Jan)
<input type="radio"/> <i>Eucalyptus</i> sp.	Gum	Myrt-	-   -   -	-	ZP	--
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i> )	California Coffee Berry	Rhamn-	-   -   -	-	NP	Coastal-sage scrub, chaparral, forest, woodland (May-Jul)
<input type="radio"/> <i>Fraxinus</i> sp.	Ash	Ole-	-   -   -	-	NP	--
<input type="radio"/> <i>Garrya elliptica</i>	Coast Silk Tassel	Garry-	-   -   B	-	NP	Seacliffs, sand dunes, chaparral, foothill-pine woodland (Jan-Mar)
<input type="radio"/> <i>Genista monspessulana</i>	French Broom	Fab-	-   -   -	H	ZP	Common. Disturbed places. (Mar-Jun)
<input type="radio"/> <i>Hedera helix</i> (JM93: subsp. <i>helix</i> )	English Ivy	Arali-	-   -   -	H	ZP	Woodland, open, disturbed areas (Aug-Nov)

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b><i>Heteromeles</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Heteromeles</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	-   -   -	-	NP	Chaparral, oak woodland, mixed-evergreen forest (May)Jun–Aug)
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	-   -   -	-	NP	Moist woodland edges, rocky slopes (May–Aug)
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i> )	Hairy Vine Honeysuckle	Caprifoli-	-   -   -	-	NP	Canyons, streamsides, woodland (May–Jun)
<input type="radio"/> <i>Lonicera subspicata</i> var. <i>denudata</i>	Southern Honeysuckle	Caprifoli-	-   -   B	-	NP	Chaparral slopes (Jun–Jul)
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	-   -   -	-	NP	--
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>albifrons</i>	Bush Lupine	Fab-	-   -   -	-	NP	Common. Chaparral, foothill woodland (Mar–Jun)
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>collinus</i>	Bay Area Silver Lupine	Fab-	-   -   -	-	NP	Cliffs, forest openings (Mar–Jun)
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	-   -   -	-	NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	-   -   -	-	NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)
<input type="radio"/> <i>Olea europaea</i>	Olive	Ole-	-   -   -	L	ZP	Generally waif, persisting from cultivation (Feb–Jun)
<input type="radio"/> <i>Physocarpus capitatus</i>	Pacific Ninebark	Ros-	-   -   C	-	NP	Moist banks, n-facing slopes, mixed-conifer forest (May–Jul)
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	-   -   -	-	NP	Closed-cone-pine forest, oak woodland -
<input type="radio"/> <i>Pinus sabiniana</i>	Gray Pine	Pin-	-   -   -	-	NP	Foothill woodland, n oak woodland, chaparral, infertile soils in mixed-conifer and hardwood forests -
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	-   -   B	-	NP	Common. Streamsides, canyons, arroyos (Feb–Apr)
<input type="radio"/> <i>Populus fremontii</i> subsp. <i>fremontii</i>	Fremont Cottonwood	Salic-	-   -   -	-	NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)
<input type="radio"/> <i>Prunus cerasifera</i>	Cherry Plum	Ros-	-   -   -	L	ZP	Roadsides, streambanks, chaparral as waif (Feb–Mar)
<input type="radio"/> <i>Prunus dulcis</i>	Almond	Ros-	-   -   -	-	ZP	Canyons, roadsides, grassland (as waif) (Feb–Mar)
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Ros-	-   -   C	-	NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Ros-	-   -   -	-	NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)
<input type="radio"/> <i>Ptelea crenulata</i>	Western Hop Tree	Rut-	-   -   B	-	NP	Scrub, woodland (Apr–May)
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	-   -   A2	-	NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	-   -   C	-	NP	Dry slopes, chaparral (Feb–Apr)

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Quercus</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Quercus</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	-   -   -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus durata</i> var. <i>durata</i>	Leather Oak	Fag-	-   -   A2	- NP	Chaparral, generally serpentine (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	-   -   B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus palmeri</i>	Palmer Oak	Fag-	-   -   A2	- NP	Uncommon. Rocky slopes, flats (Apr–May)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Rhamn-	-   -   C	- NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	-   -   -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes divaricatum</i> var. <i>pubiflorum</i>	Straggle Gooseberry	Grossulari-	-   -   B	- NP	Uncommon. Coastal bluffs, forest edges (Mar–May)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	-   -   -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes sanguineum</i> var. <i>glutinosum</i>	Red-flowering Currant	Grossulari-	-   -   C	- NP	Many habitats (Feb–Apr)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Fab-	-   -   -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	-   -   -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	-   -   -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa</i> sp.	Rose	Ros-	-   -   -	- NP	--	
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i> )	Himalayan Blackberry	Ros-	-   -   -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	-   -   -	- NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	-   -   -	- NP	Open, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Salic-	-   -   -	- NP	Common. Floodplains, sandy gravel (Apr–May)	
<input type="radio"/> <i>Salix laevigata</i>	Red Willow	Salic-	-   -   -	- NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)	
<input type="radio"/> <i>Salix lasiandra</i> var. <i>lasiandra</i> (JM93: <i>S. lucida</i> subsp. <i>Lasiandra</i> )	Pacific Willow	Salic-	-   -   -	- NP	Wet meadows, lakeshores, riverbanks (May–Jun)	
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	-   -   -	- NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="radio"/> <i>Salix</i> sp.	Willow	Salic-	-   -   -	- ?P	--	
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i> )	Blue Elderberry	Adox-	-   -   -	- NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="radio"/> <i>Sequoia sempervirens</i>	Redwood	Cupress-	-   -   B	- NP	Redwood forest -	
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	-   -   -	- NP	Shrubland, mixed-evergreen forest, woodland (All year)	

<b>Woody</b>	<b>Grouped by Growth Form</b>						<b>Woody</b>
<b><i>Symphoricarpos</i></b>	<b>Sorted Alphabetically by Scientific Name</b>						<b><i>Symphoricarpos</i></b>
<input checked="" type="checkbox"/> <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>	
<input type="checkbox"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	-   -   -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="checkbox"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	-   -   -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="checkbox"/> <i>Tamarix parviflora</i>	Smallflower Tamarisk	Tamaric-	-   -   -	H	ZP	Common. Washes, streambanks, slopes, roadsides (Mar–Apr)	
<input type="checkbox"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	-   -   -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="checkbox"/> <i>Umbellularia californica</i>	California Bay	Laur-	-   -   -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)	

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Acer negundo* var. *californicum* (no var.)

*Adenostoma fasciculatum* (var. *fasciculatum*)

*Agoseris grandiflora* (var. *grandiflora*)

*Agoseris heterophylla* (var. *cryptopleura*)

*Agrostis viridis* (*Polypogon*)

*Alisma plantago-aquatica* (*A. triviale*)

*Amsinckia menziesii* var. *intermedia* (*A. intermedia*)

*Amsinckia menziesii* var. *menziesii* (no var.)

*Antirrhinum vexillo-calyculatum* ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

*Arabis breweri* var. *breweri* (*Boechera breweri* subsp. *breweri*)

*Arabis glabra* var. *glabra* (*Turritis glabra*)

*Arctostaphylos tomentosa* ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

*Aster chilensis* (*Symphyotrichum chilense*)

*Aster lentus* (*Symphyotrichum lentum*)

*Aster radulinus* (*Eurybia radulina*)

*Aster subulatus* var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

*Atriplex joaquiniana* (*A. joaquinana*)

*Atriplex lentiformis* ssp. *lentiformis* (no subsp.)

*Atriplex triangularis* (*A. prostrata*)

*Azolla mexicana* (*A. microphylla*)

*Baccharis douglasii* (*B. glutinosa*)

*Baccharis pilularis* (subsp. *consanguinea*)

*Baccharis salicifolia* (subsp. *salicifolia*)

*Balsamorhiza macrolepis* var. *macrolepis* (no var.)

*Blepharizonia plumosa* ssp. *plumosa* (no subsp.)

*Blepharizonia plumosa* ssp. *viscida* (*Blepharizonia laxa*)

*Boschniakia strobilacea* (*Kopsiopsis*)

*Bromus catharticus* (var. *catharticus*)

*Camissonia boothii* ssp. *decorticans* (*Eremothera*)

*Camissonia graciliflora* (*Tetrapteron graciliflorum*)

*Camissonia intermedia* (*Camissoniopsis*)

*Camissonia micrantha* (*Camissoniopsis*)

*Camissonia ovata* (*Taraxia*)

*Cardaria draba* (*Lepidium*)

*Carduus pycnocephalus* (subsp. *pycnocephalus*)

*Carex deweyana* ssp. *leptopoda* (*C. leptopoda*)

*Carex lanuginosa* (*C. pellita*)

*Carex ovalis* (*C. leporina*)

*Ceanothus thyrsoiflorus* (var. *thyrsoiflorus*)

*Centaurium davyi* (*Zeltnera*)

*Centaurium muehlenbergii* (*Zeltnera*)

*Cephalanthus occidentalis* var. *californicus* (no var.)

*Chamomilla suaveolens* (*Matricaria discoidea*)

*Chenopodium ambrosioides* (*Dysphania*)

*Chenopodium macrospermum* var. *halophilum* (no var.)

*Chenopodium multifidum* (*Dysphania multifida*)

*Chenopodium rubrum* (var. *rubrum*)

*Chrysothamnus nauseosus* ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

*Cirsium cymosum* (var. *cymosum*)

*Claytonia parviflora* ssp. *parviflora* (subsp. *utahensis*)

*Collinsia bartsifolia* var. *davidsonii* (var. *stricta*)

*Collinsia heterophylla* (var. *heterophylla*)

*Conyza bilbaoana* (*Erigeron sumatrensis*)

*Conyza bonariensis* (*Erigeron*)

*Conyza canadensis* (*Erigeron*)

*Conyza coulteri* (*Laennecia*)

*Corallorhiza maculata* (var. *maculata*)

*Cordylanthus mollis* ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

*Coronopus didymus* (*Lepidium didymum*)

*Corylus cornuta* var. *californica* (subsp. *californica*)

*Cotoneaster pannosa* (*C. pannosus*)

*Cryptantha decipiens* (*C. rattanii*)

*Cryptantha intermedia* (var. *intermedia*)

*Cryptantha muricata* (var. *muricata*)

*Cryptantha torreyana* (var. *pumila*)

*Cupressus macrocarpa* (*Hesperocyparis*)

*Cuscuta salina* var. *major* (*C. pacifica* var. *pacifica*)

*Cynara cardunculus* (subsp. *flavescens*)

*Danthonia californica* var. *californica* (no var.)

*Dicentra chrysantha* (*Ehrendorferia*)

*Disporum hookeri* (*Prosartes*)

*Disporum smithii* (*Prosartes*)

*Draba cuneifolia* var. *integrifolia* (no var.)

*Duchesnea indica* (no var.)

*Epilobium pygmaeum* (*E. campestre*)

*Eragrostis curvula* var. *curvula* (no var.)

*Erechtites glomerata* (*Senecio glomeratus*)

*Erechtites minima* (*Senecio minimus*)

*Eremocarpus setigerus* (*Croton*)

*Erigeron philadelphicus* (var. *philadelphicus*)

*Eriogonum umbellatum* var. *bahiiforme* (var. *smallianum*)

*Eriophyllum lanatum* var. *achillaeoides* (var. *achilleoides*)

*Eryngium aristulatum* var. *aristulatum* (*E. jepsonii*)

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

*Erysimum capitatum* var. *angustatum* (var. *capitatum*)

*Filago californica* (*Logfia filaginoides*)

*Filago gallica* (*Logfia*)

*Fritillaria affinis* var. *affinis* (no var.)

*Galium trifidum* var. *pacificum* (subsp. *columbianum*)

*Gastridium ventricosum* (*G. phleoides*)

*Gnaphalium bicolor* (*Pseudognaphalium biolettii*)

*Gnaphalium californicum* (*Pseudognaphalium*)

*Gnaphalium canescens* ssp. *beneolens*  
(*Pseudognaphalium beneolens*)

*Gnaphalium canescens* ssp. *microcephalum*  
(*Pseudognaphalium microcephalum*)

*Gnaphalium luteo-album* (*Pseudognaphalium luteoalbum*)

*Gnaphalium purpureum* (*Gamochaeta ustulata*)

*Gnaphalium ramosissimum* (*Pseudognaphalium*)

*Gnaphalium stramineum* (*Pseudognaphalium*)

*Grindelia camporum* var. *camporum* (no var.)

*Grindelia hirsutula* var. *davyi* (*G. camporum*)

*Grindelia hirsutula* var. *hirsutula* (no var.)

*Guillenia flavescens* (*Caulanthus*)

*Guillenia lasiophylla* (*Caulanthus lasiophyllus*)

*Hedera helix* ssp. *helix* (no subsp.)

*Heliotropium curassavicum* (var. *oculatum*)

*Hemizonia congesta* ssp. *congesta* (subsp. *lutescens*)

*Hemizonia corymbosa* ssp. *corymbosa* (*Deinandra corymbosa*)

*Hemizonia fitchii* (*Centromadia*)

*Hemizonia lobbii* (*Deinandra*)

*Hemizonia parryi* ssp. *congdonii* (*Centromadia*)

*Hemizonia parryi* ssp. *parryi* (*Centromadia*)

*Hemizonia pungens* ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

*Hemizonia pungens* ssp. *pungens* (*Centromadia*)

*Heracleum lanatum* (*H. maximum*)

*Herniaria hirsuta* ssp. *cinerea* (var. *cinerea*)

*Holodiscus discolor* (var. *discolor*)

*Hordeum jubatum* (subsp. *jubatum*)

*Horkelia californica* ssp. *californica* (var. *californica*)

*Horkelia californica* ssp. *dissita* (var. *elata*)

*Horkelia californica* ssp. *frondosa* (var. *frondosa*)

*Isopyrum occidentale* (*Enemion*)

*Iva axillaris* ssp. *robustior* (no subsp.)

*Juglans californica* var. *hindsii* (*J. hindsii*)

*Juncus balticus* (subsp. *ater*)

*Juncus effusus* var. *pacificus* (subsp. *pacificus*)

*Juncus lesueurii* (*J. lescurii*)

*Koeleria phleoides* (*K. gerardii*)

*Lagophylla ramosissima* ssp. *ramosissima* (no subsp.)

*Lasthenia californica* (subsp. *californica*)

*Lavatera cretica* (*Malva pseudolavatera*)

*Lepidium dictyotum* var. *dictyotum* (no var.)

*Lepidium latipes* var. *latipes* (no var.)

*Lepidium nitidum* var. *nitidum* (no var.)

*Lepidium nitidum* var. *oreganum* (no var.)

*Leptochloa fascicularis* (*L. fusca* subsp. *fascicularis*)

*Lessingia filaginifolia* var. *californica* (*Corethrogyne filaginifolia*)

*Lessingia filaginifolia* var. *filaginifolia*  
(*Corethrogyne filaginifolia*)

*Lewisia rediviva* (var. *rediviva*)

*Leymus condensatus* (*Elymus*)

*Leymus triticoides* (*Elymus*)

*Lilaea scilloides* (*Triglochin*)

*Limosella subulata* (*Limosella australis*)

*Linanthus acicularis* (*Leptosiphon*)

*Linanthus ambiguus* (*Leptosiphon*)

*Linanthus androsaceus* (*Leptosiphon*)

*Linanthus bicolor* (*Leptosiphon*)

*Linanthus ciliatus* (*Leptosiphon*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus parviflorus* (*Leptosiphon*)

*Linaria canadensis* (*Nuttallanthus texanus*)

*Lolium multiflorum* (*Festuca perennis*)

*Lolium perenne* (*Festuca perennis*)

*Lolium temulentum* (*Festuca temulenta*)

*Lonicera hispidula* var. *vacillans* (no var.)

*Lotus humistratus* (*Acmispon brachycarpus*)

*Lotus micranthus* (*Acmispon parviflorus*)

*Lotus purshianus* var. *purshianus* (*Acmispon americanus* var. *americanus*)

*Lotus scoparius* var. *scoparius* (*Acmispon glaber* var. *glaber*)

*Lotus strigosus* (*Acmispon*)

*Lotus wrangelianus* (*Acmispon*)

*Luzula comosa* (var. *comosa*)

*Madia madioides* (*Anisocarpus*)

*Malacothamnus fasciculatus* (var. *nuttallii*)

*Marah fabaceus* (*M. fabacea*)

*Marah oreganus* (*M. oregana*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Melilotus alba* (*M. albus*)

*Melilotus indica* (*M. indicus*)

*Mentha spicata* var. *spicata* (no var.)

*Mimulus aurantiacus* (var. *aurantiacus*)

*Mirabilis californica* (*M. laevis* var. *crassifolia*)

*Monardella antonina* ssp. *antonina* (*M. villosa* subsp. *villosa*)

*Monardella douglasii* ssp. *douglasii* (no subsp.)

*Monardella villosa* ssp. *globosa* (subsp. *villosa*)

*Montia fontana* ssp. *chondrosperma* (no subsp.)

*Myrica californica* (*Morella*)

*Nassella cernua* (*Stipa*)

*Nassella lepida* (*Stipa*)

*Nassella pulchra* (*Stipa*)

*Osmorhiza chilensis* (*O. berteroi*)

*Oxalis albicans* ssp. *pilosa* (*O. pilosa*)

*Parvisedum pentandrum* (*Sedella pentandra*)

*Phacelia ramosissima* var. *latifolia* (no var.)

*Phacelia ramosissima* var. *ramosissima* (no var.)

*Phlox gracilis* (*Microsteris*)

*Phoradendron macrophyllum* (*P. serotinum* subsp. *macrophyllum*)

*Phoradendron villosum* (*P. serotinum* subsp. *tomentosum*)

*Phyla nodiflora* var. *incisa* (no var.)

*Phyla nodiflora* var. *nodiflora* (no var.)

*Picris echioides* (*Helminthotheca*)

*Piperia elegans* (subsp. *elegans*)

*Piptatherum miliaceum* (*Stipa miliacea* var. *miliacea*)

*Plagiobothrys canescens* (var. *canescens*)

*Plagiobothrys fulvus* (var. *campestris*)

*Plectritis brachystemon* (*P. congesta* subsp. *brachystemon*)

*Plectritis ciliosa* ssp. *ciliosa* (no subsp.)

*Plectritis ciliosa* ssp. *insignis* (no subsp.)

*Plectritis congesta* (subsp. *congesta*)

*Pleuropogon californicus* (var. *californicus*)

*Pluchea odorata* (var. *odorata*)

*Poa bulbosa* (subsp. *vivipara*)

*Polygonum amphibium* var. *emersum* (*Persicaria amphibia*)

*Polygonum amphibium* var. *stipulaceum* (*Persicaria amphibia*)

*Polygonum arenastrum* (*P. aviculare* subsp. *depressum*)

*Polygonum hydropiperoides* (*Persicaria*)

*Polygonum lapathifolium* (*Persicaria lapathifolia*)

*Polygonum persicaria* (*Persicaria maculosa*)

*Polygonum punctatum* (*Persicaria punctata*)

*Populus balsamifera* ssp. *trichocarpa* (*P. trichocarpa*)

*Potamogeton pectinatus* (*Stuckenia pectinata*)

*Potentilla glandulosa* ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

*Psilocarphus tenellus* var. *tenellus* (no var.)

*Ranunculus aquatilis* var. *capillaceus* (var. *diffusus*)

*Ranunculus californicus* (var. *californicus*)

*Ranunculus canus* (var. *canus*)

*Ranunculus occidentalis* (var. *occidentalis*)

*Ranunculus sceleratus* (var. *sceleratus*)

*Rhamnus californica* ssp. *californica* (*Frangula*)

*Rhamnus tomentella* ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

*Rhus trilobata* (*R. aromatica*)

*Ribes menziesii* (var. *menziesii*)

*Rorippa nasturtium-aquaticum* (*Nasturtium officinale*)

*Rorippa palustris* var. *occidentalis* (subsp. *palustris*)

*Rosa gymnocarpa* (var. *gymnocarpa*)

*Rubus discolor* (*R. armeniacus*)

*Rumex maritimus* (*R. fueginus*)

*Rumex salicifolius* var. *crassus* (*R. crassus*)

*Rumex salicifolius* var. *salicifolius* (no var.)

*Rumex salicifolius* var. *transitorius* (*R. transitorius*)

*Salicornia europaea* (*S. depressa*)

*Salicornia subterminalis* (*Arthrocnemum subterminale*)

*Salicornia virginica* (*S. pacifica*)

*Salix exigua* (var. *hindsiana*)

*Salix lucida* ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

*Sambucus mexicana* (*S. nigra* subsp. *caerulea*)

*Satureja douglasii* (*Clinopodium*)

*Saxifraga californica* (*Micranthes*)

*Scirpus acutus* var. *occidentalis* (*Schoenoplectus*)

*Scirpus americanus* (*Schoenoplectus*)

*Scirpus californicus* (*Schoenoplectus*)

*Scirpus cernuus* (*Isolepis*)

*Scirpus maritimus* (*Bolboschoenus maritimus* subsp. *paludosus*)

*Scirpus robustus* (*Bolboschoenus*)

*Scrophularia californica* ssp. *californica* (no subsp.)

*Scrophularia californica* ssp. *floribunda* (no subsp.)

*Senecio breweri* (*Packera*)

*Senecio mikanioides* (*Delairea odorata*)

*Silene californica* (*S. laciniata* subsp. *californica*)



---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Smilacina racemosa* (*Maianthemum racemosum*)

*Smilacina stellata* (*Maianthemum stellatum*)

*Solidago californica* (*S. velutina* subsp. *californica*)

*Solidago canadensis* ssp. *elongata* (*S. elongata*)

*Sparganium erectum* ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

*Sparganium eurycarpum* ssp. *eurycarpum* (var. *eurycarpum*)

*Spergula arvensis* ssp. *arvensis* (no subsp.)

*Spergularia media* ( var. *media*)

*Stachys ajugoides* var. *ajugoides* (no var.)

*Stachys ajugoides* var. *rigida* (*Stachys rigida* var. *quercetorum*)

*Streptanthus albidus* ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

*Stylomecon heterophylla* (*Papaver heterophyllum*)

*Taeniatherum caput-medusae* (*Elymus*)

*Thysanocarpus laciniatus* (var. *laciniatus*)

*Tolmiea menziesii* (*T. diplomenziesii*)

*Trifolium albopurpureum* var. *albopurpureum* (no var.)

*Trifolium albopurpureum* var. *dichotomum* (*T. dichotomum*)

*Trifolium albopurpureum* var. *olivaceum* (*T. olivaceum*)

*Trifolium barbigerum* var. *andrewsii* (*T. grayi*)

*Trifolium barbigerum* var. *barbigerum* (no var.)

*Trifolium gracilentum* var. *gracilentum* (no var.)

*Vicia americana* var. *americana* (subsp. *americana*)

*Vicia ludoviciana* var. *ludoviciana* (subsp. *ludoviciana*)

*Viola adunca* (var. *adunca*)

*Vulpia bromoides* (*Festuca*)

*Vulpia microstachys* var. *ciliata* (*Festuca microstachys*)

*Vulpia microstachys* var. *confusa* (*Festuca microstachys*)

*Vulpia microstachys* var. *microstachys* (*Festuca microstachys*)

*Vulpia microstachys* var. *pauciflora* (*Festuca microstachys*)

*Vulpia myuros* var. *hirsuta* (*Festuca myuros*)

*Vulpia myuros* var. *myuros* (*Festuca myuros*)

*Vulpia octoflora* var. *hirtella* (*Festuca octoflora*)

*Vulpia octoflora* var. *octoflora* (*Festuca octoflora*)

*Zigadenus fremontii* (*Toxicoscordion*)