

# Las Trampas Regional Wilderness

## Checklist of Wild Plants

### Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Las Trampas Regional Wilderness. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
<b>Scientific Name</b>	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> ( <a href="http://ucjeps.berkeley.edu/IJM.html">ucjeps.berkeley.edu/IJM.html</a> )
<b>(JM93 if different)</b>	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
<b>Common Name</b>	According to <i>JM2</i> and other references (not standardized)
<b>Family</b>	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
<b>Status</b>	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): <b>Federal/California (Fed./Calif.)   California Native Plant Society (CNPS)   East Bay chapter of the CNPS (EBCNPS)</b>  <b>Fed./Calif.:</b> FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare  <b>CNPS</b> (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California  <b>EBCNPS</b> (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9
<b>Inv</b>	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
<b>OL</b>	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
<b>Habitat (Bloom)</b>	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, [wlegard@ebparks.org](mailto:wlegard@ebparks.org)). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

# Las Trampas Regional Wilderness Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name					<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	-   -   -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Cystopteris fragilis</i>	Brittle Fern	Woodsi-	-   -   C	- NP	Shady, moist rock crevices, meadows, streamsides -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	-   -   -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum arvense</i>	Common Horsetail	Equiset-	-   -   -	- NP	Streambanks, wet meadows, springs, other wet, shaded places -	
<input type="radio"/> <i>Equisetum telmateia subsp. braunii</i>	Giant Horsetail	Equiset-	-   -   -	- NP	Streambanks, roadside ditches, seepage areas -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	-   -   -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis subsp. triangularis</i>	Goldenback Fern	Pterid-	-   -   -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	-   -   -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Dryopterid-	-   -   -	- NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -	
<input type="radio"/> <i>Pteridium aquilinum var. pubescens</i>	Bracken Fern	Dennstaedti-	-   -   -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	-   -   -	- NP	Near streams, springs, seeps -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	-   -   -	- NP	Common. Open meadows, woodland, forest, subalpine (Jun–Aug)
<input type="radio"/> <i>Aira caryophyllea</i>	Silver Hair Grass	Po-	-   -   -	- ZA	Sandy soils, open or disturbed sites (Apr–Jun)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	-   -   -	M ZA	Disturbed sites (Mar–Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	-   -   -	M ZA	Disturbed sites (Apr–Jun)
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	-   -   -	- NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr–Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	-   -   -	M ZA	Open, gen disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	-   -   -	L ZA	Fields, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	-   -   -	- NP	Shrubland, conifer forest, shaded streambanks, roadsides (May–Jul)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>madritensis</i>	Foxtail Chess	Po-	-   -   -	- ZA	Disturbed areas, roadsides (Apr–Jan)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	-   -   -	H ZA	Disturbed areas, roadsides (Mar–Jun)
<input type="radio"/> <i>Carex</i> sp.	Sedge	Cyper-	-   -   -	- -	
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	-   -   -	M ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Po-	-   -   -	- NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	-   -   C	- NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus trachycaulus</i> subsp. <i>trachycaulus</i>	Slender Wheat Grass	Po-	-   -   B	- NP	Dry to moist, open areas, forest, woodland (Jun–Aug)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i> )	Beardless Wild Rye	Po-	-   -   -	- NP	Dry to moist, often saline, meadows ( Jun–Jul)
<input type="radio"/> <i>Eragrostis curvula</i> (JM93: var. <i>curvula</i> )	Weeping Love Grass	Po-	-   -   -	- NP	Roadsides (Aug–Oct)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	-   -   -	M ZP	Disturbed places (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	-   -   C	- NP	Dry, open forest, moist streambanks, chaparral (May–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i> )	Rattail Sixweeks Grass	Po-	-   -   -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca octoflora</i> (JM93: <i>Vulpia</i> , var. <i>hirtella</i> , <i>octoflora</i> )	Slender Six-weeks Fescue	Po-	-   -   A2	- NA	Sandy to rocky soils, open sites (Mar–Jun)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i> )	Rye Grass	Po-	-   -   -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Po-	-   -   B	- NP	Sand dunes, grassland, subalpine forest (May–Jul)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	-   -   -	- NA	- -
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	-   -   -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Junc-	-   -   -	- NA	Damp sunny ground, gen disturbed (May–Sep)

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like
<i>Juncus</i>	Sorted Alphabetically by Scientific Name				<i>Juncus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Junc-	-   -   -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	-   -   -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	-   -   -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	-   -   C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	-   -   -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	-   -   C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	-   -   -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	-   -   -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Panicum</i> sp.	Panic Grass	Po-	-   -   -	- -	-
<input type="radio"/> <i>Phalaris aquatica</i>	Harding Grass	Po-	-   -   -	M ZP	Disturbed areas, roadsides (Apr–Aug)
<input type="radio"/> <i>Phalaris arundinacea</i>	Reed Canary Grass	Po-	-   -   A2	- NP	Moist habitats, woodland (May–Sep)
<input type="radio"/> <i>Phalaris minor</i>	Little-seeded Canary Grass	Po-	-   -   -	- ZA	Disturbed areas, cultivated fields (Apr–Jul)
<input type="radio"/> <i>Phalaris paradoxa</i>	Hood Canary Grass	Po-	-   -   -	- ZA	Disturbed areas, cultivated field (May–Aug)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	-   -   -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa howellii</i>	Howell Blue Grass	Po-	-   -   B	- NA	Rocky banks, shaded slopes of woodland, chaparral, disturbed places (Apr–Jun)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	-   -   -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	-   -   -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Polypogon viridis</i> (JM93: <i>Agrostis</i> )	Water Beard Grass	Po-	-   -   -	- ZP	Common. Disturbed areas, wet areas, ponds, streambanks (May–Jun)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i> )	Foothill Needle Grass	Po-	-   -   C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i> )	Purple Needle Grass	Po-	-   -   C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)

Herbaceous <i>Acaena</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Acaena</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Acaena pinnatifida</i> var. <i>californica</i>	California Acaena	Ros-	-   -   C	- NP	Coastal grassland, open, rocky slopes (Mar–May)
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	-   -   -	- NP	Many habitats (Apr–Sep)
<input type="radio"/> <i>Achyrachaena mollis</i>	Blow Wives	Aster-	-   -   -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i> )	Colchita	Fab-	-   -   -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i> )	Deerweed	Fab-	-   -   -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)
<input type="radio"/> <i>Acmispon</i> or <i>Hosackia</i> sp. (JM93: <i>Lotus</i> sp.)	Lotus	Fab-	-   -   -	- -	- -
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i> )	Small-flower Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i> )	California Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	-   -   -	- NP	Grassland, scrub, woodland (Apr–Jul)
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	-   -   -	- NP	Common. Grassy slopes (Apr–May)
<input type="radio"/> <i>Allophyllum gilioides</i> subsp. <i>gilioides</i>	Purple False Gilia	Polemoni-	-   -   B	- NA	Open, sandy, gen damp or grassy areas (Apr–Jun)
<input type="radio"/> <i>Amaranthus albus</i>	Tumbleweed	Amaranth-	-   -   -	- ZA	Disturbed areas, roadsides, riverbanks, sandy places, agricultural fields (Jun–Oct)
<input type="radio"/> <i>Amaranthus retroflexus</i>	Redroot Pigweed	Amaranth-	-   -   -	- ZA	Common. Wet fields, roadside ditches, waste places, agricultural fields (Jun–Nov)
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i> )	Common Fiddleneck	Boragin-	-   -   -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)
<input type="radio"/> <i>Amsinckia lunaris</i>	Bent-flowered Fiddleneck	Boragin-	-   1B.2   *A2	- NA	Gravelly slopes, grassland, openings in woodland (Mar–Jun)
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	-   -   -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	-   -   -	- NP	Generally wooded areas (Jun–Aug)
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	-   -   -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	-   -   -	- ZA	Generally shady places (Apr–Jun)
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	-   -   -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)
<input type="radio"/> <i>Apocynum androsaemifolium</i>	Bitter Dogbane	Apocyn-	-   -   B	- NP	Open slopes, rocky places, with conifers, chaparral (May–Oct)
<input type="radio"/> <i>Aristolochia californica</i>	Dutchman's Pipe	Aristolochi-	-   -   C	- NP	Streamsides, forest, chaparral (Jan–Apr)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Artemisia</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Artemisia</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	-   -   -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	-   -   B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)	
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	-   -   -	- NP	Dry ground, valleys, foothills (May–Oct)	
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvetch	Fab-	-   -   -	- NA	Open, grassy areas, scrub (Mar–Jul)	
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	-   -   -	- NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	-   -   -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	-   -   -	L ZA	Disturbed grassland. (Apr–Jun)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	-   -   -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	-   -   -	M ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	-   -   -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	-   -   -	- NP	Common. Shady to open woodland, scrub (Apr–Jun)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	-   -   C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus umbellatus</i>	Oakland Star-tulip	Lili-	-   4.2   *A2	- NP	Open chaparral or woodland, gen on serpentine (Mar–May)	
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	-   -   B	- NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calystegia malacophylla subsp. pedicellata</i>	Woolly Morning-glory	Convolvul-	-   -   A2	- NP	Dry slopes, chaparral (Apr–Jul)	
<input type="radio"/> <i>Calystegia purpurata subsp. purpurata</i>	Climbing Morning-glory	Convolvul-	-   -   -	- NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	-   -   -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	-   -   -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	-   -   -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	-   -   -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	-   -   -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	-   -   -	- NA	Open fields, grassland (Mar–May)	
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	-   -   -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)	
<input type="radio"/> <i>Castilleja rubicundula subsp. lithospermoides</i>	Yellow Cream Sacs	Orobanch-	-   -   B	- NA	Open grassland (Apr–Jun)	

Herbaceous <i>Caulanthus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Caulanthus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i> )	California Mustard	Brassic-	-   -   -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	-   -   -	M ZAB	Pastures, disturbed places (Apr–Nov)
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	-   -   -	M ZA	Disturbed fields, open woodland (Apr–Jul)
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	-   -   -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	-   -   -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chamaesyce ocellata</i> subsp. <i>ocellata</i>	Valley Spurge	Euphorbi-	-   -   B	- NA	Dry, sandy places (May–Sep)
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	-   -   -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Aster-	-   -   -	- NB	Disturbed areas, grassland, woodland (May–Jul)
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	-   -   -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia affinis</i>	Small Clarkia	Onagr-	-   -   C	- NA	Openings in woodland, chaparral (May–Jun)
<input type="radio"/> <i>Clarkia concinna</i> subsp. ?	Red Ribbons	Onagr-	-   -   -	- NA	--
<input type="radio"/> <i>Clarkia purpurea</i> subsp. ?	Winecup Clarkia	Onagr-	-   -   -	- NA	--
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	-   -   -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	-   -   -	- NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	-   -   -	- NA	--
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	-   -   -	- NA	Common. Woodland (Apr–Sep)
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>exigua</i>	Common Pale Claytonia	Monti-	-   -   C	- NA	Dry or moist, disturbed bare clay to sandy soils, often serpentine (Apr–Jul)
<input type="radio"/> <i>Claytonia parviflora</i> subsp. ?	Miner's Lettuce	Monti-	-   -   -	- NA	--
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	-   -   -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i> )	Yerba Buena	Lami-	-   -   -	- NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. ?	White Chinese Houses	Plantagin-	-   -   -	- NA	--
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. <i>bartsiiifolia</i>	White Chinese Houses	Plantagin-	-   -   A1	- NA	Open, sandy places (Mar–Jun)
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	-   -   -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	-   -   C	- NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)

Herbaceous <i>Conium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Conium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	-   -   -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)	
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	-   -   -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)	
<input type="radio"/> <i>Corallorhiza maculata</i> var. <i>maculata</i> (JM93: no var.)	Spotted Coralroot	Orchid-	-   -   A2	- NP	Shaded mixed-evergreen or conifer forest, in decomposing lf litter (May–Aug)	
<input type="radio"/> <i>Cordylanthus pilosus</i> subsp. <i>pilosus</i>	Hairy Bird's Beak	Orobanch-	-   -   B	- NA	Open foothill woodland, chaparral, on serpentine (Jul–Sep)	
<input type="radio"/> <i>Corethrogyne filaginifolia</i> (JM93: <i>Lessingia</i> , var. <i>filaginifolia</i> )	Common California-aster	Aster-	-   -   C	- NP	Coastal scrub, chaparral, grassland, foothill woodland, forest (Jul–Nov)	
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	-   -   -	- NA	Open areas (Feb–May)	
<input type="radio"/> <i>Crepis vesicaria</i> subsp. <i>taraxacifolia</i>	Dandelion-leaf Hawksbeard	Aster-	-   -   -	- ZAB	Sandy clearings, hillsides, disturbed places (Feb–Oct)	
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i> )	Turkey-mullein	Euphorbi-	-   -   -	- NA	Dry, open, often disturbed areas (May–Oct)	
<input type="radio"/> <i>Cryptantha microstachys</i>	Hairy Cryptantha	Boragin-	-   -   A2	- NA		
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	-   -   -	M ZP	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	-   -   -	- NP	Chaparral, woodland (Feb–May)	
<input type="radio"/> <i>Datisca glomerata</i>	Durango Root	Datic-	-   -   A2	- NP	Dry streambeds or washes (May–Jul)	
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Ranuncul-	-   -   C	- NP	Open coastal grassland, chaparral (Mar–May)	
<input type="radio"/> <i>Delphinium nudicaule</i>	Red Or Orange Larkspur	Ranuncul-	-   -   C	- NP	Moist talus, wooded, rocky slopes (Mar–Jun)	
<input type="radio"/> <i>Delphinium variegatum</i> subsp. <i>variegatum</i>	Royal Larkspur	Ranuncul-	-   -   C	- NP	Grassland, open oak woodland (Mar–May)	
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	-   -   -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	-   -   -	- NP	Open woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	-   -   -	- NP	Gen in shady sites (Mar–Jul)	
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i> )	Sticky Cinquefoil	Ros-	-   -   -	- NP	Gen ± shady or moist areas (May–Jul)	
<input type="radio"/> <i>Emmenanthe penduliflora</i> var. <i>penduliflora</i>	Whispering Bells	Boragin-	-   -   B	- NP	Chaparral to creosote-bush scrub, rocky, sandy, decomposed granite, serpentine soils (Apr–Jul)	
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	-   -   -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)	
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	-   -   -	- NP	Dry slopes, ridges (Jun–Dec)	
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Aster-	-   -   -	- NP	Grassy dunes, chaparral, oak woodland (May–Oct)	
<input type="radio"/> <i>Erigeron petrophilus</i> var. <i>petrophilus</i>	Rock-loving Fleabane	Aster-	-   -   A2	- NP	Rocky foothills to montane forest, sometimes on serpentine (May–Sep)	
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	-   -   -	- NP	- -	



<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Eriophyllum</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Eriophyllum</i></b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
○ <i>Eriophyllum lanatum</i> var. <i>achilleoides</i> (JM93: var. <i>achillaeoides</i> )	Common Woolly Sunflower	Aster-	-   -   C	- NP	Dry, often rocky sites, chaparral, forest (May–Jul)	
○ <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)	
○ <i>Erodium brachycarpum</i>	Southern European Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)	
○ <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	-   -   -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)	
○ <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	-   -   -	- ZA	Open, disturbed sites (Feb–Sep)	
○ <i>Erysimum capitatum</i> var. <i>capitatum</i> (JM93: var. <i>capitatum</i> , var. <i>angustatum</i> )	Western Wallflower	Brassic-	-   -   C	- NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)	
○ <i>Eschscholzia californica</i>	California Poppy	Papaver-	-   -   -	- NP	Grassy, open areas (Feb–Sep)	
○ <i>Eucrypta chrysanthemifolia</i> var. <i>chrysanthemifolia</i>	Common Eucrypta	Boragin-	-   -   B	- NA	Roadsides, burns, coastal bluffs, ravines (Mar–Jun)	
○ <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i> )	Broadleaf Aster	Aster-	-   -   -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)	
○ <i>Fragaria vesca</i>	Wood Strawberry	Ros-	-   -   -	- NP	Gen partial shade in forest (Jan–Jul)	
○ <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i> )	Checker Lily	Lili-	-   -   -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)	
○ <i>Galium aparine</i>	Goose Grass	Rubi-	-   -   -	- NA	Grassy, ± shady places (Apr–Jun)	
○ <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	-   -   -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)	
○ <i>Galium murale</i>	Tiny Bedstraw	Rubi-	-   -   -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)	
○ <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	-   -   -	- NP	Among shrubs in chaparral, forest (May–Aug)	
○ <i>Geranium carolinianum</i>	Carolina Geranium	Gerani-	-   -   -	- NA	Open to shaded sites, grassland, scrub, forest (Feb–Aug)	
○ <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	-   -   -	M ZA	Open, disturbed sites (Mar–Jul)	
○ <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	-   -   -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)	
○ <i>Helenium puberulum</i>	Rosilla	Aster-	-   -   -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)	
○ <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	-   1B.2   *A2	- NP	Open, grassy sites (Apr–Jun)	
○ <i>Helminthotheca echioides</i> (JM93: <i>Picris</i> )	Bristly Ox-tongue	Aster-	-   -   -	L ZAB	Common. Disturbed areas (All year)	
○ <i>Hemizonia congesta</i> subsp. <i>lutescens</i> (JM93: subsp. <i>congesta</i> )	Yellow Hayfield Tarweed	Aster-	-   -   C	- NA	Grassland, barrens, openings in chaparral and woodland, often on serpentine (Apr–Dec)	
○ <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i> )	Cow Parsnip	Api-	-   -   -	- NP	Moist places, wooded or open (Apr–Jul)	
○ <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Aster-	-   -   -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)	

Herbaceous <i>Hesperocnide</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Hesperocnide</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	-   -   -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. ?	Sessileflower Goldenaster	Aster-	-   -   -	- NP	--
<input type="radio"/> <i>Heuchera micrantha</i>	Small-flower Alumroot	Saxifrag-	-   -   C	- NP	Moist, rocky banks and cliffs (Apr–Jul)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	-   -   -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	-   -   -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	-   -   -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	-   -   -	- ZA	Roadsides, grassland (Jul–Nov)
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	-   -   -	- ZA	Abundant. Disturbed places (May–Oct)
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	-   -   -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: subsp. <i>ramosissima</i> )	Common Hare-leaf	Aster-	-   -   -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lasthenia californica</i> subsp. <i>californica</i> (JM93: no subsp.)	California Goldfields	Aster-	-   -   -	- NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lasthenia glabrata</i> subsp. <i>glabrata</i>	Yellow-ray Goldfields	Aster-	-   -   B	- NA	Saline places, vernal pools (Mar–May)
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	-   -   -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	-   -   C	- NA	Many habitats (Feb–Jul)
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i> )	Threadleaf Peppergrass	Brassic-	-   -   -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)
<input type="radio"/> <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	-   -   -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i> )	Pinklobe Leptosiphon	Polemoni-	-   -   C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	-   -   -	- NP	Open, grassy slopes (Mar–Apr)
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	-   -   -	- NP	Shaded slopes (Feb–Jun)
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i> )	California Fluffweed	Aster-	-   -   -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i> )	Daggerleaf Cottonrose	Aster-	-   -   -	- ZA	Bare or grassy openings, burns (Mar–Jul)
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Api-	-   -   B	- NP	Woodland, brushy slopes (Apr–Jun)
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	-   -   -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)

Herbaceous <i>Lomatium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Lomatium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	-   -   -	- NP	Open grassy slopes, meadows, woodland (Feb–May)
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	-   -   -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)
<input type="radio"/> <i>Madia elegans</i> (JM93: <i>subsp. elegans</i> , <i>densiflora</i> , <i>vernalis</i> )	Elegant Tarweed	Aster-	-   -   C	- NA	
<input type="radio"/> <i>Madia exigua</i>	Thread-stem Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites, in sandy or clayey soils, including serpentine (Apr–Jul)
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	-   -   -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)
<input type="radio"/> <i>Madia</i> or <i>Anisocarpus</i> sp. (JM93: <i>Madia</i> sp.)	Tarweed	Aster-	-   -   -	- NA	--
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites (May–Oct)
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i> )	Starry False Solomon's Seal	Rusc-	-   -   -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Malv-	-   -   -	- ZA	Disturbed places (Mar–Jun)
<input type="radio"/> <i>Malva parviflora</i>	Cheeseweed	Malv-	-   -   -	- ZA	Common. Disturbed places (Mar–May)
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i> )	California Man-root	Cucurbit-	-   -   -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	-   -   -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i> )	Pineapple Weed	Aster-	-   -   -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	-   -   -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i> )	California Saxifrage	Saxifrag-	-   -   -	- NP	Moist, shady places (Feb–May(Jun))
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	-   -   -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i> )	Slender Annual Phlox	Polemoni-	-   -   -	- NA	Dry to moist areas (Mar–Aug)
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	-   -   -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i> )	Coyote-mint	Lami-	-   -   -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Nasturtium</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Nasturtium</i></b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i> )	Water Cress	Brassic-	-   -   -	-	NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	-   -   C	-	NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Polemoni-	-   -   C	-	NA	Open, slopes, gravel, clay (May–Jul)
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	-   -   -	-	NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	-   -   -	-	NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Boragin-	-   -   -	-	NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)
<input type="radio"/> <i>Nemophila parviflora</i> var. <i>parviflora</i>	Small-flower Nemophila	Boragin-	-   -   C	-	NA	Woodland, forest, roadsides, slopes (Mar–Jul)
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i> )	Sweet-cicely	Api-	-   -   -	-	NP	Conifer forest, woodland, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Papaver californicum</i>	Fire Poppy	Papaver-	-   -   A2	-	NA	Burns, disturbed areas, open woodland (Apr–May)
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i> )	Wind Poppy	Papaver-	-   -   -	-	NA	Grassy areas, openings in chaparral (Apr–May)
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	-   -   -	-	NP	Dry chaparral, oak/pine forest (Mar–May)
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	-   -   -	-	NP	Open grassland, serpentine outcrops (Jul–Aug)
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	-   -   -	-	NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	-   -   -	-	NA	Common. Clay to rocky soils, slopes (Mar–May)
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	-   -   -	-	NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)
<input type="radio"/> <i>Phacelia malvifolia</i>	Stinging Phacelia	Boragin-	-   -   A2	-	NA	Sandy or gravelly soils, scrub, conifer forest (Apr–Jul)
<input type="radio"/> <i>Pholistoma membranaceum</i>	White Fiesta Flower	Boragin-	-   -   -	-	NA	Beaches, bluffs, ravines, wooded slopes, desert washes (Feb–May)
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>macrophyllum</i> (JM93: <i>P. macrophyllum</i> )	Big-leaf Mistletoe	Visc-	-   -   C	-	NP	On trees other than <i>Quercus</i> (especially <i>Alnus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Platanus</i> , <i>Populus</i> , <i>Robinia</i> , <i>Salix</i> ) (Dec–Mar)
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>tomentosum</i> (JM93: <i>P. villosum</i> )	Oak Mistletoe	Visc-	-   -   -	-	NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Boragin-	-   -   -	-	NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	-   -   -	-	NA	Common; open woodland, grassland (Mar–May)
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	-   -   -	-	- -	

Herbaceous <i>Plantago</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Plantago</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	-   -   -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	-   -   -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	-   -   -	- NA	Open grassland, sandy soil, burns (Mar–May)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: <i>subsp. ciliosa, insignis</i> )	Longspur Plectritis	Valerian-	-   -   B	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	-   -   -	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	-   -   -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Polycarpon tetraphyllum</i>	Four-leaved Allseed	Caryophyll-	-   -   -	- ZA	Disturbed shaded areas, roadsides (Spring–fall)	
<input type="radio"/> <i>Polygonum aviculare subsp. depressum</i> (JM93: <i>P. arenastrum</i> )	Knotweed	Polygon-	-   -   -	- ZA	Disturbed places (May–Nov)	
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i> )	Hooker Fairy Bells	Lili-	-   -   C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)	
<input type="radio"/> <i>Prunella vulgaris</i> var. ?	European Selfheal	Lami-	-   -   -	- ZP	--	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i> )	California Everlasting	Aster-	-   -   -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i> )	Cotton-batting Plant	Aster-	-   -   -	- NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	-   -   -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: <i>no var.</i> )	California Buttercup	Ranuncul-	-   -   -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	-   -   -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	-   -   -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)	
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	-   -   -	L ZA	Disturbed areas, fields (May–Jul)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	-   -   -	M ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	-   -   -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	-   -   -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rumex</i> sp.	Dock	Polygon-	-   -   -	-	--	
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	-   -   -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	-   -   -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	-   -   -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula tuberosa</i>	Turkey Pea Sanicle	Api-	-   -   C	- NP	Open gravelly meadows, chaparral, woodland, pine forest (Mar–Jul)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Scandix</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Scandix</b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	-   -   -	- ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: <i>subsp. californica, floribunda</i> )	California Figwort	Scrophulari-	-   -   -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberosus Skullcap	Lami-	-   -   C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sedum spathulifolium</i>	Broadleaf Stonecrop	Crassul-	-   -   B	- NP	Outcrops, often in shade (Apr–Aug)	
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	-   -   C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	-   -   -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	-   -   -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sidalcea malviflora subsp. ?</i>	Common Checkerbloom	Malv-	-   -   -	- NP	--	
<input type="radio"/> <i>Sidalcea malviflora subsp. malviflora</i>	Common Checkerbloom	Malv-	-   -   B	- NP	Coastal prairie, scrub, open forest (Mar–Jul)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	-   -   -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	-   -   -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sinapis arvensis</i>	Charlock	Brassic-	-   -   -	L ZA	Disturbed areas (Mar–Sep)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	-   -   -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	-   -   -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solanum nigrum</i>	Black Nightshade	Solan-	-   -   -	- ZA	Disturbed places (Mar–Oct)	
<input type="radio"/> <i>Solidago velutina subsp. californica</i> (JM93: <i>S. californica</i> )	California Goldenrod	Aster-	-   -   -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	-   -   -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper subsp. asper</i>	Prickly Sow Thistle	Aster-	-   -   -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	-   -   -	- ZA	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Stachys rigida var. quercetorum</i> (JM93: <i>S. ajugoides var. rigida</i> )	Common Rigid Hedge-nettle	Lami-	-   -   -	- NP	Moist to ± dry places (Mar–Oct)	
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	-   -   -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	-   -   C	- NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)	
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	-   -   -	- ZBP	Abundant. Esp disturbed areas (All year)	
<input type="radio"/> <i>Taraxia ovata</i> (JM93: <i>Camissonia</i> )	Golden Eggs Suncup	Onagr-	-   -   C	- NP	Grassy fields, gen clay soil (Mar–Jun)	
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	-   -   C	- NP	Chaparral, pine/oak woodland (Mar–May)	

Herbaceous <i>Thalictrum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Thalictrum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	-   -   -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	-   -   -	- NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)	
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	-   -   -	M ZA	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	-   -   -	- ZA	Disturbed places (Apr–Jun)	
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i> )	Common Star Lily	Melanthi-	-   -   -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)	
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	-   -   -	- ZBP	Common. Disturbed places (Mar–Nov)	
<input type="radio"/> <i>Trichostema lanceolatum</i>	Vinegar Weed	Lami-	-   -   -	- NA	Dry, open, gen disturbed habitats (Jun–Nov)	
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i> )	Rancheria Clover	Fab-	-   -   C	- NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Fab-	-   -   -	- NA	Open, grassy areas, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	-   -   -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Fab-	-   -   -	- NA	Grassland, coastal woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Fab-	-   -   -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	-   -   -	- ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	-   -   -	M ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	-   -   -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium obtusiflorum</i>	Clammy Clover	Fab-	-   -   B	- NA	Moist disturbed areas, gravel bars, marshes (Apr–Jul)	
<input type="radio"/> <i>Trifolium olivaceum</i> (JM93: <i>T. albopurpureum</i> var. <i>olivaceum</i> )	Olive Clover	Fab-	-   -   B	- NA	Grassy slopes, valley meadows, disturbed areas (Apr–May)	
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Fab-	-   -   -	- ZA	Meadows, roadsides, disturbed areas (Mar–Apr)	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	-   -   -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	-   -   -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	-   -   -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	-   -   -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	-   -   -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Brassic-	-   -   B	- NP	Grassy banks, open fields, roadsides, pastures (Mar–May)	

Herbaceous <i>Uropappus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Uropappus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	-   -   -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urtica dioica subsp. holosericea</i>	Hoary Nettle	Urtic-	-   -   -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	-   -   -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	-   -   -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Veronica americana</i>	American Brooklime	Plantagin-	-   -   C	- NP	Common. Moist to wet soil, springs, slow streams, meadows, lakeshores (May–Aug)	
<input type="radio"/> <i>Veronica peregrina subsp. xalapensis</i>	Purslane Speedwell	Plantagin-	-   -   -	- NA	Moist places (Apr–Aug)	
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	-   -   -	- ZA	Wet, disturbed areas, fields (Feb–May)	
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	-   -   -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sp.</i>	Vetch	Fab-	-   -   -	- - -		
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)	
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Apocyn-	-   -   -	M ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	-   -   C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	-   -   -	- NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	-   -   C	- NP	Gen shady sites (Mar–Jun)	
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	-   -   -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))	
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	-   -   -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Api-	-   -   C	- NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)	



<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Acer</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Acer</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	-   -   -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	-   -   -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	-   -   -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	-   -   -	- NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	-   -   C	- NP	Conifer, oak forests (Mar–May)	
<input type="radio"/> <i>Arctostaphylos crustacea</i> subsp. <i>crustacea</i> (JM93: <i>A. tomentosa</i> subsp. <i>crustacea</i> )	Brittle-leaf Manzanita	Eric-	-   -   C	- NP	Chaparral, conifer forest (Feb–Apr)	
<input type="radio"/> <i>Arctostaphylos glandulosa</i> subsp. <i>glandulosa</i>	Eastwood Manzanita	Eric-	-   -   B	- NP	Chaparral, conifer forest (Jan–Apr)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Eric-	-   -   C	- NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Arctostaphylos</i> sp.	Manzanita	Eric-	-   -   -	- NP	--	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	-   -   -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	-   -   -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Berberis pinnata</i> subsp. <i>pinnata</i>	Shinyleaf Oregon-grape	Berberid-	-   -   C	- NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
<input type="radio"/> <i>Berberis</i> sp.	Oregon Grape	Berberid-	-   -   -	- NP	--	
<input type="radio"/> <i>Ceanothus cuneatus</i> var. <i>cuneatus</i>	Buckbrush	Rhamn-	-   -   C	- NP	Sandy to rocky flats, slopes, ridges (Feb–May)	
<input type="radio"/> <i>Ceanothus oliganthus</i> var. <i>sorediatus</i>	Jim Brush	Rhamn-	-   -   -	- NP	Slopes, ridges, chaparral, conifer forest (Jan–May)	
<input type="radio"/> <i>Ceanothus thyrsiflorus</i> var. <i>thyrsiflorus</i> (JM93: no var.)	Blue Blossom	Rhamn-	-   -   -	- NP	Bluffs, slopes, canyons, chaparral, coastal scrub, closed-cone-pine forest (Mar–Jun)	
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	-   -   C	- NP	Dry, rocky slopes, chaparral (Mar–May)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	-   -   -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Corn-	-   -   C	- NP	Generally moist places (May–Jul)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Corn-	-   -   C	- NP	Generally moist habitats (May–Jul)	
<input type="radio"/> <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i> )	California Hazelnut	Betul-	-   -   -	- NP	Common. Many habitats, esp moist, shady places (Jan–Mar)	
<input type="radio"/> <i>Eriodictyon californicum</i>	California Yerba Santa	Boragin-	-   -   -	- NP	Slopes, fields, roadsides, woodland, chaparral (Apr–Jul)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	-   -   -	- NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	-   -   -	M ZP	Common. Disturbed areas (Oct–Jan)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i> )	California Coffee Berry	Rhamn-	-   -   -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Frangula</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Frangula</b>
✓ <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>tomentella</i> (JM93: <i>Rhamnus tomentella</i> subsp. <i>tomentella</i> )	Hoary Coffee Berry	Rhamn-	-   -   B	- NP	Chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Garrya elliptica</i>	Coast Silk Tassel	Garry-	-   -   B	- NP	Seacliffs, sand dunes, chaparral, foothill-pine woodland (Jan–Mar)	
<input type="radio"/> <i>Helianthemum scoparium</i>	Peak Rush-rose	Cist-	-   -   B	- NP	Dry sandy or rocky soil of hills, slopes, ridges (Feb–Sep)	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	-   -   -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	-   -   -	- NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i> )	N. California Black Walnut	Jugland-	-   1B.1   *A2	- NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Juniperus californica</i>	California Juniper	Cupress-	-   -   B	- NP	Dry slopes, flats, pinyon/juniper woodland -	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i> )	Hairy Vine Honeysuckle	Caprifoli-	-   -   -	- NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	-   -   -	- NP	- -	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	-   -   -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	-   -   -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	-   -   B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Ros-	-   -   C	- NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	-   -   A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	-   -   C	- NP	Dry slopes, chaparral (Feb–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	-   -   -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	-   -   -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	
<input type="radio"/> <i>Quercus garryana</i> var. <i>garryana</i>	Oregon / Garry Oak	Fag-	-   -   B	- NP	Slopes, mixed-evergreen or conifer forest (Apr–Jun)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	-   -   -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	-   -   B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>frutescens</i>	Dwarf Interior Live Oak	Fag-	-   -   C	- NP	Valleys, chaparral (Mar–May)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	-   -   -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Rhamnus</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Rhamnus</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Rhamn-	-   -   C	- NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	-   -   C	- NP	Chaparral, montane forest (Mar–Jun)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	-   -   -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes malvaceum</i> var. <i>malvaceum</i>	Chaparral Currant	Grossulari-	-   -   B	- NP	Chaparral, oak woodland (Oct–Apr)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	-   -   -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	-   -   -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	-   -   -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa spithamea</i>	Coast Ground Rose	Ros-	-   -   C	- NP	Open forest, chaparral, especially after fire (Apr–Aug)	
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i> )	Himalayan Blackberry	Ros-	-   -   -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	-   -   -	- NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	-   -   -	- NP	Open, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	-   -   -	- NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="radio"/> <i>Salvia mellifera</i>	Black Sage	Lami-	-   -   -	- NP	Coastal-sage scrub, lower chaparral (Mar–Jun)	
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i> )	Blue Elderberry	Adox-	-   -   -	- NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	-   -   -	- NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="radio"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	-   -   -	- NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="radio"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	-   -   -	- NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	-   -   -	- NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="radio"/> <i>Umbellularia californica</i>	California Bay	Laur-	-   -   -	- NP	Common. Canyons, valleys, chaparral (Nov–May)	
<input type="radio"/> <i>Vitis californica</i>	California Wild Grape	Vit-	-   -   -	- NP	Streamsides, springs, canyons (May–Jun)	

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Acer negundo* var. *californicum* (no var.)

*Adenostoma fasciculatum* (var. *fasciculatum*)

*Agoseris grandiflora* (var. *grandiflora*)

*Agoseris heterophylla* (var. *cryptopleura*)

*Agrostis viridis* (*Polypogon*)

*Alisma plantago-aquatica* (*A. triviale*)

*Amsinckia menziesii* var. *intermedia* (*A. intermedia*)

*Amsinckia menziesii* var. *menziesii* (no var.)

*Antirrhinum vexillo-calyculatum* ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

*Arabis breweri* var. *breweri* (*Boechera breweri* subsp. *breweri*)

*Arabis glabra* var. *glabra* (*Turritis glabra*)

*Arctostaphylos tomentosa* ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

*Aster chilensis* (*Symphyotrichum chilense*)

*Aster lentus* (*Symphyotrichum lentum*)

*Aster radulinus* (*Eurybia radulina*)

*Aster subulatus* var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

*Atriplex joaquiniana* (*A. joaquinana*)

*Atriplex lentiformis* ssp. *lentiformis* (no subsp.)

*Atriplex triangularis* (*A. prostrata*)

*Azolla mexicana* (*A. microphylla*)

*Baccharis douglasii* (*B. glutinosa*)

*Baccharis pilularis* (subsp. *consanguinea*)

*Baccharis salicifolia* (subsp. *salicifolia*)

*Balsamorhiza macrolepis* var. *macrolepis* (no var.)

*Blepharizonia plumosa* ssp. *plumosa* (no subsp.)

*Blepharizonia plumosa* ssp. *viscida* (*Blepharizonia laxa*)

*Boschniakia strobilacea* (*Kopsiopsis*)

*Bromus catharticus* (var. *catharticus*)

*Camissonia boothii* ssp. *decorticans* (*Eremothera*)

*Camissonia graciliflora* (*Tetrapteron graciliflorum*)

*Camissonia intermedia* (*Camissoniopsis*)

*Camissonia micrantha* (*Camissoniopsis*)

*Camissonia ovata* (*Taraxia*)

*Cardaria draba* (*Lepidium*)

*Carduus pycnocephalus* (subsp. *pycnocephalus*)

*Carex deweyana* ssp. *leptopoda* (*C. leptopoda*)

*Carex lanuginosa* (*C. pellita*)

*Carex ovalis* (*C. leporina*)

*Ceanothus thyrsoiflorus* (var. *thyrsoiflorus*)

*Centaurium davyi* (*Zeltnera*)

*Centaurium muehlenbergii* (*Zeltnera*)

*Cephalanthus occidentalis* var. *californicus* (no var.)

*Chamomilla suaveolens* (*Matricaria discoidea*)

*Chenopodium ambrosioides* (*Dysphania*)

*Chenopodium macrospermum* var. *halophilum* (no var.)

*Chenopodium multifidum* (*Dysphania multifida*)

*Chenopodium rubrum* (var. *rubrum*)

*Chrysothamnus nauseosus* ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

*Cirsium cymosum* (var. *cymosum*)

*Claytonia parviflora* ssp. *parviflora* (subsp. *utahensis*)

*Collinsia bartsifolia* var. *davidsonii* (var. *stricta*)

*Collinsia heterophylla* (var. *heterophylla*)

*Conyza bilbaoana* (*Erigeron sumatrensis*)

*Conyza bonariensis* (*Erigeron*)

*Conyza canadensis* (*Erigeron*)

*Conyza coulteri* (*Laennecia*)

*Corallorhiza maculata* (var. *maculata*)

*Cordylanthus mollis* ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

*Coronopus didymus* (*Lepidium didymum*)

*Corylus cornuta* var. *californica* (subsp. *californica*)

*Cotoneaster pannosa* (*C. pannosus*)

*Cryptantha decipiens* (*C. rattanii*)

*Cryptantha intermedia* (var. *intermedia*)

*Cryptantha muricata* (var. *muricata*)

*Cryptantha torreyana* (var. *pumila*)

*Cupressus macrocarpa* (*Hesperocyparis*)

*Cuscuta salina* var. *major* (*C. pacifica* var. *pacifica*)

*Cynara cardunculus* (subsp. *flavescens*)

*Danthonia californica* var. *californica* (no var.)

*Dicentra chrysantha* (*Ehrendorferia*)

*Disporum hookeri* (*Prosartes*)

*Disporum smithii* (*Prosartes*)

*Draba cuneifolia* var. *integrifolia* (no var.)

*Duchesnea indica* (no var.)

*Epilobium pygmaeum* (*E. campestre*)

*Eragrostis curvula* var. *curvula* (no var.)

*Erechtites glomerata* (*Senecio glomeratus*)

*Erechtites minima* (*Senecio minimus*)

*Eremocarpus setigerus* (*Croton*)

*Erigeron philadelphicus* (var. *philadelphicus*)

*Eriogonum umbellatum* var. *bahiiforme* (var. *smallianum*)

*Eriophyllum lanatum* var. *achillaeoides* (var. *achilleoides*)

*Eryngium aristulatum* var. *aristulatum* (*E. jepsonii*)

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

*Erysimum capitatum* var. *angustatum* (var. *capitatum*)

*Filago californica* (*Logfia filaginoides*)

*Filago gallica* (*Logfia*)

*Fritillaria affinis* var. *affinis* (no var.)

*Galium trifidum* var. *pacificum* (subsp. *columbianum*)

*Gastridium ventricosum* (*G. phleoides*)

*Gnaphalium bicolor* (*Pseudognaphalium biolettii*)

*Gnaphalium californicum* (*Pseudognaphalium*)

*Gnaphalium canescens* ssp. *beneolens*  
(*Pseudognaphalium beneolens*)

*Gnaphalium canescens* ssp. *microcephalum*  
(*Pseudognaphalium microcephalum*)

*Gnaphalium luteo-album* (*Pseudognaphalium luteoalbum*)

*Gnaphalium purpureum* (*Gamochaeta ustulata*)

*Gnaphalium ramosissimum* (*Pseudognaphalium*)

*Gnaphalium stramineum* (*Pseudognaphalium*)

*Grindelia camporum* var. *camporum* (no var.)

*Grindelia hirsutula* var. *davyi* (*G. camporum*)

*Grindelia hirsutula* var. *hirsutula* (no var.)

*Guillenia flavescens* (*Caulanthus*)

*Guillenia lasiophylla* (*Caulanthus lasiophyllus*)

*Hedera helix* ssp. *helix* (no subsp.)

*Heliotropium curassavicum* (var. *oculatum*)

*Hemizonia congesta* ssp. *congesta* (subsp. *lutescens*)

*Hemizonia corymbosa* ssp. *corymbosa* (*Deinandra corymbosa*)

*Hemizonia fitchii* (*Centromadia*)

*Hemizonia lobbii* (*Deinandra*)

*Hemizonia parryi* ssp. *congdonii* (*Centromadia*)

*Hemizonia parryi* ssp. *parryi* (*Centromadia*)

*Hemizonia pungens* ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

*Hemizonia pungens* ssp. *pungens* (*Centromadia*)

*Heracleum lanatum* (*H. maximum*)

*Herniaria hirsuta* ssp. *cinerea* (var. *cinerea*)

*Holodiscus discolor* (var. *discolor*)

*Hordeum jubatum* (subsp. *jubatum*)

*Horkelia californica* ssp. *californica* (var. *californica*)

*Horkelia californica* ssp. *dissita* (var. *elata*)

*Horkelia californica* ssp. *frondosa* (var. *frondosa*)

*Isopyrum occidentale* (*Enemion*)

*Iva axillaris* ssp. *robustior* (no subsp.)

*Juglans californica* var. *hindsii* (*J. hindsii*)

*Juncus balticus* (subsp. *ater*)

*Juncus effusus* var. *pacificus* (subsp. *pacificus*)

*Juncus lesueurii* (*J. lescurii*)

*Koeleria phleoides* (*K. gerardii*)

*Lagophylla ramosissima* ssp. *ramosissima* (no subsp.)

*Lasthenia californica* (subsp. *californica*)

*Lavatera cretica* (*Malva pseudolavatera*)

*Lepidium dictyotum* var. *dictyotum* (no var.)

*Lepidium latipes* var. *latipes* (no var.)

*Lepidium nitidum* var. *nitidum* (no var.)

*Lepidium nitidum* var. *oreganum* (no var.)

*Leptochloa fascicularis* (*L. fusca* subsp. *fascicularis*)

*Lessingia filaginifolia* var. *californica* (*Corethrogyne filaginifolia*)

*Lessingia filaginifolia* var. *filaginifolia*  
(*Corethrogyne filaginifolia*)

*Lewisia rediviva* (var. *rediviva*)

*Leymus condensatus* (*Elymus*)

*Leymus triticoides* (*Elymus*)

*Lilaea scilloides* (*Triglochin*)

*Limosella subulata* (*Limosella australis*)

*Linanthus acicularis* (*Leptosiphon*)

*Linanthus ambiguus* (*Leptosiphon*)

*Linanthus androsaceus* (*Leptosiphon*)

*Linanthus bicolor* (*Leptosiphon*)

*Linanthus ciliatus* (*Leptosiphon*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus parviflorus* (*Leptosiphon*)

*Linaria canadensis* (*Nuttallanthus texanus*)

*Lolium multiflorum* (*Festuca perennis*)

*Lolium perenne* (*Festuca perennis*)

*Lolium temulentum* (*Festuca temulenta*)

*Lonicera hispidula* var. *vacillans* (no var.)

*Lotus humistratus* (*Acmispon brachycarpus*)

*Lotus micranthus* (*Acmispon parviflorus*)

*Lotus purshianus* var. *purshianus* (*Acmispon americanus* var. *americanus*)

*Lotus scoparius* var. *scoparius* (*Acmispon glaber* var. *glaber*)

*Lotus strigosus* (*Acmispon*)

*Lotus wrangelianus* (*Acmispon*)

*Luzula comosa* (var. *comosa*)

*Madia madioides* (*Anisocarpus*)

*Malacothamnus fasciculatus* (var. *nuttallii*)

*Marah fabaceus* (*M. fabacea*)

*Marah oreganus* (*M. oregana*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Melilotus alba* (*M. albus*)

*Melilotus indica* (*M. indicus*)

*Mentha spicata* var. *spicata* (no var.)

*Mimulus aurantiacus* (var. *aurantiacus*)

*Mirabilis californica* (*M. laevis* var. *crassifolia*)

*Monardella antonina* ssp. *antonina* (*M. villosa* subsp. *villosa*)

*Monardella douglasii* ssp. *douglasii* (no subsp.)

*Monardella villosa* ssp. *globosa* (subsp. *villosa*)

*Montia fontana* ssp. *chondrosperma* (no subsp.)

*Myrica californica* (*Morella*)

*Nassella cernua* (*Stipa*)

*Nassella lepida* (*Stipa*)

*Nassella pulchra* (*Stipa*)

*Osmorhiza chilensis* (*O. berteroi*)

*Oxalis albicans* ssp. *pilosa* (*O. pilosa*)

*Parvisedum pentandrum* (*Sedella pentandra*)

*Phacelia ramosissima* var. *latifolia* (no var.)

*Phacelia ramosissima* var. *ramosissima* (no var.)

*Phlox gracilis* (*Microsteris*)

*Phoradendron macrophyllum* (*P. serotinum* subsp. *macrophyllum*)

*Phoradendron villosum* (*P. serotinum* subsp. *tomentosum*)

*Phyla nodiflora* var. *incisa* (no var.)

*Phyla nodiflora* var. *nodiflora* (no var.)

*Picris echioides* (*Helminthotheca*)

*Piperia elegans* (subsp. *elegans*)

*Piptatherum miliaceum* (*Stipa miliacea* var. *miliacea*)

*Plagiobothrys canescens* (var. *canescens*)

*Plagiobothrys fulvus* (var. *campestris*)

*Plectritis brachystemon* (*P. congesta* subsp. *brachystemon*)

*Plectritis ciliosa* ssp. *ciliosa* (no subsp.)

*Plectritis ciliosa* ssp. *insignis* (no subsp.)

*Plectritis congesta* (subsp. *congesta*)

*Pleuropogon californicus* (var. *californicus*)

*Pluchea odorata* (var. *odorata*)

*Poa bulbosa* (subsp. *vivipara*)

*Polygonum amphibium* var. *emersum* (*Persicaria* *amphibia*)

*Polygonum amphibium* var. *stipulaceum* (*Persicaria* *amphibia*)

*Polygonum arenastrum* (*P. aviculare* subsp. *depressum*)

*Polygonum hydropiperoides* (*Persicaria*)

*Polygonum lapathifolium* (*Persicaria lapathifolia*)

*Polygonum persicaria* (*Persicaria maculosa*)

*Polygonum punctatum* (*Persicaria punctata*)

*Populus balsamifera* ssp. *trichocarpa* (*P. trichocarpa*)

*Potamogeton pectinatus* (*Stuckenia pectinata*)

*Potentilla glandulosa* ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

*Psilocarphus tenellus* var. *tenellus* (no var.)

*Ranunculus aquatilis* var. *capillaceus* (var. *diffusus*)

*Ranunculus californicus* (var. *californicus*)

*Ranunculus canus* (var. *canus*)

*Ranunculus occidentalis* (var. *occidentalis*)

*Ranunculus sceleratus* (var. *sceleratus*)

*Rhamnus californica* ssp. *californica* (*Frangula*)

*Rhamnus tomentella* ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

*Rhus trilobata* (*R. aromatica*)

*Ribes menziesii* (var. *menziesii*)

*Rorippa nasturtium-aquaticum* (*Nasturtium officinale*)

*Rorippa palustris* var. *occidentalis* (subsp. *palustris*)

*Rosa gymnocarpa* (var. *gymnocarpa*)

*Rubus discolor* (*R. armeniacus*)

*Rumex maritimus* (*R. fueginus*)

*Rumex salicifolius* var. *crassus* (*R. crassus*)

*Rumex salicifolius* var. *salicifolius* (no var.)

*Rumex salicifolius* var. *transitorius* (*R. transitorius*)

*Salicornia europaea* (*S. depressa*)

*Salicornia subterminalis* (*Arthrocnemum subterminale*)

*Salicornia virginica* (*S. pacifica*)

*Salix exigua* (var. *hindsiana*)

*Salix lucida* ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

*Sambucus mexicana* (*S. nigra* subsp. *caerulea*)

*Satureja douglasii* (*Clinopodium*)

*Saxifraga californica* (*Micranthes*)

*Scirpus acutus* var. *occidentalis* (*Schoenoplectus*)

*Scirpus americanus* (*Schoenoplectus*)

*Scirpus californicus* (*Schoenoplectus*)

*Scirpus cernuus* (*Isolepis*)

*Scirpus maritimus* (*Bolboschoenus maritimus* subsp. *paludosus*)

*Scirpus robustus* (*Bolboschoenus*)

*Scrophularia californica* ssp. *californica* (no subsp.)

*Scrophularia californica* ssp. *floribunda* (no subsp.)

*Senecio breweri* (*Packera*)

*Senecio mikanioides* (*Delairea odorata*)

*Silene californica* (*S. laciniata* subsp. *californica*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Smilacina racemosa* (*Maianthemum racemosum*)

*Smilacina stellata* (*Maianthemum stellatum*)

*Solidago californica* (*S. velutina* subsp. *californica*)

*Solidago canadensis* ssp. *elongata* (*S. elongata*)

*Sparganium erectum* ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

*Sparganium eurycarpum* ssp. *eurycarpum* (var. *eurycarpum*)

*Spergula arvensis* ssp. *arvensis* (no subsp.)

*Spergularia media* ( var. *media*)

*Stachys ajugoides* var. *ajugoides* (no var.)

*Stachys ajugoides* var. *rigida* (*Stachys rigida* var. *quercetorum*)

*Streptanthus albidus* ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

*Stylomecon heterophylla* (*Papaver heterophyllum*)

*Taeniatherum caput-medusae* (*Elymus*)

*Thysanocarpus laciniatus* (var. *laciniatus*)

*Tolmiea menziesii* (*T. diplomenziesii*)

*Trifolium albopurpureum* var. *albopurpureum* (no var.)

*Trifolium albopurpureum* var. *dichotomum* (*T. dichotomum*)

*Trifolium albopurpureum* var. *olivaceum* (*T. olivaceum*)

*Trifolium barbigerum* var. *andrewsii* (*T. grayi*)

*Trifolium barbigerum* var. *barbigerum* (no var.)

*Trifolium gracilentum* var. *gracilentum* (no var.)

*Vicia americana* var. *americana* (subsp. *americana*)

*Vicia ludoviciana* var. *ludoviciana* (subsp. *ludoviciana*)

*Viola adunca* (var. *adunca*)

*Vulpia bromoides* (*Festuca*)

*Vulpia microstachys* var. *ciliata* (*Festuca microstachys*)

*Vulpia microstachys* var. *confusa* (*Festuca microstachys*)

*Vulpia microstachys* var. *microstachys* (*Festuca microstachys*)

*Vulpia microstachys* var. *pauciflora* (*Festuca microstachys*)

*Vulpia myuros* var. *hirsuta* (*Festuca myuros*)

*Vulpia myuros* var. *myuros* (*Festuca myuros*)

*Vulpia octoflora* var. *hirtella* (*Festuca octoflora*)

*Vulpia octoflora* var. *octoflora* (*Festuca octoflora*)

*Zigadenus fremontii* (*Toxicoscordion*)