

Morgan Territory Regional Preserve

Checklist of Wild Plants

Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Morgan Territory Regional Preserve. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Family	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

Morgan Territory Regional Preserve

Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name					<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	- - -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Aspidotis californica</i>	California Lace Fern	Pterid-	- - A2	- NP	Rock outcrops, crevices -	
<input type="radio"/> <i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	Western Lady Fern	Woods-	- - -	- NP	Woodland, along streams, seepage area -	
<input type="radio"/> <i>Cystopteris fragilis</i>	Brittle Fern	Woods-	- - C	- NP	Shady, moist rock crevices, meadows, streamsides -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	- - -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	- - -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Pterid-	- - C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Pterid-	- - -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	- - -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	- - -	- NP	Near streams, springs, seeps -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	- - -	-	NP	Common. Open meadows, woodland, forest, subalpine (Jun–Aug)
<input type="radio"/> <i>Agrostis sp.</i>	Bent Grass	Po-	- - -	-	--	
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	- - -	M	ZA	Disturbed sites (Mar–Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	- - -	M	ZA	Disturbed sites (Apr–Jun)
<input type="radio"/> <i>Brachypodium distachyon</i>	Purple False-brome	Po-	- - -	M	ZA	Disturbed areas, dry slopes (Apr–Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	- - -	-	ZA	Shaded or moist, open sites (Apr–Jul)
<input type="radio"/> <i>Bromus carinatus var. carinatus</i>	California Brome	Po-	- - -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr–Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	- - -	M	ZA	Open, gen disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	- - -	L	ZA	Fields, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus madritensis subsp. madritensis</i>	Foxtail Chess	Po-	- - -	-	ZA	Disturbed areas, roadsides (Apr–Jan)
<input type="radio"/> <i>Bromus madritensis subsp. rubens</i>	Red Brome	Po-	- - -	H	ZA	Disturbed areas, roadsides (Mar–Jun)
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	- - B	-	NP	Seasonally wet places (May–Aug)
<input type="radio"/> <i>Crypsis schoenoides</i>	Swamp Prickle Grass	Po-	- - -	-	ZA	Wet places (Jun–Oct)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	- - -	M	ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	- - -	L	ZP	Disturbed, often moist sites (May–Aug)
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Po-	- - -	-	NP	Salt marshes, coastal dunes, moist, alkaline areas (Apr–Sep)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	- - -	-	NP	Common. Fresh to brackish wetland (Spring–summer)
<input type="radio"/> <i>Elymus caput-medusae (JM93: Taeniatherum)</i>	Medusa Head	Po-	- - -	H	ZA	Disturbed areas (Apr–Jul)
<input type="radio"/> <i>Elymus glaucus subsp. glaucus</i>	Western Wild-rye	Po-	- - -	-	NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	- - C	-	NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus stebbinsii</i>	Parish's Wheat Grass	Po-	- - A2	-	NP	Dry slopes, chaparral, conifer forest (Jun–Jul)
<input type="radio"/> <i>Elymus trachycaulus subsp. trachycaulus</i>	Slender Wheat Grass	Po-	- - B	-	NP	Dry to moist, open areas, forest, woodland (Jun–Aug)
<input type="radio"/> <i>Elymus triticoides (JM93: Leymus)</i>	Beardless Wild Rye	Po-	- - -	-	NP	Dry to moist, often saline, meadows (Jun–Jul)
<input type="radio"/> <i>Festuca bromoides (JM93: Vulpia)</i>	Brome Fescue	Po-	- - -	-	ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Po-	- - C	-	NP	Dry, open or shady places (Jul–Sep)
<input type="radio"/> <i>Festuca microstachys (JM93: Vulpia, var. ciliata, confusa, microstachys, pauciflora)</i>	Hairy Fescue	Po-	- - -	-	NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros (JM93: Vulpia, var. hirsuta, myuros)</i>	Rattail Sixweeks Grass	Po-	- - -	-	ZA	Common. Generally open places, sandy soils (Feb–May)

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i>)	Rye Grass	Po-	- - -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca</i> sp. (perennial)	Perennial Fescue	Po-	- - -	- -	
<input type="radio"/> <i>Gastridium phleoides</i> (JM93: <i>G. ventricosum</i>)	Nit Grass	Po-	- - -	- ZA	Open, generally dry, disturbed sites (Apr–Nov)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	- - -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	- - -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	- - -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>murinum</i>	Wall Barley	Po-	- - -	- ZA	Moist, gen disturbed sites (Feb–May)
<input type="radio"/> <i>Juncus bufonius</i> var. ?	Toad Rush	Junc-	- - -	- NA	- -
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Junc-	- - -	- NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Junc-	- - -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus occidentalis</i>	Western Rush	Junc-	- - -	- NP	Moist gen sunny areas (May–Sep)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	- - -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus tenuis</i>	Slender Rush	Junc-	- - -	- NP	Uncommon. Damp places (Jun–Sep)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	- - -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	- - C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	- - -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	- - C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	- - -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	- - -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Phalaris aquatica</i>	Harding Grass	Po-	- - -	M ZP	Disturbed areas, roadsides (Apr–Aug)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	- - -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa bulbosa</i> subsp. <i>vivipara</i> (JM93: no subsp.)	Bulbous Blue Grass	Po-	- - -	- ZP	Disturbed places (Mar–Jul)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	- - -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Po-	- - -	- ZP	Common. Streambanks (May–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	- - -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i>)	Foothill Needle Grass	Po-	- - C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Po-	- - C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like
<i>Triglochin</i>	Sorted Alphabetically by Scientific Name				<i>Triglochin</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Triglochin scilloides</i> (JM93: <i>Lilaea</i>)	Flowering-quillwort	Juncagin-	- - C	- NA	Vernal pools, streams, ponds, lake margins (Mar–Oct)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	- - -	- NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Achillea</i>	Sorted Alphabetically by Scientific Name					<i>Achillea</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	- - -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Aster-	- - -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)	
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i>)	Spanish Clover	Fab-	- - -	- NA		
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Fab-	- - -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Fab-	- - -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i>)	Small-flower Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	- - -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Aster-	- - -	- NA	Many open habitats (May–Jun)	
<input type="radio"/> <i>Allium amplexans</i>	Narrowleaf Onion	Alli-	- - A2	- NP	Clays including serpentine, open or wooded places (Apr–Jul)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	- - -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Alli-	- - C	- NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)	
<input type="radio"/> <i>Amaranthus californicus</i>	California Amaranth	Amaranth-	- - A2	- NA	Seasonally moist flats, lake margins, disturbed areas (Jul–Oct)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Boragin-	- - -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Amsinckia menziesii</i> (JM93: var. <i>menziesii</i>)	Small-flowered Fiddleneck	Boragin-	- - -	- NA	Shade-tolerant, open, disturbed areas at forest/woodland edges (May–Jul)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	- - -	- NP	Generally wooded areas (Jun–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	- - -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	- - -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Apocynum cannabinum</i>	Indian Hemp	Apocyn-	- - A2	- NP	Moist places, near streams, springs, or as weed in orchards (Apr–Oct)	

Herbaceous	Grouped by Growth Form				Herbaceous
<i>Aquilegia</i>	Sorted Alphabetically by Scientific Name				<i>Aquilegia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	- - C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	- - -	- NP	Common. Open to shady areas, often in drainages (May–Nov)
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	- - B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	- - -	- NP	Dry ground, valleys, foothills (May–Oct)
<input type="radio"/> <i>Aster sp.</i>	Aster	Aster-	- - -	- NP	--
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	- - -	- NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)
<input type="radio"/> <i>Balsamorhiza deltoidea</i>	Balsamroot	Aster-	- - A2	- NP	Common. Grassy slopes, open forests, shrubby areas (Apr–Jul)
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	- - -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	- - -	L ZA	Disturbed grassland. (Apr–Jun)
<input type="radio"/> <i>Blennosperma nanum var. nanum</i>	Glue-seed	Aster-	- - C	- NA	Open, grassy areas, often margins of seeps or vernal pools (Jan–May)
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	- - -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	- - -	M ZA	Common. Disturbed areas, fields (Apr–Sep)
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Themid-	- - -	- NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	- - -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)
<input type="radio"/> <i>Calochortus pulchellus</i>	Mount Diablo Fairy-lantern	Lili-	- 1B.2 *A2	- NP	Wooded slopes, rarely chaparral, gen n aspect (Apr–Jun)
<input type="radio"/> <i>Calochortus sp.</i>	Mariposa Lily	Lili-	- - -	- NP	--
<input type="radio"/> <i>Calochortus splendens</i>	Splendid Lilac Mariposa Lily	Lili-	- - A1	- NP	Dry, often granitic soils, grassland, chaparral, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	- - -	- NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	- - B	- NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calycadenia truncata</i>	Rosinweed	Aster-	- - A2	- NA	Common. Dry, open hillsides, rocky ridges, talus, grassland, openings in foothill woodland, scrub (May–Oct)
<input type="radio"/> <i>Calystegia sp.</i>	Morning-glory	Convolvul-	- - -	- NP	--
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	- - -	- NP	Dry, open scrub or woodland (Apr–Jun)

Herbaceous <i>Capsella</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Capsella</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	- - -	- ZA	Disturbed areas (Jan–Oct)
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	- - -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	- - -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	- - -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	- - -	- NP	Chaparral, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Orobanch-	- - -	- NA	Grassland (Mar–May)
<input type="radio"/> <i>Castilleja exserta subsp. exserta</i>	Purple Owl's-clover	Orobanch-	- - -	- NA	Open fields, grassland (Mar–May)
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	- - -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i>)	California Mustard	Brassic-	- - -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	- - -	M ZA	Disturbed fields, open woodland (Apr–Jul)
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	- - -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	- - -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	- - -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Aster-	- - -	- NB	Disturbed areas, grassland, woodland (May–Jul)
<input type="radio"/> <i>Cirsium</i> sp.	Thistle	Aster-	- - -	- --	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	- - -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	- - -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	- - -	- NA	--
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	- - -	- NA	Common. Woodland (Apr–Sep)
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>exigua</i>	Common Pale Claytonia	Monti-	- - C	- NA	Dry or moist, disturbed bare clay to sandy soils, often serpentine (Apr–Jul)
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>glauca</i>	Glaucus Pale Claytonia	Monti-	- - A1	- NA	
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i>)	Small-flower Claytonia	Monti-	- - -	- NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	- - -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)

Herbaceous <i>Claytonia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Claytonia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	- - -	- NA	Vernally moist, often shady or disturbed sites (Jan–May)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i>)	Yerba Buena	Lami-	- - -	- NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	- - -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)
<input type="radio"/> <i>Collinsia parviflora</i>	Blue-eyed Mary	Plantagin-	- - A2	- NA	Common. Moist, ± shady places, montane (Mar–Jul)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	- - C	- NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>sparsiflora</i>	Few-flowered Collinsia	Plantagin-	- - C	- NA	Grassy, sometimes disturbed or rocky places, chaparral, oak woodland, dry mixed woodland (Mar–May)
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	- - -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	- - -	- NA	Open areas (Feb–May)
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i>)	Turkey-mullein	Euphorbi-	- - -	- NA	Dry, open, often disturbed areas (May–Oct)
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	- - -	M ZP	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	- - -	- NP	Chaparral, woodland (Feb–May)
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>hesperium</i>	Western Larkspur	Ranuncul-	- - B	- NP	Oak woodland, w slope coast ranges (Mar–Jun)
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>palescens</i>	Pale-flowered Western Larkspur	Ranuncul-	- - C	- NP	Oak woodland, e slope coast ranges (Mar–May)
<input type="radio"/> <i>Delphinium variegatum</i> subsp. <i>variegatum</i>	Royal Larkspur	Ranuncul-	- - C	- NP	Grassland, open oak woodland (Mar–May)
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	- - -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	- - -	- NP	Open woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	- - -	- NP	Gen in shady sites (Mar–Jul)
<input type="radio"/> <i>Downingia</i> sp.	Downingia	Campanul-	- - -	- NA	--
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i>)	Sticky Cinquefoil	Ros-	- - -	- NP	Gen ± shady or moist areas (May–Jul)
<input type="radio"/> <i>Emmenanthe penduliflora</i> var. <i>penduliflora</i>	Whispering Bells	Boragin-	- - B	- NP	Chaparral to creosote-bush scrub, rocky, sandy, decomposed granite, serpentine soils (Apr–Jul)
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	- - -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	- - -	- NP	Dry slopes, ridges (Jun–Dec)
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Onagr-	- - -	- NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)

Herbaceous <i>Epilobium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Epilobium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Epilobium minutum</i>	Chaparral Willowherb	Onagr-	- - -	- NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)
<input type="radio"/> <i>Epilobium torreyi</i>	Narrowleaf Willowherb	Onagr-	- - A2	- NA	Seasonally moist streambanks, seeps, roadside ditches (May–Aug)
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Aster-	- - -	- NP	Grassy dunes, chaparral, oak woodland (May–Oct)
<input type="radio"/> <i>Erigeron petrophilus</i> var. <i>petrophilus</i>	Rock-loving Fleabane	Aster-	- - A2	- NP	Rocky foothills to montane forest, sometimes on serpentine (May–Sep)
<input type="radio"/> <i>Erigeron philadelphicus</i> var. <i>philadelphicus</i> (JM93: no var.)	Philadelphia Fleabane	Aster-	- - C	- NBP	Streamsides, other moist habitats (May–Jun)
<input type="radio"/> <i>Erigeron</i> sp. (JM93: <i>Conyza</i>)	Fleabane Daisy	Aster-	- - -	- --	
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	- - A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Eriogonum</i> sp.	Wild Buckwheat	Polygon-	- - -	- NAP	--
<input type="radio"/> <i>Eriophyllum lanatum</i> var. <i>achilleoides</i> (JM93: var. <i>achillaeoides</i>)	Common Woolly Sunflower	Aster-	- - C	- NP	Dry, often rocky sites, chaparral, forest (May–Jul)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	- - -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	- - -	- ZA	Open, disturbed sites (Feb–Sep)
<input type="radio"/> <i>Eryngium jepsonii</i> (JM93: <i>E. aristulatum</i> var. <i>aristulatum</i>)	Button Celery	Api-	- - -	- NP	Moist clay soil, vernal pools, lake shores, drying lakes, wet depressions (Apr–Aug)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	- - -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	- - -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i>)	Broadleaf Aster	Aster-	- - -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i>)	Checker Lily	Lili-	- - -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Galium andrewsii</i> subsp. <i>gatense</i>	Phlox-leaf Serpentine Bedstraw	Rubi-	- 4.2 *A2	- NP	Dry, rocky places in serpentine soil, chaparral or open oak/pine woodland (Apr–Jun)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	- - -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Rubi-	- - -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	- - -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	- - -	M ZA	Open, disturbed sites (Mar–Jul)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	- - -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>multicaulis</i>	Many-stem California Gilia	Polemoni-	- - -	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Feb–Jun)
<input type="radio"/> <i>Gilia clivorum</i>	Purplespot Gilia	Polemoni-	- - C	- NA	Common. Open, grassy areas (Feb–Jun)
<input type="radio"/> <i>Gilia tricolor</i> subsp. ?	Bird's-eye Gilia	Polemoni-	- - -	- NA	--

Herbaceous <i>Githopsis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Githopsis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Githopsis specularioides</i>	Looking-glass Bluecup	Campanul-	- - B	- NA	Chaparral, oak woodland (Apr–May)
<input type="radio"/> <i>Gnaphalium</i> or <i>Pseudognaphalium</i> sp. (JM93: <i>Gnaphalium</i> sp.)	Everlasting Cudweed	Aster-	- - -	- - -	
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i>)	Great Valley Gumplant	Aster-	- - -	- NP	Sandy or saline bottomland, roadsides (May–Nov)
<input type="radio"/> <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	- 1B.2 *A2	- NP	Open, grassy sites (Apr–Jun)
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i>)	Bristly Ox-tongue	Aster-	- - -	L ZAB	Common. Disturbed areas (All year)
<input type="radio"/> <i>Hemizonia</i> , <i>Centromadia</i> , or <i>Deinandra</i> sp. (JM93: <i>Hemizonia</i> sp.)	Spikeweed	Aster-	- - -	- NAP	- -
<input type="radio"/> <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Aster-	- - -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)
<input type="radio"/> <i>Hesperolinon breweri</i>	Brewer's Western Flax	Lin-	- 1B.2 *A2	- NA	Chaparral or grassland, occ on serpentine (May–Jun)
<input type="radio"/> <i>Heterotheca sessiliflora</i> subsp. ?	Sessileflower Goldenaster	Aster-	- - -	- NP	- -
<input type="radio"/> <i>Holocarpha heermannii</i>	Heermann Tarplant	Aster-	- - -	- NA	Grassland (May–Nov)
<input type="radio"/> <i>Holocarpha obconica</i>	San Joaquin Tarplant	Aster-	- - B	- NA	Grassland (Apr–Nov)
<input type="radio"/> <i>Horkelia californica</i> var. <i>frondosa</i> (JM93: subsp. <i>frondosa</i>)	Unlobed California Horkelia	Ros-	- - C	- NP	Coastal scrub, canyons, poison-oak thickets (May–Oct)
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	- - -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	- - -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	- - -	- ZA	Roadsides, grassland (Jul–Nov)
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	- - -	- ZA	Abundant. Disturbed places (May–Oct)
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: subsp. <i>ramosissima</i>)	Common Hare-leaf	Aster-	- - -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lasthenia californica</i> subsp. <i>californica</i> (JM93: no subsp.)	California Goldfields	Aster-	- - -	- NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lasthenia microglossa</i>	Small-ray Goldfields	Aster-	- - B	- NA	Shaded slopes of woodland, chaparral, desert scrub (Mar–May)
<input type="radio"/> <i>Lathyrus jepsonii</i> var. <i>californicus</i>	Bluff Pea	Fab-	- - B	- NP	Forest, open areas (Apr–Aug)
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	- - -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Layia gaillardiioides</i>	Yellow Woodland Tidytops	Aster-	- - A2	- NA	Open or semi-shaded slopes, in sandy or clayey soil (including serpentine) (Mar–Aug)

Herbaceous <i>Lepidium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lepidium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i>)	Threadleaf Peppergrass	Brassic-	- - -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i>)	Pinklobe Leptosiphon	Polemoni-	- - C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)	
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i>)	Bicolor Leptosiphon	Polemoni-	- - -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Leptosiphon ciliatus</i> (JM93: <i>Linanthus</i>)	Whisker Brush	Polemoni-	- - B	- NA	Common. Open or wooded areas (Mar–Jul)	
<input type="radio"/> <i>Linanthus</i> or <i>Leptosiphon</i> sp. (JM93: <i>Linanthus</i> sp.)	Linanthus	Polemoni-	- - -	- NA	--	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	- - -	- NP	Open, grassy slopes (Mar–Apr)	
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	- - -	- NP	Shaded slopes (Feb–Jun)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Aster-	- - -	- ZA	Bare or grassy openings, burns (Mar–Jul)	
<input type="radio"/> <i>Lomatium caruifolium</i> var. <i>caruifolium</i>	Alkali Desertparsley	Api-	- - B	- NP	Wet, clay depressions, open grassland (Mar–May)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	- - -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium</i> sp.	Biscuit Root	Api-	- - -	- NP	--	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	- - -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus</i> sp.	Lupine	Fab-	- - -	- NAP	--	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	- - -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	- - -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia</i> or <i>Anisocarpus</i> sp. (JM93: <i>Madia</i> sp.)	Tarweed	Aster-	- - -	- NA	--	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i>)	Fat False Solomon's Seal	Rusc-	- - -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i>)	Starry False Solomon's Seal	Rusc-	- - -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i>)	California Man-root	Cucurbit-	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	

Herbaceous <i>Matricaria</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Matricaria</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i>)	Pineapple Weed	Aster-	- - -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Fab-	- - -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i>)	California Saxifrage	Saxifrag-	- - -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Aster-	- - -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	- - -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris</i> sp.	Silverpuffs	Aster-	- - -	- NA	- -	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i>)	Slender Annual Phlox	Polemoni-	- - -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	- - -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Minuartia douglasii</i>	Douglas' Stitchwort	Caryophyll-	- - B	- NA	Dry, rocky, slopes, flats in chaparral, oak and pine woodland, often serpentine (Spring–early summer)	
<input type="radio"/> <i>Minuartia pusilla</i>	Annual Sandwort	Caryophyll-	- - A2	- NA	Plains, open pine forest, chaparral slopes (Spring–summer)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i>)	Coyote-mint	Lami-	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Montia fontana</i>	Water Chickweed	Monti-	- - C	- NA	Common. Ponds, streams, vernal pools, seeps (Jul–Aug)	
<input type="radio"/> <i>Muilla maritima</i>	Common Muilla	Themid-	- - B	- NP	Coastal scrub, grassland, woodland, valleys (Mar–Jun)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i>)	Water Cress	Brassic-	- - -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia intertexta</i> subsp. <i>intertexta</i>	Needle-leaved Navarretia	Polemoni-	- - A1	- NA	Open, seasonally wet areas, meadows, vernal pools (May–Jul)	
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	- - C	- NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)	
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Polemoni-	- - C	- NA	Open, slopes, gravel, clay (May–Jul)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	- - -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Boragin-	- - -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Nemophila parviflora</i> var. <i>parviflora</i>	Small-flower Nemophila	Boragin-	- - C	- NA	Woodland, forest, roadsides, slopes (Mar–Jul)	

Herbaceous	Grouped by Growth Form					Herbaceous
Nemophila	Sorted Alphabetically by Scientific Name					Nemophila
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Nemophila pedunculata</i>	Meadow Nemophila	Boragin-	- - B	- NA	Common. Ocean bluffs, grassland, slopes, meadows, sandbars, fields, woodland, streambanks (Feb–Jul)	
<input type="radio"/> <i>Orobanche bulbosa</i>	Chaparral Broom-rape	Orobanch-	- - A2	- NP	Openings in chaparral, generally on <i>Adenostoma fasciculatum</i> (Apr–Jun)	
<input type="radio"/> <i>Orobanche fasciculata</i>	Clustered Broomrape	Orobanch-	- - B	- NP	Dry, gen bare places. Root parasite on various shrubs (Apr–Jul)	
<input type="radio"/> <i>Orobanche uniflora</i>	Naked Broom-rape	Orobanch-	- - C	- NP	Generally moist places, on herbs, especially <i>Sedum</i> , <i>Saxifragaceae</i> , <i>Asteraceae</i> (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Api-	- - -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza brachypoda</i>	California Sweet Cicely	Api-	- - B	- NP	Moist ravines, conifer forest, woodland (Mar–May)	
<input type="radio"/> <i>Packera breweri</i> (JM93: <i>Senecio</i>)	Brewer's Ragwort	Aster-	- - B	- NP	Common. Seasonally damp grassland, oak savanna, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i>)	Wind Poppy	Papaver-	- - -	- NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Pectocarya pusilla</i>	Little Pectocarya	Boragin-	- - B	- NA	Dry, semi-barren sites in grassland, chaparral, woodland, roadsides (Mar–Jun)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	- - -	- NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Pediomelum californicum</i>	Indian Breadroot	Fab-	- - A1	- NP	Open chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	- - -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Phacelia ciliata</i>	Field Phacelia	Boragin-	- - B	- NA	Clay or gravelly slopes in grassland, fields (Feb–Jun)	
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	- - -	- NA	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Phacelia ramosissima</i> (JM93: var. <i>ramosissima</i> , <i>latifolia</i>)	Branched Phacelia	Boragin-	- - A2	- NP	Diverse habitats incl salt marshes, canyons (Apr–Oct)	
<input type="radio"/> <i>Pholistoma membranaceum</i>	White Fiesta Flower	Boragin-	- - -	- NA	Beaches, bluffs, ravines, wooded slopes, desert washes (Feb–May)	
<input type="radio"/> <i>Phoradendron serotinum subsp. macrophyllum</i> (JM93: <i>P. macrophyllum</i>)	Big-leaf Mistletoe	Visc-	- - C	- NP	On trees other than <i>Quercus</i> (especially <i>Alnus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Platanus</i> , <i>Populus</i> , <i>Robinia</i> , <i>Salix</i>) (Dec–Mar)	
<input type="radio"/> <i>Phoradendron serotinum subsp. tomentosum</i> (JM93: <i>P. villosum</i>)	Oak Mistletoe	Visc-	- - -	- NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)	
<input type="radio"/> <i>Piperia elegans subsp. elegans</i> (JM93: no subsp.)	Elegant Rein Orchid	Orchid-	- - B	- NP	Generally dry, open sites, scrub, conifer forest (May–Sep)	
<input type="radio"/> <i>Piperia michaelii</i>	Michael's Rein Orchid	Orchid-	- 4.2 *A2	- NP		

Herbaceous <i>Piperia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Piperia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Piperia unalascensis</i>	Alaska Rein Orchid	Orchid-	- - A1	- NP	Generally dry sites, scrub, woodland, forest (May–Aug)	
<input type="radio"/> <i>Plagiobothrys canescens</i> var. <i>canescens</i> (JM93: no var.)	Valley Popcornflower	Boragin-	- - C	- NA	Common. Grassland, woodland, coastal scrub, desert scrub, roadsides (Mar–Jun)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	- - -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	- - -	- -		
<input type="radio"/> <i>Plagiobothrys stipitatus</i> var. <i>micranthus</i>	Valley Popcornflower	Boragin-	- - -	- NA	Common. Vernal pools, wet sites in grassland, conifer forest, sagebrush scrub (Apr–Jul)	
<input type="radio"/> <i>Plagiobothrys stipitatus</i> var. <i>stipitatus</i>	Showy Great Popcornflower	Boragin-	- - C	- NA	Vernal pools, wet sites in grassland (Mar–Jun)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	- - -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: subsp. <i>ciliosa</i> , <i>insignis</i>)	Longspur Plectritis	Valerian-	- - B	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis congesta</i> subsp. <i>brachystemon</i> (JM93: <i>P. brachystemon</i>)	Pale Plectritis	Valerian-	- - B	- NA	Common. Coastal bluffs, open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis congesta</i> subsp. <i>congesta</i>	Sea Blush	Valerian-	- - A2	- NA	Coastal bluffs, open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	- - -	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i>)	Hooker Fairy Bells	Lili-	- - C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)	
<input type="radio"/> <i>Prunella vulgaris</i> var. ?	European Selfheal	Lami-	- - -	- ZP	--	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i>)	California Everlasting	Aster-	- - -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i>)	Weedy Cudweed	Aster-	- - -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)	
<input type="radio"/> <i>Psilocarphus tenellus</i> (JM93: var. <i>tenellus</i>)	Slender Woolly-marbles	Aster-	- - -	- NA	Common. Dry, seasonally moist slopes, flats, burns, trails, rarely vernal pools (Mar–Jul)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	- - -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	- - B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)	
<input type="radio"/> <i>Ranunculus aquatilis</i> var. <i>diffusus</i> (JM93: var. <i>capillaceus</i>)	Thread-leaf Water Buttercup	Ranuncul-	- - -	- NP	Ponds, lakes, streams (Mar–Sep)	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	- - -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	- - -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Ranunculus</i>	Sorted Alphabetically by Scientific Name					<i>Ranunculus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Ranunculus occidentalis</i> var. <i>occidentalis</i> (JM93: no var.)	Western Buttercup	Ranuncul-	- - A2	- NP	Grassy slopes in meadows or open woodland (Mar–Jul)	
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	- - -	L ZA	Disturbed areas, fields (May–Jul)	
<input type="radio"/> <i>Rigiopappus leptocladus</i>	Bristle-head	Aster-	- - B	- NA	Grassy sites, openings among shrubs or trees (Apr–Jul)	
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	- - -	M ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	- - -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	- - -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i>)	Thin-leaf Willow Dock	Polygon-	- - B	- NP	Wet places, margins, rocky slopes (May–Jul)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	- - -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	- - -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	- - -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula tuberosa</i>	Turkey Pea Sanicle	Api-	- - C	- NP	Open gravelly meadows, chaparral, woodland, pine forest (Mar–Jul)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica</i> , <i>floribunda</i>)	California Figwort	Scrophulari-	- - -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Sedella pentandra</i> (JM93: <i>Parvisedum pentandrum</i>)	Annual Stonecrop	Crassul-	- - A2	- NA	Compacted ground, slate, shale, sandstone or serpentine outcrops (Mar–Jun)	
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	- - C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	- - -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	- - -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	- - -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silene laciniata</i> subsp. <i>californica</i> (JM93: <i>S. californica</i>)	California Pink	Caryophyll-	- - B	- NP	Chaparral, oak woodland, conifer forest, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sinapis arvensis</i>	Charlock	Brassic-	- - -	L ZA	Disturbed areas (Mar–Sep)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	- - -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	- - -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solidago</i> sp.	Goldenrod	Aster-	- - -	- NP	--	
<input type="radio"/> <i>Solidago velutina</i> subsp. <i>californica</i> (JM93: <i>S. californica</i>)	California Goldenrod	Aster-	- - -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	

Herbaceous Sonchus	Grouped by Growth Form				Herbaceous Sonchus
Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	- - -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	- - -	- ZA	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Spergula arvensis</i> (JM93: subsp. <i>arvensis</i>)	Stickwort	Caryophyll-	- - -	- ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	- - -	- ZAP	Forest, meadows, mud flats, disturbed
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Lami-	- - C	- NP	Streambanks, springs, pine/oak forest (Jun–Oct)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i>)	Common Rigid Hedge-nettle	Lami-	- - -	- NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	- - -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	- - C	- NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	- - C	- NP	Chaparral, pine/oak woodland (Mar–May)
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	- - -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	- - -	- NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)
<input type="radio"/> <i>Thysanocarpus radians</i>	Ribbed Fringepod	Brassic-	- - A2	- NA	Moist slopes, pastures, open meadows, fields (Mar–Apr)
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	- - -	M ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	- - -	- ZA	Disturbed places (Apr–Jun)
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i>)	Common Star Lily	Melanthi-	- - -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)
<input type="radio"/> <i>Trichostema lanceolatum</i>	Vinegar Weed	Lami-	- - -	- NA	Dry, open, gen disturbed habitats (Jun–Nov)
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i>)	Rancheria Clover	Fab-	- - C	- NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decepiens</i>	Deceiving Clover	Fab-	- - -	- NA	Open, grassy areas, forest (Apr–Jun)
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	- - -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Fab-	- - -	- NA	Grassland, coastal woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>depauperatum</i>	Dwarf Sack Clover	Fab-	- - B	- NA	Wet meadows, grassland, roadsides, open spring-moist, heavy soils (Mar–May)
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Fab-	- - -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium dichotomum</i> (JM93: <i>T. albopurpureum</i> var. <i>dichotomum</i>)	Branched Indian Clover	Fab-	- - B	- NA	Coastal dunes, open slopes, meadows, oak woodland, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	- - -	- ZA	Agricultural, disturbed areas, lawns (Spring)

Herbaceous <i>Trifolium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Trifolium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	- - -	M ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	- - -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	- - -	- NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Fab-	- - C	- NA	Woody or shrubby slopes, roadsides (Mar–Jun)	
<input type="radio"/> <i>Trifolium olivaceum</i> (JM93: <i>T. albopurpureum</i> var. <i>olivaceum</i>)	Olive Clover	Fab-	- - B	- NA	Grassy slopes, valley meadows, disturbed areas (Apr–May)	
<input type="radio"/> <i>Trifolium variegatum</i> var. ? (JM93: no var.)	White-tip Clover	Fab-	- - -	- NA	--	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	- - -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Triphysaria eriantha</i> subsp. <i>eriantha</i>	Yellow Johnny-tuck	Orobanch-	- - -	- NA	Grassland, foothills (Mar–May)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	- - -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	- - -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	- - -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Brassic-	- - B	- NP	Grassy banks, open fields, roadsides, pastures (Mar–May)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	- - -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urtica dioica</i> subsp. <i>holosericea</i>	Hoary Nettle	Urtic-	- - -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	- - -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbena lasiostachys</i> var. ?	Western Vervain	Verben-	- - -	- NP	--	
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	- - -	- ZA	Wet, disturbed areas, fields (Feb–May)	
<input type="radio"/> <i>Vicia americana</i> subsp. <i>americana</i> (JM93: var. <i>americana</i>)	American Vetch	Fab-	- - -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>nigra</i>	Narrow-leaved Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>sativa</i>	Spring Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	- - C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	

Herbaceous <i>Wyethia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Wyethia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	- - C	- NP	Gen shady sites (Mar–Jun)	
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	- - -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))	
<input type="radio"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Api-	- - C	- NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)	

Woody	Grouped by Growth Form					Woody
Acer	Sorted Alphabetically by Scientific Name					Acer
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	- - -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	- - -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	- - -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	- - C	- NP	Conifer, oak forests (Mar–May)	
<input type="radio"/> <i>Arctostaphylos auriculata</i>	Mount Diablo Manzanita	Eric-	- 1B.3 *A2	- NP	Sandstone, upland chaparral near coast. Mount Diablo and vicinity (Feb–Mar)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Eric-	- - C	- NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Arctostaphylos manzanita</i> subsp. <i>laevigata</i>	Contra Costa Manzanita	Eric-	- 1B.2 *A2	- NP	Chaparral, rocky outcrops (Feb–May)	
<input type="radio"/> <i>Arctostaphylos manzanita</i> subsp. <i>manzanita</i>	Common / Giant Manzanita	Eric-	- - C	- NP	Chaparral, conifer forest (Feb–May)	
<input type="radio"/> <i>Arctostaphylos</i> sp.	Manzanita	Eric-	- - -	- NP	--	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	- - -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	- - -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Berberis aquifolium</i> var. <i>dictyota</i>	Hollyleaf Oregon Grape	Berberid-	- - A2	- NP	Slopes, canyons, conifer forest, oak woodland, chaparral (Mar–May)	
<input type="radio"/> <i>Berberis pinnata</i> subsp. <i>pinnata</i>	Shinyleaf Oregon-grape	Berberid-	- - C	- NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
<input type="radio"/> <i>Berberis</i> sp.	Oregon Grape	Berberid-	- - -	- NP	--	
<input type="radio"/> <i>Ceanothus cuneatus</i> var. <i>cuneatus</i>	Buckbrush	Rhamn-	- - C	- NP	Sandy to rocky flats, slopes, ridges (Feb–May)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	- - -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
<input type="radio"/> <i>Eriodictyon californicum</i>	California Yerba Santa	Boragin-	- - -	- NP	Slopes, fields, roadsides, woodland, chaparral (Apr–Jul)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	- - -	- NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i>)	California Coffee Berry	Rhamn-	- - -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Fraxinus dipetala</i>	California Ash	Ole-	- - A2	- NP	Canyons, slopes, chaparral, oak/pine woodland (Apr–Jun)	
<input type="radio"/> <i>Fremontodendron californicum</i> (JM93: subsp. <i>californicum</i>)	Calif. Flannelbush/fremontia	Malv-	- - A1	- NP		
<input type="radio"/> <i>Garrya flavescens</i>	Ashy Silk Tassel	Garry-	- - A2	- NP		
<input type="radio"/> <i>Garrya fremontii</i>	Green-leaf Silk Tassel	Garry-	- - A1	- NP	Chaparral, foothill woodland, montane forest (Jan–Apr)	

Woody	Grouped by Growth Form					Woody
Gutierrezia	Sorted Alphabetically by Scientific Name					Gutierrezia
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	- - C	- NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Helianthemum scoparium</i>	Peak Rush-rose	Cist-	- - B	- NP	Dry sandy or rocky soil of hills, slopes, ridges (Feb–Sep)	
<input type="radio"/> <i>Hesperocyparis macrocarpa</i> (JM93: <i>Cupressus</i>)	Monterey Cypress	Cupress-	- - -	N NP	Closed-cone-pine/cypress forests -	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	- - -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	- - -	- NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i>)	N. California Black Walnut	Jugland-	- 1B.1 *A2	- NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Juniperus californica</i>	California Juniper	Cupress-	- - B	- NP	Dry slopes, flats, pinyon/juniper woodland -	
<input type="radio"/> <i>Lepechinia calycina</i>	Pitcher Sage	Lami-	- - B	- NP	Common. Rocky slopes; chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Caprifoli-	- - -	- NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Malacothamnus fremontii</i>	Fremont Bush-mallow	Malv-	- - B	- NP	Open chaparral to pine woodland (May–Jul)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	- - -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	- - -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Pickeringia montana</i> var. <i>montana</i>	Chaparral Pea	Fab-	- - B	- NP	Chaparral, open woodland (May–Aug)	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	- - B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Populus fremontii</i> subsp. <i>fremontii</i>	Fremont Cottonwood	Salic-	- - -	- NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)	
<input type="radio"/> <i>Prunus dulcis</i>	Almond	Ros-	- - -	- ZP	Canyons, roadsides, grassland (as waif) (Feb–Mar)	
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Ros-	- - C	- NP	Mixed-evergreen or conifer forest (Mar–May)	
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Ros-	- - -	- NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Ptelea crenulata</i>	Western Hop Tree	Rut-	- - B	- NP	Scrub, woodland (Apr–May)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	- - A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	- - C	- NP	Dry slopes, chaparral (Feb–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	- - -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	- - -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	

Woody	Grouped by Growth Form					Woody
Quercus	Sorted Alphabetically by Scientific Name					Quercus
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	- - -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	- - B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>frutescens</i>	Dwarf Interior Live Oak	Fag-	- - C	- NP	Valleys, chaparral (Mar–May)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	- - -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Rhamn-	- - C	- NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	- - C	- NP	Chaparral, montane forest (Mar–Jun)	
<input type="radio"/> <i>Rhus aromatica</i> (JM93: <i>R. trilobata</i>)	Skunk Bush	Anacardi-	- - C	- NP	Slopes, washes, scrub (Mar–May)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	- - -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	- - -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Fab-	- - -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	- - -	- NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	- - -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i>)	Himalayan Blackberry	Ros-	- - -	H ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	- - -	- NP	Open, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	- - -	- NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="radio"/> <i>Salix</i> sp.	Willow	Salic-	- - -	- ?P	--	
<input type="radio"/> <i>Salvia mellifera</i>	Black Sage	Lami-	- - -	- NP	Coastal-sage scrub, lower chaparral (Mar–Jun)	
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i>)	Blue Elderberry	Adox-	- - -	- NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	- - -	- NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="radio"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	- - -	- NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="radio"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	- - -	- NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	- - -	- NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="radio"/> <i>Umbellularia californica</i>	California Bay	Laur-	- - -	- NP	Common. Canyons, valleys, chaparral (Nov–May)	
<input type="radio"/> <i>Vitis californica</i>	California Wild Grape	Vit-	- - -	- NP	Streamsides, springs, canyons (May–Jun)	

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex dewayana ssp. *leptopoda* (*C. leptopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thrysiflorus (var. *thrysiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria amphibia*)

Polygonum amphibium var. *stipulaceum* (*Persicaria amphibia*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)