

Ohlone Regional Wilderness

Checklist of Wild Plants

Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Ohlone Regional Wilderness. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Family	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

Ohlone Regional Wilderness Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name					<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	- - -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Azolla filiculoides</i>	Mosquito Fern	Azoll-	- - -	- NP	Common. Ponds, slow streams -	
<input type="radio"/> <i>Cheilanthes covillei</i>	Coville's Lip Fern	Pterid-	- - A2	- NP	Crevices, bases of rocks, sun or shade -	
<input type="radio"/> <i>Cheilanthes intertexta</i>	Coastal Lip Fern	Pterid-	- - A2	- NP	Crevices, bases of rocks -	
<input type="radio"/> <i>Cystopteris fragilis</i>	Brittle Fern	Woods-	- - C	- NP	Shady, moist rock crevices, meadows, streamsides -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	- - -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	- - -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata var. mucronata</i>	Bird's-foot Fern	Pterid-	- - C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis subsp. triangularis</i>	Goldenback Fern	Pterid-	- - -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	- - -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Selaginella bigelovii</i>	Spike-moss	Selaginell-	- - A2	- NP	Open sites, mineral soil on rock outcrops, amid shrubs -	

Grasses & Grass-like <i>Aira</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Aira</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	- - -	- ZA	Sandy soils, open or disturbed sites (Apr–Jun)
<input type="radio"/> <i>Alopecurus saccatus</i>	Pacific Foxtail	Po-	- - B	- NA	Vernal pools, moist, open meadows (Mar–May)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	- - -	M ZA	Disturbed sites (Mar–Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	- - -	M ZA	Disturbed sites (Apr–Jun)
<input type="radio"/> <i>Bromus arenarius</i>	Australian Chess	Po-	- - -	- ZA	Open, disturbed places (Apr–Jul)
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	- - -	- NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr–Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	- - -	M ZA	Open, gen disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	- - -	L ZA	Fields, disturbed areas (Apr–Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	- - -	- NP	Shrubland, conifer forest, shaded streambanks, roadsides (May–Jul)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>madritensis</i>	Foxtail Chess	Po-	- - -	- ZA	Disturbed areas, roadsides (Apr–Jan)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	- - -	H ZA	Disturbed areas, roadsides (Mar–Jun)
<input type="radio"/> <i>Bromus tectorum</i>	Cheat Grass	Po-	- - -	H ZA	Open, disturbed areas (May–Aug)
<input type="radio"/> <i>Carex leporina</i> (JM93: <i>C. ovalis</i>)	Tracy Sedge	Cyper-	- - A1	- NP	Seasonally wet soil (May–Aug)
<input type="radio"/> <i>Carex serratodens</i>	Bifid Sedge	Cyper-	- - B	- NP	Moist places (Apr–Jul)
<input type="radio"/> <i>Carex</i> sp.	Sedge	Cyper-	- - -	- -	
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	- - -	M ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	- - -	L ZP	Disturbed, often moist sites (May–Aug)
<input type="radio"/> <i>Danthonia californica</i> (JM93: var. <i>californica</i>)	California Oat Grass	Po-	- - C	- NP	Gen moist meadows, open woodland (Apr–Aug)
<input type="radio"/> <i>Deschampsia cespitosa</i> subsp. <i>cespitosa</i>	Tufted Hair Grass	Po-	- - C	- NP	Meadows, streambanks, coastal marshes, forest (Jul–Aug)
<input type="radio"/> <i>Deschampsia danthonioides</i>	Annual Hair Grass	Po-	- - C	- NA	Moist to drying, open sites, meadows, streambanks, vernal pools, occ alkali soil (Mar–Aug)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	- - -	- NP	Common. Fresh to brackish wetland (Spring–summer)
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Po-	- - -	- NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	- - C	- NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i>)	Beardless Wild Rye	Po-	- - -	- NP	Dry to moist, often saline, meadows (Jun–Jul)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i>)	Brome Fescue	Po-	- - -	- ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	- - C	- NP	Dry, open forest, moist streambanks, chaparral (May–Jun)
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Po-	- - C	- NP	Dry, open or shady places (Jul–Sep)

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i>)	Hairy Fescue	Po-	- - -	- NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i>)	Rattail Sixweeks Grass	Po-	- - -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i>)	Rye Grass	Po-	- - -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	- - -	- NA	--
<input type="radio"/> <i>Festuca</i> sp. (perennial)	Perennial Fescue	Po-	- - -	-	--
<input type="radio"/> <i>Glyceria occidentalis</i>	Western Manna Grass	Po-	- - A2	- NP	Freshwater marshes, ponds and steams (Jun–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	- - -	- NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	- - -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	- - -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Junc-	- - -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	- - -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus tenuis</i>	Slender Rush	Junc-	- - -	- NP	Uncommon. Damp places (Jun–Sep)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	- - -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	- - C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Lamarckia aurea</i>	Goldentop	Po-	- - -	- ZA	Open ground, moist seeps, rocky hillsides, sandy soil (Feb–May)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	- - -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	- - C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	- - -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Paspalum distichum</i>	Knot Grass	Po-	- - -	- ZP	Edges of lakes, ponds, rice fields, wet roadside ditches (Jun–Oct)
<input type="radio"/> <i>Pleuropogon californicus</i> var. <i>californicus</i> (JM93: no var.)	California Semaphore Grass	Po-	- - B	- NA	Wet places, vernal pools, wet grassland (Mar–Jun)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	- - -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa bulbosa</i> subsp. <i>vivipara</i> (JM93: no subsp.)	Bulbous Blue Grass	Po-	- - -	- ZP	Disturbed places (Mar–Jul)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	- - -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	- - -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Po-	- - C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)

Grasses & Grass-like <i>Typha</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Typha</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	- - -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)

Herbaceous <i>Acaena</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Acaena</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acaena pinnatifida</i> var. <i>californica</i>	California Acaena	Ros-	- - C	- NP	Coastal grassland, open, rocky slopes (Mar–May)	
<input type="radio"/> <i>Acanthomintha lanceolata</i>	Santa Clara Thornmint	Lami-	- 4.2 *A2	- NP	Arid, rocky slopes, openings in scrub or woodland, gen on serpentine (Mar–Jun)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	- - -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Aster-	- - -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)	
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Fab-	- - -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Fab-	- - -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Agoseris apargioides</i> var. <i>apargioides</i>	Seaside Dandelion	Aster-	- - A1	- NP	Coastal dunes, sand hills (Apr–May)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	- - -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Allium bolanderi</i> var. <i>bolanderi</i>	Bolander's Onion	Alli-	- - A1	- NP	Uncommon. Rocky clays including serpentine (May–Aug)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	- - -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Alli-	- - C	- NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)	
<input type="radio"/> <i>Allophyllum divaricatum</i>	Stragglng Gilia	Polemoni-	- - A2	- NA	Sandy areas, chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Allophyllum gilioides</i> subsp. <i>gilioides</i>	Purple False Gilia	Polemoni-	- - B	- NA	Open, sandy, gen damp or grassy areas (Apr–Jun)	
<input type="radio"/> <i>Amaranthus californicus</i>	California Amaranth	Amaranth-	- - A2	- NA	Seasonally moist flats, lake margins, disturbed areas (Jul–Oct)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Boragin-	- - -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	- - -	- NP	Generally wooded areas (Jun–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	- - -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	- - -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	- - C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	- - -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	- - B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)	

Herbaceous <i>Asclepias</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Asclepias</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	- - -	- NP	Dry ground, valleys, foothills (May–Oct)	
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvech	Fab-	- - -	- NA	Open, grassy areas, scrub (Mar–Jul)	
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	- - -	- NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	- - -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	- - -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Themid-	- - -	- NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	- - -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Callitriche marginata</i>	California Water-starwort	Callitrich-	- - -	- NA	Becoming stranded (often in vernal pools) (Mar–Jun)	
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	- - -	- NP	Common. Shady to open woodland, scrub (Apr–Jun)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	- - C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	- - -	- NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	- - B	- NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calycadenia truncata</i>	Rosinweed	Aster-	- - A2	- NA	Common. Dry, open hillsides, rocky ridges, talus, grassland, openings in foothill woodland, scrub (May–Oct)	
<input type="radio"/> <i>Calystegia purpurata subsp. purpurata</i>	Climbing Morning-glory	Convolvul-	- - -	- NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia sp.</i>	Morning-glory	Convolvul-	- - -	- NP	--	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	- - -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Camissonia contorta</i>	Contorted Suncup	Onagr-	- - B	- NA	Sandy soil, slopes, flats, often disturbed, grassland, chaparral, pinyon/juniper woodland (May–Ju)	
<input type="radio"/> <i>Camissonia strigulosa</i>	Hairy Suncup	Onagr-	- - A1	- NA		
<input type="radio"/> <i>Camissoniopsis micrantha (JM93: Camissonia)</i>	Small-flower Suncup	Onagr-	- - B	- NA	Coastal dunes, beaches, sandy fields, washes (Mar–May)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	- - -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	- - -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus (JM93: no subsp.)</i>	Italian Thistle	Aster-	- - -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	

Herbaceous <i>Castilleja</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Castilleja</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Orobanch-	- - -	- NA	Grassland (Mar–May)
<input type="radio"/> <i>Castilleja exserta</i> subsp. <i>exserta</i>	Purple Owl's-clover	Orobanch-	- - -	- NA	Open fields, grassland (Mar–May)
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i>)	California Mustard	Brassic-	- - -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	- - -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	- - -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chaenactis glabriuscula</i> var. <i>megacephala</i>	Big-headed Yellow Pincushion	Aster-	- - A1	- NA	Dry, often sandy slopes, openings in chaparral, woodland (Mar–Jun)
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	- - -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Chorizanthe membranacea</i>	Pink Spineflower	Polygon-	- - A2	- NA	Common. Sand, gravel or rocks (Apr–Jul)
<input type="radio"/> <i>Chorizanthe polygonoides</i> var. <i>polygonoides</i>	Knotweed Spineflower	Polygon-	- - A1	- NA	Sand or gravel (Apr–Jun)
<input type="radio"/> <i>Cicuta douglasii</i>	Western Water-hemlock	Api-	- - B	- NP	Wet places, gen aquatic (Jun–Sep)
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Aster-	- - -	- NB	Disturbed areas, grassland, woodland (May–Jul)
<input type="radio"/> <i>Cirsium</i> sp.	Thistle	Aster-	- - -	- -	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	- - -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>automixa</i>	Santa Clara Red Ribbons	Onagr-	- 4.3 *A1	- NA	Woodland (Apr–Jun)
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>concinna</i>	Red Ribbons	Onagr-	- - C	- NA	Mixed-evergreen forest, woodland, coastal scrub (Apr–Jul)
<input type="radio"/> <i>Clarkia gracilis</i> subsp. <i>gracilis</i>	Herald-of-summer	Onagr-	- - B	- NA	Common. Openings in woodland, forest (Apr–Jul)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	- - -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	- - -	- NA	--
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	- - -	- NA	Common. Woodland (Apr–Sep)
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>exigua</i>	Common Pale Claytonia	Monti-	- - C	- NA	Dry or moist, disturbed bare clay to sandy soils, often serpentine (Apr–Jul)
<input type="radio"/> <i>Claytonia gypsophiloides</i>	Santa Lucia Claytonia	Monti-	- - A2	- NA	Moist, bare, often stony sites, in sun or shade, often serpentine (Mar–May)
<input type="radio"/> <i>Claytonia parviflora</i> subsp. ?	Miner's Lettuce	Monti-	- - -	- NA	--
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i>)	Small-flower Claytonia	Monti-	- - -	- NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i>)	Yerba Buena	Lami-	- - -	- NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. <i>bartsiiifolia</i>	White Chinese Houses	Plantagin-	- - A1	- NA	Open, sandy places (Mar–Jun)

Herbaceous <i>Collinsia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Collinsia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	- - -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)
<input type="radio"/> <i>Collinsia parviflora</i>	Blue-eyed Mary	Plantagin-	- - A2	- NA	Common. Moist, ± shady places, montane (Mar–Jul)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	- - C	- NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	- - -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)
<input type="radio"/> <i>Crassula aquatica</i>	Water Pygmy-weed	Crassul-	- - C	- NA	Salt marshes, vernal pools, margins of lakes, ponds (Mar–Jun(Aug))
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	- - -	- NA	Open areas (Feb–May)
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i>)	Turkey-mullein	Euphorbi-	- - -	- NA	Dry, open, often disturbed areas (May–Oct)
<input type="radio"/> <i>Cryptantha flaccida</i>	Weak-stemmed Cryptantha	Boragin-	- - -	- NA	Common; semi-barren, gravelly, loose soils, rocky sites, washes, slopes, ridges (Apr–Jul)
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	- - -	- NP	Chaparral, woodland (Feb–May)
<input type="radio"/> <i>Datisca glomerata</i>	Durango Root	Datisc-	- - A2	- NP	Dry streambeds or washes (May–Jul)
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	- - -	- NA	Rocky or sandy places (Apr–Jun)
<input type="radio"/> <i>Delphinium californicum</i> subsp. <i>californicum</i>	California Larkspur	Ranuncul-	- - A2	- NP	Generally slopes in dense chaparral, w side of coast ranges (Apr–Jun)
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Ranuncul-	- - C	- NP	Open coastal grassland, chaparral (Mar–May)
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>pallescens</i>	Pale-flowered Western Larkspur	Ranuncul-	- - C	- NP	Oak woodland, e slope coast ranges (Mar–May)
<input type="radio"/> <i>Delphinium nudicaule</i>	Red Or Orange Larkspur	Ranuncul-	- - C	- NP	Moist talus, wooded, rocky slopes (Mar–Jun)
<input type="radio"/> <i>Delphinium patens</i> subsp. <i>patens</i>	Spreading Larkspur	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Delphinium</i> sp.	Larkspur	Ranuncul-	- - -	- NP	--
<input type="radio"/> <i>Delphinium variegatum</i> subsp. <i>variegatum</i>	Royal Larkspur	Ranuncul-	- - C	- NP	Grassland, open oak woodland (Mar–May)
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	- - -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	- - -	- NP	Open woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	- - -	- NP	Gen in shady sites (Mar–Jul)
<input type="radio"/> <i>Downingia</i> sp.	Downingia	Campanul-	- - -	- NA	--
<input type="radio"/> <i>Draba verna</i>	Spring Whitlow Grass	Brassic-	- - B	- NA	Open or disturbed areas (Feb–May)
<input type="radio"/> <i>Dudleya cymosa</i> subsp. <i>paniculata</i>	Common / Hot Rock Dudleya	Crassul-	- - B	- NP	Uncommon. Rocky outcrops, canyons (May–Jun)
<input type="radio"/> <i>Elatine californica</i>	California Waterwort	Elatin-	- - A1	- NA	Pools, ponds, rice fields, streambanks (Mar–Aug)
<input type="radio"/> <i>Enemion occidentale</i> (JM93: <i>Isopyrum</i>)	Western Rue-anemone	Ranuncul-	- - A1	- NP	Shaded slopes, chaparral, oak woodland, conifer forest (Mar–May)

Herbaceous <i>Enemion</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Enemion</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Enemion stipitatum</i> (JM93: <i>Isopyrum</i>)	Siskiyou Rue-anemone	Ranuncul-	- - A1	- NP	Shaded slopes, chaparral, oak woodland, mixed-evergreen forest (Feb–Apr)
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	- - -	- NP	Dry slopes, ridges (Jun–Dec)
<input type="radio"/> <i>Epilobium torreyi</i>	Narrowleaf Willowherb	Onagr-	- - A2	- NA	Seasonally moist streambanks, seeps, roadside ditches (May–Aug)
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Aster-	- - -	- NP	Grassy dunes, chaparral, oak woodland (May–Oct)
<input type="radio"/> <i>Eriogonum gracile</i> var. <i>gracile</i>	Slender Woolly Wild Buckwheat	Polygon-	- - B	- NA	Common. Sand (All year)
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>luteolum</i>	Golden-carpet Wild Buckwheat	Polygon-	- - A2	- NA	Common. Serpentine (Jul–Nov)
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	- - -	- NP	--
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	- - A1	- NP	Common. Sand or gravel (May–Oct)
<input type="radio"/> <i>Eriogonum umbellatum</i> var. <i>smallianum</i> (JM93: var. <i>bahiiiforme</i>)	Small's Sulphur Flower	Polygon-	- - *A2	- NP	Serpentine (Jul–Sep)
<input type="radio"/> <i>Eriogonum wrightii</i> var. <i>trachyonum</i>	Rough-node Bastard-sage	Polygon-	- - A2	- NP	Common. Gravel (Jul–Oct)
<input type="radio"/> <i>Eriophyllum lanatum</i> var. <i>achilleoides</i> (JM93: var. <i>achillaeoides</i>)	Common Woolly Sunflower	Aster-	- - C	- NP	Dry, often rocky sites, chaparral, forest (May–Jul)
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	- - -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)
<input type="radio"/> <i>Erysimum capitatum</i> var. <i>capitatum</i> (JM93: var. <i>capitatum</i> , var. <i>angustatum</i>)	Western Wallflower	Brassic-	- - C	- NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	- - -	- NP	Grassy, open areas (Feb–Sep)
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Euphorbi-	- - -	- NA	Open, gen disturbed places (Mar–Jun)
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i>)	Broadleaf Aster	Aster-	- - -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	- - -	H ZP	Roadsides, disturbed sites (May–Sep)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i>)	Checker Lily	Lili-	- - -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Galium andrewsii</i> subsp. <i>gatense</i>	Phlox-leaf Serpentine Bedstraw	Rubi-	- 4.2 *A2	- NP	Dry, rocky places in serpentine soil, chaparral or open oak/pine woodland (Apr–Jun)
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	- - -	- NA	Grassy, ± shady places (Apr–Jun)
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	- - -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)
<input type="radio"/> <i>Galium parisiense</i>	Wall Bedstraw	Rubi-	- - -	- ZA	Warm, dry, generally rocky soil to moist areas, coastal-sage scrub, chaparral, grassy hillsides with oaks, roadsides (Apr–Aug)
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	- - -	- NP	Among shrubs in chaparral, forest (May–Aug)
<input type="radio"/> <i>Galium</i> sp.	Bedstraw	Rubi-	- - -	- --	
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	- - -	M ZA	Open, disturbed sites (Mar–Jul)

Herbaceous <i>Geranium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Geranium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	- - -	-	ZA	Open to shaded sites, disturbed ground (Feb–Aug)
<input type="radio"/> <i>Gilia achilleifolia subsp. achilleifolia</i>	California Gilia	Polemoni-	- - B	-	NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)
<input type="radio"/> <i>Gilia achilleifolia subsp. multicaulis</i>	Many-stem California Gilia	Polemoni-	- - -	-	NA	Open or shaded, gen grassy places, sandy or rocky soil (Feb–Jun)
<input type="radio"/> <i>Gilia tricolor subsp. diffusa</i>	Diffuse Bird's-eye Gilia	Polemoni-	- - B	-	NA	Open grassland, hills, valleys (Mar–Jul)
<input type="radio"/> <i>Gilia tricolor subsp. tricolor</i>	Bird's-eye Gilia	Polemoni-	- - B	-	NA	Open, grassland, hills, valleys (Mar–May)
<input type="radio"/> <i>Githopsis diffusa subsp. robusta</i>	Southern Bluecup	Campanul-	- - A2	-	NA	Shaded or disturbed areas, burns (Apr–Jun)
<input type="radio"/> <i>Githopsis specularioides</i>	Looking-glass Bluecup	Campanul-	- - B	-	NA	Chaparral, oak woodland (Apr–May)
<input type="radio"/> <i>Hemizonia, Centromadia, or Deinandra sp.</i> (JM93: <i>Hemizonia sp.</i>)	Spikeweed	Aster-	- - -	-	NAP	- -
<input type="radio"/> <i>Hesperolinon sp.</i>	Flax	Lin-	- - -	-	NA	
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	- - -	M	ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Holocarpha sp.</i>	Tarplant	Aster-	- - -	-	NA	- -
<input type="radio"/> <i>Holocarpha virgata subsp. virgata</i>	Sticky Tarplant	Aster-	- - -	-	NA	Grassland (May–Nov)
<input type="radio"/> <i>Horkelia californica var. elata</i> (JM93: <i>subsp. dissita</i>)	Many-lobed Calif. Horkelia	Ros-	- - A2	-	NP	Shady meadow edges, seasonal streams, open chaparral (Jun–Sep)
<input type="radio"/> <i>Hydrophyllum occidentale</i>	Western Waterleaf	Boragin-	- - A2	-	NP	Moist, shaded slopes, woodland, meadows, streambanks, chaparral (May–Jul)
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	- - -	L	ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	- - -	M	ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Idahoia scapigera</i>	Flat-pod	Brassic-	- - A1	-	NA	Moist ledges, slopes, meadows, foothills (Mar–May)
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	- - -	-	ZA	Roadsides, grassland (Jul–Nov)
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: <i>subsp. ramosissima</i>)	Common Hare-leaf	Aster-	- - -	-	NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lasthenia californica subsp. californica</i> (JM93: <i>no subsp.</i>)	California Goldfields	Aster-	- - -	-	NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lathyrus sp.</i>	Pea	Fab-	- - -	-	- -	
<input type="radio"/> <i>Lathyrus vestitus var. vestitus</i>	Pale Purple Pacific Pea	Fab-	- - -	-	NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Layia gaillardiioides</i>	Yellow Woodland Tidytops	Aster-	- - A2	-	NA	Open or semi-shaded slopes, in sandy or clayey soil (including serpentine) (Mar–Aug)
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	- - C	-	NA	Many habitats (Feb–Jul)

Herbaceous Legenere	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous Legenere
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Legenere limosa</i>	Legenere	Campanul-	- 1B.1 *A1	- NA	Wet areas, vernal pools, ponds (May–Jun)	
<input type="radio"/> <i>Lemna minor</i>	Common Lesser Duckweed	Lemn-	- - -	- NP	Common. Freshwater (Aug)	
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i>)	Threadleaf Peppergrass	Brassic-	- - -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
<input type="radio"/> <i>Lepidium oxycarpum</i>	Sharp-pod Peppergrass	Brassic-	- - B	- NA	Fields, vernal pool margins, alkaline flats (Mar–May)	
<input type="radio"/> <i>Leptosiphon ambiguus</i> (JM93: <i>Linanthus</i>)	Serpentine Leptosiphon	Polemoni-	- 4.2 *A2	- NA	Grassy areas gen serpentine (Apr–May)	
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i>)	Pinklobe Leptosiphon	Polemoni-	- - C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)	
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i>)	Bicolor Leptosiphon	Polemoni-	- - -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Leptosiphon ciliatus</i> (JM93: <i>Linanthus</i>)	Whisker Brush	Polemoni-	- - B	- NA	Common. Open or wooded areas (Mar–Jul)	
<input type="radio"/> <i>Lessingia nemaclada</i>	Funnel-flower Lessingia	Aster-	- - A1	- NA	Open fields, roadsides, woodland borders, occasionally on serpentine soil (Jul–Oct)	
<input type="radio"/> <i>Lewisia rediviva</i> var. <i>rediviva</i> (JM93: no var.)	Bitter Root	Monti-	- - A2	- NP	Rocky, sandy ground, open conifer woodland, scrub (Mar–Jun)	
<input type="radio"/> <i>Limosella acaulis</i>	Southern Mudwort	Scrophulari-	- - A2	- NA	Wet, muddy places, generally fresh water (May–Oct)	
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. ? (JM93: no subsp.)	Evening Snow	Polemoni-	- - A2	- NA	--	
<input type="radio"/> <i>Linanthus</i> or <i>Leptosiphon</i> sp. (JM93: <i>Linanthus</i> sp.)	Linanthus	Polemoni-	- - -	- NA	--	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	- - -	- NP	Open, grassy slopes (Mar–Apr)	
<input type="radio"/> <i>Lithophragma parviflorum</i> var. <i>parviflorum</i>	Prairie Starflower	Saxifrag-	- - B	- NP	Common. Open areas (Mar–Jul)	
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Aster-	- - -	- ZA	Bare or grassy openings, burns (Mar–Jul)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Api-	- - -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Api-	- - -	- NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium nudicaule</i>	Swollen-stem Biscuit Root	Api-	- - -	- NP	Rocky slopes, flats, gen pine woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium</i> sp.	Biscuit Root	Api-	- - -	- NP	--	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	- - -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Fab-	- - -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Lupinus</i>	Sorted Alphabetically by Scientific Name					<i>Lupinus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)	
<input type="radio"/> <i>Lupinus</i> sp.	Lupine	Fab-	- - -	- NAP	-	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	- - -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Madia exigua</i>	Thread-stem Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites, in sandy or clayey soils, including serpentine (Apr–Jul)	
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	- - -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i>)	California Man-root	Cucurbit-	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i>)	Pineapple Weed	Aster-	- - -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Fab-	- - -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Mentzelia affinis</i>	Hydra Blazing Star	Loas-	- - A2	- NA	Rocky or gray-white soils in grassland, woodland, creosote-bush scrub (Apr–May)	
<input type="radio"/> <i>Mentzelia lindleyi</i>	Lindley Blazing Star	Loas-	- - A2	- NA	Rocky, open slopes, coastal-sage scrub, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i>)	California Saxifrage	Saxifrag-	- - -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Aster-	- - -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	- - -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris</i> sp.	Silverpuffs	Aster-	- - -	- NA	-	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i>)	Slender Annual Phlox	Polemoni-	- - -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus douglasii</i>	Mickey Mouse Monkeyflower	Phrym-	- - A2	- NA	Bare clay, serpentine or granitic soils; generally along upper banks of small creeks (Feb–Apr)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	- - -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	

Herbaceous <i>Minuartia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Minuartia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Minuartia californica</i>	California Sandwort	Caryophyll-	- - A2	- NA	Gravelly, sandy slopes, grassy ridges, chaparral, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Minuartia douglasii</i>	Douglas' Stitchwort	Caryophyll-	- - B	- NA	Dry, rocky, slopes, flats in chaparral, oak and pine woodland, often serpentine (Spring–early summer)	
<input type="radio"/> <i>Minuartia pusilla</i>	Annual Sandwort	Caryophyll-	- - A2	- NA	Plains, open pine forest, chaparral slopes (Spring–summer)	
<input type="radio"/> <i>Moehringia macrophylla</i>	Big-leaf Sandwort	Caryophyll-	- - A2	- NP	Moist, shaded slopes, rocky ridges, summits, pine, oak forests, serpentine (Spring–summer)	
<input type="radio"/> <i>Monardella villosa subsp. villosa</i> (JM93: <i>M. villosa subsp. villosa, globosa, M. antonina subsp. antonina</i>)	Coyote-mint	Lami-	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Montia fontana</i>	Water Chickweed	Monti-	- - C	- NA	Common. Ponds, streams, vernal pools, seeps (Jul–Aug)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i>)	Water Cress	Brassic-	- - -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia intertextata subsp. intertextata</i>	Needle-leaved Navarretia	Polemoni-	- - A1	- NA	Open, seasonally wet areas, meadows, vernal pools (May–Jul)	
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Polemoni-	- - C	- NA	Open, slopes, gravel, clay (May–Jul)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	- - -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii var. menziesii</i>	Baby Blue-eyes	Boragin-	- - -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Nemophila parviflora var. parviflora</i>	Small-flower Nemophila	Boragin-	- - C	- NA	Woodland, forest, roadsides, slopes (Mar–Jul)	
<input type="radio"/> <i>Nemophila pedunculata</i>	Meadow Nemophila	Boragin-	- - B	- NA	Common. Ocean bluffs, grassland, slopes, meadows, sandbars, fields, woodland, streambanks (Feb–Jul)	
<input type="radio"/> <i>Orobanche fasciculata</i>	Clustered Broomrape	Orobanch-	- - B	- NP	Dry, gen bare places. Root parasite on various shrubs (Apr–Jul)	
<input type="radio"/> <i>Orobanche uniflora</i>	Naked Broom-rape	Orobanch-	- - C	- NP	Generally moist places, on herbs, especially Sedum, Saxifragaceae, Asteraceae (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Api-	- - -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza brachypoda</i>	California Sweet Cicely	Api-	- - B	- NP	Moist ravines, conifer forest, woodland (Mar–May)	
<input type="radio"/> <i>Packera breweri</i> (JM93: <i>Senecio</i>)	Brewer's Ragwort	Aster-	- - B	- NP	Common. Seasonally damp grassland, oak savanna, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i>)	Wind Poppy	Papaver-	- - -	- NA	Grassy areas, openings in chaparral (Apr–May)	

Herbaceous <i>Pectocarya</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Pectocarya</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Pectocarya pusilla</i>	Little Pectocarya	Boragin-	- - B	- NA	Dry, semi-barren sites in grassland, chaparral, woodland, roadsides (Mar–Jun)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	- - -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Phacelia breweri</i>	Brewer Phacelia	Boragin-	- - A2	- NP	Rocky soils, slopes, chaparral, oak woodland (Mar–Jun)	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	- - -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	- - -	- NA	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Phacelia divaricata</i>	Divaricate Phacelia	Boragin-	- - A2	- NA	Open areas, chaparral, woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Phacelia imbricata subsp. imbricata</i>	Rock Phacelia	Boragin-	- - -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Phacelia malvifolia</i>	Stinging Phacelia	Boragin-	- - A2	- NA	Sandy or gravelly soils, scrub, conifer forest (Apr–Jul)	
<input type="radio"/> <i>Phacelia ramosissima</i> (JM93: var. <i>ramosissima</i> , <i>latifolia</i>)	Branched Phacelia	Boragin-	- - A2	- NP	Diverse habitats incl salt marshes, canyons (Apr–Oct)	
<input type="radio"/> <i>Phoradendron serotinum subsp. macrophyllum</i> (JM93: <i>P. macrophyllum</i>)	Big-leaf Mistletoe	Visc-	- - C	- NP	On trees other than <i>Quercus</i> (especially <i>Alnus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Platanus</i> , <i>Populus</i> , <i>Robinia</i> , <i>Salix</i>) (Dec–Mar)	
<input type="radio"/> <i>Phoradendron serotinum subsp. tomentosum</i> (JM93: <i>P. villosum</i>)	Oak Mistletoe	Visc-	- - -	- NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)	
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Boragin-	- - -	- NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)	
<input type="radio"/> <i>Plagiobothrys fulvus</i> var. <i>campestris</i> (JM93: no var.)	Field Popcornflower	Boragin-	- - B	- NA	Grassland, open woodland, loamy, sandy, or gravelly soil (Mar–May)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	- - -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	- - -	- -		
<input type="radio"/> <i>Plagiobothrys stipitatus</i> var. <i>stipitatus</i>	Showy Great Popcornflower	Boragin-	- - C	- NA	Vernal pools, wet sites in grassland (Mar–Jun)	
<input type="radio"/> <i>Plagiobothrys tenellus</i>	Delicate Popcorn Flower	Boragin-	- - A2	- NA		
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	- - -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	- - -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	- - -	- NA	Open grassland, sandy soil, burns (Mar–May)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: subsp. <i>ciliosa</i> , <i>insignis</i>)	Longspur Plectritis	Valerian-	- - B	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis congesta subsp. brachystemon</i> (JM93: <i>P. brachystemon</i>)	Pale Plectritis	Valerian-	- - B	- NA	Common. Coastal bluffs, open, partly shaded slopes (Mar–Jun)	

Herbaceous <i>Plectritis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Plectritis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	- - -	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	- - -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Psilocarphus brevissimus</i> var. <i>brevissimus</i>	Dwarf Woollyheads	Aster-	- - B	- NA	Drying edges of vernal pools, mud flats, drainages (May–Jun)	
<input type="radio"/> <i>Psilocarphus oregonus</i>	Oregon Woollyheads	Aster-	- - B	- NA	Drying clay of vernal pools, drainages, moist rocky slopes (Mar–Jul)	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	- - -	- NA	Common. Sand or gravel (Mar–Jul)	
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	- - B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)	
<input type="radio"/> <i>Ranunculus aquatilis</i> var. <i>diffusus</i> (JM93: var. <i>capillaceus</i>)	Thread-leaf Water Buttercup	Ranuncul-	- - -	- NP	Ponds, lakes, streams (Mar–Sep)	
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	- - -	- NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Rigiopappus leptocladus</i>	Bristle-head	Aster-	- - B	- NA	Grassy sites, openings among shrubs or trees (Apr–Jul)	
<input type="radio"/> <i>Rumex californicus</i> (JM93: <i>R. salicifolius</i> var. <i>denticulatus</i>)	Toothed Willow Dock	Polygon-	- - A2	- NP		
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	- - -	L ZP	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	- - -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="radio"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i>)	Thin-leaf Willow Dock	Polygon-	- - B	- NP	Wet places, margins, rocky slopes (May–Jul)	
<input type="radio"/> <i>Rumex</i> sp.	Dock	Polygon-	- - -	- -		
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Caryophyll-	- - -	- NA	Sandy disturbed areas, river bars, streamsides (Spring–early summer)	
<input type="radio"/> <i>Sagina decumbens</i> subsp. <i>occidentalis</i>	Western Pearlwort	Caryophyll-	- - C	- NA	Dry streams, chaparral, grassy areas, rock outcrops, vernal pools (Spring–early summer)	
<input type="radio"/> <i>Salvia columbariae</i>	Chia	Lami-	- - B	- NA	Dry, disturbed sites, chaparral, coastal-sage scrub (Mar–Jun)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	- - -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	- - -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	- - -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula tuberosa</i>	Turkey Pea Sanicle	Api-	- - C	- NP	Open gravelly meadows, chaparral, woodland, pine forest (Mar–Jul)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: subsp. <i>californica</i> , <i>floribunda</i>)	California Figwort	Scrophulari-	- - -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	

Herbaceous <i>Scutellaria</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Scutellaria</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Scutellaria siphocampyloides</i>	Grayleaf Skullcap	Lami-	- - A1	- NP	Open sites, seeps, dry stream beds, scrub, woodland (May–Jul)	
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	- - C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	- - -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	- - -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sidalcea diploscypha</i>	Fringed Checkerbloom	Malv-	- - A2	- NA	Grassland, open woodland, valleys, occasionally near vernal pools, generally serpentine (Apr–May)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>laciniata</i>	Geranium-leaved Checkerbloom	Malv-	- - C	- NP	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	- - -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silene laciniata</i> subsp. <i>californica</i> (JM93: <i>S. californica</i>)	California Pink	Caryophyll-	- - B	- NP	Chaparral, oak woodland, conifer forest, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	- - -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	- - -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	- - -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	- - -	- ZAP	Forest, meadows, mud flats, disturbed	
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Lami-	- - C	- NP	Streambanks, springs, pine/oak forest (Jun–Oct)	
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i>)	Common Rigid Hedge-nettle	Lami-	- - -	- NP	Moist to ± dry places (Mar–Oct)	
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	- - -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	- - C	- NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)	
<input type="radio"/> <i>Stephanomeria virgata</i> subsp. <i>pleurocarpa</i>	Twiggy Wreath Plant	Aster-	- - C	- NA	Chaparral openings, grassland (Jun–Nov)	
<input type="radio"/> <i>Streptanthus glandulosus</i> subsp. <i>glandulosus</i> (JM93: incl. <i>S. albidus</i> subsp. <i>peramoenus</i>)	Bristly Jewel Flower	Brassic-	- - B	- NA	Serpentine, bare slopes, chaparral & woodland openings (Apr–Jul)	
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	- - -	- ZBP	Abundant. Esp disturbed areas (All year)	
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	- - C	- NP	Chaparral, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Tetrapteron graciliflorum</i> (JM93: <i>Camissonia graciliflora</i>)	Hill Sun Cup	Onagr-	- - A2	- NA	Open or shrubby slopes, generally clay soils, grassland, oak and Joshua-tree woodland (Mar–Apr)	
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	- - -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	

Herbaceous <i>Thysanocarpus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Thysanocarpus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	- - -	- NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)
<input type="radio"/> <i>Thysanocarpus laciniatus</i> var. <i>laciniatus</i> (JM93: no var.)	Narrowleaf Fringepod	Brassic-	- - B	- NA	Oak woodland, rocky ridges, slopes, chaparral, washes (Mar–May)
<input type="radio"/> <i>Tonella tenella</i>	Small-flowered Tonella	Plantagin-	- - A2	- NA	Moist, shaded places in chaparral, oak and mixed woodland (Mar–Jun)
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	- - -	M ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	- - -	- ZA	Disturbed places (Apr–Jun)
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i>)	Rancheria Clover	Fab-	- - C	- NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)
<input type="radio"/> <i>Trifolium barbigerum</i> (JM93: var. <i>barbigerum</i>)	Bearded Clover	Fab-	- - B	- NP	Wet meadows, open, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decepiens</i>	Deceiving Clover	Fab-	- - -	- NA	Open, grassy areas, forest (Apr–Jun)
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	- - -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Trifolium dichotomum</i> (JM93: <i>T. albopurpureum</i> var. <i>dichotomum</i>)	Branched Indian Clover	Fab-	- - B	- NA	Coastal dunes, open slopes, meadows, oak woodland, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	- - -	- ZA	Agricultural, disturbed areas, lawns (Spring)
<input type="radio"/> <i>Trifolium fucatum</i>	Bull Clover	Fab-	- - C	- NA	Locally abundant. Moist, open grassland, marshes, roadsides, occasionally saline or serpentine soils (Apr–Jun)
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i>)	Pinpoint Clover	Fab-	- - -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Trifolium grayi</i> (JM93: <i>T. barbigerum</i> var. <i>andrewsii</i>)	Gray's Clover	Fab-	- - A1?	- NA	Wet meadows, foothill slopes, pine woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	- - -	M ZA	Disturbed areas, roadsides (Apr–May)
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	- - -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Fab-	- - C	- NA	Woody or shrubby slopes, roadsides (Mar–Jun)
<input type="radio"/> <i>Trifolium olivaceum</i> (JM93: <i>T. albopurpureum</i> var. <i>olivaceum</i>)	Olive Clover	Fab-	- - B	- NA	Grassy slopes, valley meadows, disturbed areas (Apr–May)
<input type="radio"/> <i>Trifolium variegatum</i> var. ? (JM93: no var.)	White-tip Clover	Fab-	- - -	- NA	- -
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	- - -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Triphysaria eriantha</i> subsp. <i>eriantha</i>	Yellow Johnny-tuck	Orobanch-	- - -	- NA	Grassland, foothills (Mar–May)
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	- - -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)

Herbaceous <i>Triteleia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Triteleia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	- - -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)
<input type="radio"/> <i>Turritis glabra</i> (JM93: <i>Arabis</i> , var. <i>glabra</i>)	Tower Mustard	Brassic-	- - C	- NP	Open fields, meadows, slopes (Apr–Jul)
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	- - -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)
<input type="radio"/> <i>Urtica dioica</i> subsp. <i>holosericea</i>	Hoary Nettle	Urtic-	- - -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)
<input type="radio"/> <i>Verbascum thapsus</i>	Woolly Mullein	Scrophulari-	- - -	L ZB	Roadsides, streambanks, disturbed areas (May–Sep)
<input type="radio"/> <i>Veronica peregrina</i> subsp. <i>xalapensis</i>	Purslane Speedwell	Plantagin-	- - -	- NA	Moist places (Apr–Aug)
<input type="radio"/> <i>Vicia americana</i> subsp. <i>americana</i> (JM93: var. <i>americana</i>)	American Vetch	Fab-	- - -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Viola douglasii</i>	Douglas' Violet	Viol-	- - A1	- NP	Vernally moist flats, grassy slopes, often on serpentine (Feb–Jul)
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	- - C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)
<input type="radio"/> <i>Viola purpurea</i> subsp. <i>purpurea</i>	Mountain Violet	Viol-	- - A2	- NP	In openings or beneath shrubs, generally in yellow-pine forest or higher (Mar–Jul)
<input type="radio"/> <i>Viola purpurea</i> subsp. <i>quercetorum</i>	Oak Violet	Viol-	- - A2	- NP	Dry, grassy or brushy slopes, chaparral, generally below yellow-pine forest (Feb–Jul)
<input type="radio"/> <i>Viola sheltonii</i>	Shelton's Violet	Viol-	- - A2	- NP	Fir, pine, or oak woodland, rich or gravelly soil (Mar–Jul)
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	- - -	- NP	Grassland (Apr–Aug)
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	- - C	- NP	Gen shady sites (Mar–Jun)
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	- - -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))
<input type="radio"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Api-	- - C	- NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)

Woody	Grouped by Growth Form					Woody
Acer	Sorted Alphabetically by Scientific Name					Acer
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	- - -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	- - -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	- - -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	- - -	- NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Ros-	- - C	- NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Eric-	- - C	- NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Arctostaphylos manzanita</i> subsp. <i>manzanita</i>	Common / Giant Manzanita	Eric-	- - C	- NP	Chaparral, conifer forest (Feb–May)	
<input type="radio"/> <i>Arctostaphylos</i> sp.	Manzanita	Eric-	- - -	- NP	- -	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	- - -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	- - -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Baccharis salicifolia</i> subsp. <i>salicifolia</i> (JM93: no subsp.)	Mule Fat	Aster-	- - -	- NP	Riparian woodland, canyon bottoms, disturbed sites, often forming thickets (All year)	
<input type="radio"/> <i>Berberis pinnata</i> subsp. <i>pinnata</i>	Shinyleaf Oregon-grape	Berberid-	- - C	- NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	- - C	- NP	Dry, rocky slopes, chaparral (Mar–May)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	- - -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	- - C	- NP	Along streams, wet places (Jun–Sep)	
<input type="radio"/> <i>Ericameria linearifolia</i>	Interior Goldenbush	Aster-	- - -	- NP	Dry slopes, valleys, foothill and desert woodland, saltbush and creosote-bush scrub (Mar–May)	
<input type="radio"/> <i>Ericameria nauseosa</i> var. <i>mohavensis</i> (JM93: <i>Chrysothamnus nauseosus</i> subsp. <i>mohavensis</i>)	Mojave Rabbitbrush	Aster-	- - A2	- NP	Common. Dry scrub (Aug–Oct)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	- - -	- NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i>)	California Coffee Berry	Rhamn-	- - -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	- - C	- NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	- - -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	- - -	- NP	Moist woodland edges, rocky slopes (May–Aug)	

Woody	Grouped by Growth Form					Woody
Keckiella	Sorted Alphabetically by Scientific Name					Keckiella
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Keckiella corymbosa</i>	Redwood / Red Rock Penstemon	Plantagin-	- - A1	- NP	Rocky slopes in conifer or hardwood forests, (chaparral) (Jun–Oct)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Caprifoli-	- - -	- NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera subspicata</i> var. <i>denudata</i>	Southern Honeysuckle	Caprifoli-	- - B	- NP	Chaparral slopes (Jun–Jul)	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>albifrons</i>	Bush Lupine	Fab-	- - -	- NP	Common. Chaparral, foothill woodland (Mar–Jun)	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>collinus</i>	Bay Area Silver Lupine	Fab-	- - -	- NP	Cliffs, forest openings (Mar–Jun)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	- - -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Pinus coulteri</i>	Coulter Pine	Pin-	- - A2	- NP	Chaparral, lower mixed-conifer, mixed-hardwood forests -	
<input type="radio"/> <i>Pinus sabiniana</i>	Gray Pine	Pin-	- - -	- NP	Foothill woodland, n oak woodland, chaparral, infertile soils in mixed-conifer and hardwood forests -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	- - B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Ros-	- - C	- NP	Mixed-evergreen or conifer forest (Mar–May)	
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Ros-	- - -	- NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	- - A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	- - -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	- - -	- NP	Dry slopes, interior foothills, woodland (Apr–May)	
<input type="radio"/> <i>Quercus garryana</i> var. <i>garryana</i>	Oregon / Garry Oak	Fag-	- - B	- NP	Slopes, mixed-evergreen or conifer forest (Apr–Jun)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	- - -	- NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	- - B	- NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	- - -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Rhamn-	- - C	- NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	- - C	- NP	Chaparral, montane forest (Mar–Jun)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	- - -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	- - -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	- - -	- NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa</i> sp.	Rose	Ros-	- - -	- NP	--	

Woody	Grouped by Growth Form						Woody
Salix	Sorted Alphabetically by Scientific Name						Salix
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="checkbox"/> <i>Salix sp.</i>	Willow	Salic-	- - -	-	?P	- -	
<input type="checkbox"/> <i>Salvia mellifera</i>	Black Sage	Lami-	- - -	-	NP	Coastal-sage scrub, lower chaparral (Mar–Jun)	
<input type="checkbox"/> <i>Sambucus nigra subsp. caerulea (JM93: S. mexicana)</i>	Blue Elderberry	Adox-	- - -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="checkbox"/> <i>Senecio flaccidus var. douglasii</i>	Douglas' Threadleaf Ragwort	Aster-	- - A2	-	NP	Generally disturbed dry, open, sandy or rocky sites (Jun–Oct)	
<input type="checkbox"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	- - -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="checkbox"/> <i>Symphoricarpos albus var. laevigatus</i>	Snowberry	Caprifoli-	- - -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="checkbox"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	- - -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="checkbox"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	- - -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="checkbox"/> <i>Umbellularia californica</i>	California Bay	Laur-	- - -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)	

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex deweyana ssp. *leptopoda* (*C. leptopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thyrsoiflorus (var. *thyrsoiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria amphibia*)

Polygonum amphibium var. *stipulaceum* (*Persicaria amphibia*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)