

Redwood Regional Park

Checklist of Wild Plants

Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Redwood Regional Park. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Family	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

Redwood Regional Park Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name					<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	- - -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	Western Lady Fern	Woods-	- - -	- NP	Woodland, along streams, seepage area -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	- - -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum arvense</i>	Common Horsetail	Equiset-	- - -	- NP	Streambanks, wet meadows, springs, other wet, shaded places -	
<input type="radio"/> <i>Equisetum telmateia</i> subsp. <i>braunii</i>	Giant Horsetail	Equiset-	- - -	- NP	Streambanks, roadside ditches, seepage areas -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	- - -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Pterid-	- - C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Pterid-	- - -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	- - -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Dryopterid-	- - -	- NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -	
<input type="radio"/> <i>Pteridium aquilinum</i> var. <i>pubescens</i>	Bracken Fern	Dennstaedti-	- - -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	- - -	- NP	Near streams, springs, seeps -	

Grasses & Grass-like	Grouped by Growth Form				Grasses & Grass-like
<i>Agrostis</i>	Sorted Alphabetically by Scientific Name				<i>Agrostis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis capillaris</i>	Colonial Bent Grass	Po-	- - -	- ZP	Roadsides, open, disturbed places (Jul-Sep)
<input type="radio"/> <i>Agrostis hallii</i>	Hall's Bent Grass	Po-	- - A2	- NP	Open oak woodland, conifer forest (May-Jul)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	- - -	- NP	Common. Open meadows, woodland, forest, subalpine (Jun-Aug)
<input type="radio"/> <i>Agrostis stolonifera</i>	Creeping Bent Grass	Po-	- - -	L ZP	Ditches, lake margins, marshes (Jun-Sep)
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	- - -	- ZA	Sandy soils, open or disturbed sites (Apr-Jun)
<input type="radio"/> <i>Alopecurus saccatus</i>	Pacific Foxtail	Po-	- - B	- NA	Vernal pools, moist, open meadows (Mar-May)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	- - -	M ZA	Disturbed sites (Mar-Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	- - -	M ZA	Disturbed sites (Apr-Jun)
<input type="radio"/> <i>Briza maxima</i>	Rattlesnake Grass	Po-	- - -	L ZA	Shaded sites, roadsides, pastures, weedy on coastal dunes (Apr-Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	- - -	- ZA	Shaded or moist, open sites (Apr-Jul)
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	- - -	- NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)
<input type="radio"/> <i>Bromus catharticus</i> var. <i>catharticus</i> (JM93: no var.)	Rescue Grass	Po-	- - -	- ZAP	Open, disturbed places (Apr-Nov)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	- - -	M ZA	Open, gen disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	- - -	L ZA	Fields, disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus japonicus</i>	Japanese Brome	Po-	- - -	L ZA	Open, disturbed areas (May-Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	- - -	- NP	Shrubland, conifer forest, shaded streambanks, roadsides (May-Jul)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>madritensis</i>	Foxtail Chess	Po-	- - -	- ZA	Disturbed areas, roadsides (Apr-Jan)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	- - -	H ZA	Disturbed areas, roadsides (Mar-Jun)
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	- - B	- NP	Seasonally wet places (May-Aug)
<input type="radio"/> <i>Carex brevicaulis</i>	Short-stemmed Sedge	Cyper-	- - A1	- NP	Coastal dunes, headlands, rocky or sandy soil (Apr-May)
<input type="radio"/> <i>Carex globosa</i>	Round-fruited Sedge	Cyper-	- - A1	- NP	Well-drained soil of wooded areas, edges (Apr-Jun)
<input type="radio"/> <i>Carex leptopoda</i> (JM93: <i>C. deweyana</i> subsp. <i>leptopoda</i>)	Slender-footed Sedge	Cyper-	- - A1	- NP	Moist soil, wooded areas (May-Aug)
<input type="radio"/> <i>Carex serratodens</i>	Bifid Sedge	Cyper-	- - B	- NP	Moist places (Apr-Jul)
<input type="radio"/> <i>Carex tumulicola</i>	Foothill Sedge	Cyper-	- - C	- NP	Meadows, open woodland (Apr-Jul)
<input type="radio"/> <i>Cortaderia jubata</i>	Hairy Pampas Grass	Po-	- - -	H ZP	Disturbed sites, many habitats, esp coastal (Sep-Feb)
<input type="radio"/> <i>Cortaderia selloana</i>	Smooth Pampas Grass	Po-	- - -	H ZP	Disturbed sites (Sep-Mar)

Grasses & Grass-like <i>Cynodon</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Cynodon</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Po-	- - -	M	ZP	Disturbed sites (Jun–Aug)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	- - -	M	ZA	Open, disturbed sites (May–Jul)
<input type="radio"/> <i>Cyperus acuminatus</i>	Tapertip Flatsedge	Cyper-	- - -	-	NA	Edges of temporary pools, ponds, streams (Jun–Oct)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	- - -	-	NP	Vernal pools, streambanks (May–Nov)
<input type="radio"/> <i>Cyperus sp.</i>	Nutsedge	Cyper-	- - -	-	NAP	- -
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	- - -	L	ZP	Disturbed, often moist sites (May–Aug)
<input type="radio"/> <i>Danthonia californica</i> (JM93: <i>var. californica</i>)	California Oat Grass	Po-	- - C	-	NP	Gen moist meadows, open woodland (Apr–Aug)
<input type="radio"/> <i>Deschampsia danthonioides</i>	Annual Hair Grass	Po-	- - C	-	NA	Moist to drying, open sites, meadows, streambanks, vernal pools, occ alkali soil (Mar–Aug)
<input type="radio"/> <i>Deschampsia elongata</i>	Slender Hair Grass	Po-	- - C	-	NP	Wet sites, meadows, lakeshores, shaded slopes (May–Sep)
<input type="radio"/> <i>Digitaria sanguinalis</i>	Hairy Crab Grass	Po-	- - -	-	ZA	Disturbed areas (Jun–Sep)
<input type="radio"/> <i>Elymus glaucus subsp. glaucus</i>	Western Wild-rye	Po-	- - -	-	NP	Open areas, chaparral, woodland, forest (Jun–Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	- - C	-	NP	Open, sandy to rocky areas (May–Jul)
<input type="radio"/> <i>Elymus trachycaulus subsp. trachycaulus</i>	Slender Wheat Grass	Po-	- - B	-	NP	Dry to moist, open areas, forest, woodland (Jun–Aug)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i>)	Beardless Wild Rye	Po-	- - -	-	NP	Dry to moist, often saline, meadows (Jun–Jul)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	- - -	M	ZP	Disturbed places (May–Jun)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i>)	Brome Fescue	Po-	- - -	-	ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	- - C	-	NP	Dry, open forest, moist streambanks, chaparral (May–Jun)
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Po-	- - C	-	NP	Dry, open or shady places (Jul–Sep)
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , <i>var. ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i>)	Hairy Fescue	Po-	- - -	-	NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , <i>var. hirsuta</i> , <i>myuros</i>)	Rattail Sixweeks Grass	Po-	- - -	-	ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i>)	Rye Grass	Po-	- - -	-	ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Po-	- - B	-	NP	Sand dunes, grassland, subalpine forest (May–Jul)
<input type="radio"/> <i>Festuca sp. (annual)</i> (JM93: <i>Vulpia sp.</i>)	Annual Fescue	Po-	- - -	-	NA	- -
<input type="radio"/> <i>Festuca sp. (perennial)</i>	Perennial Fescue	Po-	- - -	-	- -	- -
<input type="radio"/> <i>Gastridium phleoides</i> (JM93: <i>G. ventricosum</i>)	Nit Grass	Po-	- - -	-	ZA	Open, generally dry, disturbed sites (Apr–Nov)

Grasses & Grass-like <i>Holcus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Holcus</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Holcus lanatus</i>	Common Velvet Grass	Po-	- - -	M	ZP	Moist sites, roadbanks, cult fields, meadows (Jun–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Po-	- - -	-	NP	Meadows, pastures, streambanks (May–Aug)
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>californicum</i>	California Barley	Po-	- - B	-	NP	Meadows, pastures, streambanks (Mar–Jul)
<input type="radio"/> <i>Hordeum jubatum</i> subsp. <i>jubatum</i> (JM93: no subsp.)	Foxtail Barley	Po-	- - B	-	NAP	Roadsides, disturbed areas, meadows, marshes (May–Jul)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	- - -	-	ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>glaucum</i>	Smooth Barley	Po-	- - -	-	ZA	Moist, generally disturbed sites (Mar–Jul)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	- - -	-	ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Juncus balticus</i> subsp. <i>ater</i> (JM93: no subsp.)	Baltic Rush	Junc-	- - -	-	NP	Moist to ± dry sites (Jul–Nov)
<input type="radio"/> <i>Juncus bufonius</i> var. ?	Toad Rush	Junc-	- - -	-	NA	- -
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Junc-	- - -	-	NA	Damp sunny ground, gen disturbed (May–Sep)
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Junc-	- - -	-	NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)
<input type="radio"/> <i>Juncus occidentalis</i>	Western Rush	Junc-	- - -	-	NP	Moist gen sunny areas (May–Sep)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	- - -	-	NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus tenuis</i>	Slender Rush	Junc-	- - -	-	NP	Uncommon. Damp places (Jun–Sep)
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	- - -	-	NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	- - C	-	NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	- - -	-	NP	Meadows, open woodland, conifer forest (Jun–Jul)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	- - C	-	NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	- - -	-	NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	- - -	-	NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Phalaris aquatica</i>	Harding Grass	Po-	- - -	M	ZP	Disturbed areas, roadsides (Apr–Aug)
<input type="radio"/> <i>Phalaris minor</i>	Little-seeded Canary Grass	Po-	- - -	-	ZA	Disturbed areas, cultivated fields (Apr–Jul)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	- - -	-	ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa howellii</i>	Howell Blue Grass	Po-	- - B	-	NA	Rocky banks, shaded slopes of woodland, chaparral, disturbed places (Apr–Jun)
<input type="radio"/> <i>Poa pratensis</i> subsp. <i>pratensis</i>	Kentucky Blue Grass	Po-	- - -	-	ZP	Common. Disturbed habitats, including saline or alkaline soils (Mar–Jul)

Grasses & Grass-like <i>Poa</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Poa</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	- - -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Po-	- - -	- ZP	Common. Streambanks (May–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	- - -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Scirpus microcarpus</i>	Smallfruit Bulrush	Cyper-	- - -	- NP	Marshes, wet meadows, streambanks, pond margins, sometimes weedy (Jun–Jul)
<input type="radio"/> <i>Sorghum bicolor</i>	Sorghum	Po-	- - -	- ZA	Disturbed areas, roadsides (All year)
<input type="radio"/> <i>Stipa cernua</i> (JM93: <i>Nassella</i>)	Nodding Needle Grass	Po-	- - C	- NP	Grassland, chaparral, juniper woodland (Feb–Jul)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i>)	Foothill Needle Grass	Po-	- - C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Po-	- - C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Trisetum canescens</i>	Tall False Oat	Po-	- - B	- NP	Open to shaded sites, meadows, chaparral, conifer forest (May–Aug)
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Typh-	- - -	- NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)

Herbaceous <i>Acaena</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Acaena</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Acaena pinnatifida</i> var. <i>californica</i>	California Acaena	Ros-	- - C	- NP	Coastal grassland, open, rocky slopes (Mar–May)
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	- - -	- NP	Many habitats (Apr–Sep)
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Aster-	- - -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i>)	Spanish Clover	Fab-	- - -	- NA	
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Fab-	- - -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Fab-	- - -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i>)	Small-flower Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Fab-	- - -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Actaea rubra</i>	Baneberry	Ranuncul-	- - B	- NP	Deep soils, moist, open to shaded sites, mixed-evergreen or conifer forests (May–Sep)
<input type="radio"/> <i>Agoseris apargioides</i> var. <i>apargioides</i>	Seaside Dandelion	Aster-	- - A1	- NP	Coastal dunes, sand hills (Apr–May)
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	- - -	- NP	Grassland, scrub, woodland (Apr–Jul)
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Aster-	- - -	- NA	Many open habitats (May–Jun)
<input type="radio"/> <i>Agoseris retrorsa</i>	Spearleaf Native Dandelion	Aster-	- - B	- NP	Scrub, oak woodland, conifer forest (Apr–Aug)
<input type="radio"/> <i>Allium falcifolium</i>	Sickle Leaf Onion	Alli-	- - A2	- NP	Common. Heavy clays including serpentine (Apr–Jun)
<input type="radio"/> <i>Allium triquetrum</i>	Three Cornered Leek	Alli-	- - -	- ZP	Locally common. Shady ± disturbed places (Mar–Apr)
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Alli-	- - C	- NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)
<input type="radio"/> <i>Allophyllum gilioides</i> subsp. <i>gilioides</i>	Purple False Gilia	Polemoni-	- - B	- NA	Open, sandy, gen damp or grassy areas (Apr–Jun)
<input type="radio"/> <i>Amaranthus deflexus</i>	Low Amaranth	Amaranth-	- - -	- ZA	Railroad right-of-ways, disturbed areas (May–Nov)
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Boragin-	- - -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)
<input type="radio"/> <i>Anaphalis margaritacea</i>	Pearly Everlasting	Aster-	- - -	- NP	Woodland, disturbed places (Jul–Oct)
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	- - -	- NP	Generally wooded areas (Jun–Aug)

Herbaceous <i>Anthemis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Anthemis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	- - -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	- - -	- ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	- - -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Apocynum androsaemifolium</i>	Bitter Dogbane	Apocyn-	- - B	- NP	Open slopes, rocky places, with conifers, chaparral (May–Oct)	
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	- - C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)	
<input type="radio"/> <i>Aralia californica</i>	Elk Clover	Arali-	- - B	- NP	Moist shade, canyons, stream sides (Jun–Aug)	
<input type="radio"/> <i>Arnica discoidea</i>	Rayless Arnica	Aster-	- - B	- NP	Chaparral, foothill woodland (May–Jul)	
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	- - -	- NP	Common. Open to shady areas, often in drainages (May–Nov)	
<input type="radio"/> <i>Asarum caudatum</i>	Wild Ginger	Aristolochi-	- - A2	- NP	Moist places in forest (Mar–Aug)	
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvetch	Fab-	- - -	- NA	Open, grassy areas, scrub (Mar–Jul)	
<input type="radio"/> <i>Baccharis glutinosa (JM93: B. douglasii)</i>	Marsh Baccharis	Aster-	- - -	- NP	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	- - -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	- - -	L ZA	Disturbed grassland. (Apr–Jun)	
<input type="radio"/> <i>Bellis perennis</i>	English Daisy	Aster-	- - -	- ZP	Damp, grassy areas (Dec–Sep)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	- - -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	- - -	M ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	- - -	L ZA	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Themid-	- - -	- NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	- - -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	- - -	- NP	Common. Shady to open woodland, scrub (Apr–Jun)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	- - C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus umbellatus</i>	Oakland Star-tulip	Lili-	- 4.2 *A2	- NP	Open chaparral or woodland, gen on serpentine (Mar–May)	
<input type="radio"/> <i>Calycadenia multiglandulosa</i>	Sticky Rosinweed	Aster-	- - A1	- NA	Common. Gen dry, open valleys, hillsides, rocky ridges (May–Oct)	

Herbaceous <i>Calystegia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Calystegia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Calystegia purpurata</i> subsp. <i>purpurata</i>	Climbing Morning-glory	Convolvul-	- - -	- NP	Chaparral, coastal scrub (May–Jun)
<input type="radio"/> <i>Calystegia subacaulis</i> subsp. <i>subacaulis</i>	Shortstem Morning-glory	Convolvul-	- - -	- NP	Dry, open scrub or woodland (Apr–Jun)
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	- - -	- ZA	Disturbed areas (Jan–Oct)
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	- - -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	- - -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)
<input type="radio"/> <i>Carduus pycnocephalus</i> subsp. <i>pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Aster-	- - -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Carduus tenuiflorus</i>	Slender Flower Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Apr–Jun)
<input type="radio"/> <i>Castilleja affinis</i> subsp. <i>affinis</i>	Common Indian Paintbrush	Orobanch-	- - -	- NP	Chaparral, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Castilleja densiflora</i> subsp. <i>densiflora</i>	Common Owl's-clover	Orobanch-	- - -	- NA	Grassland (Mar–May)
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	- - -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)
<input type="radio"/> <i>Castilleja subinclusa</i> subsp. <i>franciscana</i>	Franciscan Paintbrush	Orobanch-	- - A2	- NP	Coastal scrub (Mar–Jul)
<input type="radio"/> <i>Castilleja wightii</i>	Seaside / Coastal Paintbrush	Orobanch-	- - C	- NP	Coastal scrub (Mar–Aug)
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i>)	California Mustard	Brassic-	- - -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	- - -	M ZA	Disturbed fields, open woodland (Apr–Jul)
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	- - -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Centranthus ruber</i>	Red Valerian	Valerian-	- - -	- ZP	Disturbed places, rock or wall crevices, roadsides (Apr–Jul)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	- - -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	- - -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Cicendia quadrangularis</i>	Timwort	Gentian-	- - B	- NA	Open places (Mar–May)
<input type="radio"/> <i>Cirsium quercetorum</i>	Brownie Thistle	Aster-	- - A2	- NP	Open places, grassland, woodland (Apr–Aug)
<input type="radio"/> <i>Cirsium</i> sp.	Thistle	Aster-	- - -	- -	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	- - -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia franciscana</i>	Presidio Clarkia	Onagr-	FE/CE 1B.1 *A1	- NA	Serpentine soil (May–Jun)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>purpurea</i>	Purple Winecup Clarkia	Onagr-	- - A2	- NA	Uncommon. Grassland (Apr–Jun)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	- - -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)

Herbaceous <i>Clarkia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Clarkia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>viminea</i>	Large Clarkia	Onagr-	- - A2	- NA	Open, grassy or shrubby places (May–Jul)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	- - -	- NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>exigua</i>	Common Pale Claytonia	Monti-	- - C	- NA	Dry or moist, disturbed bare clay to sandy soils, often serpentine (Apr–Jul)
<input type="radio"/> <i>Claytonia parviflora</i> subsp. ?	Miner's Lettuce	Monti-	- - -	- NA	--
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i>)	Small-flower Claytonia	Monti-	- - -	- NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	- - -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Monti-	- - -	- NA	Vernally moist, often shady or disturbed sites (Jan–May)
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i>)	Yerba Buena	Lami-	- - -	- NP	Shady places, chaparral, woodland (Apr–Sep)
<input type="radio"/> <i>Clintonia andrewsiana</i>	Red-bead Lily	Lili-	- - A1	- NP	Shaded, damp redwood forest (May–Jul)
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	- - -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	- - -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)
<input type="radio"/> <i>Corallorhiza maculata</i> var. <i>maculata</i> (JM93: no var.)	Spotted Coralroot	Orchid-	- - A2	- NP	Shaded mixed-evergreen or conifer forest, in decomposing lf litter (May–Aug)
<input type="radio"/> <i>Corallorhiza striata</i>	Striped Coralroot	Orchid-	- - A1	- NP	Open to shaded mixed-evergreen or conifer forest, in decomposing leaf litter (Feb–Jul)
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	- - -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	- - -	- NA	Open areas (Feb–May)
<input type="radio"/> <i>Cymbalaria muralis</i>	Kenilworth Ivy	Plantagin-	- - -	- ZP	Rock walls, shady, disturbed areas (May–Sep)
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Aster-	- - -	M ZP	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	- - -	- NP	Chaparral, woodland (Feb–May)
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	- - -	- NA	Rocky or sandy places (Apr–Jun)
<input type="radio"/> <i>Delairea odorata</i> (JM93: <i>Senecio mikanioides</i>)	German Ivy	Aster-	- - -	H ZP	Shady, ± disturbed places, riparian woodland, coastal scrub (Nov–Mar)
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Ranuncul-	- - C	- NP	Open coastal grassland, chaparral (Mar–May)
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	- - -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	- - -	- NP	Open woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Digitalis purpurea</i>	Foxglove	Plantagin-	- - -	L ZB	Acid soil in open woodland, disturbed areas (May–Jul)

Herbaceous <i>Dipsacus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Dipsacus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Dipsacus sativus</i>	Fuller's Teasel	Dipsac-	- - -	M ZB	Disturbed areas, fields, vacant lots, pastures (May–Jul)	
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	- - -	- NP	Gen in shady sites (Mar–Jul)	
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i>)	Sticky Cinquefoil	Ros-	- - -	- NP	Gen ± shady or moist areas (May–Jul)	
<input type="radio"/> <i>Duchesnea indica</i> var. <i>indica</i> (JM93: no var.)	Mock-strawberry	Ros-	- - -	- ZP	Disturbed areas (May–Aug)	
<input type="radio"/> <i>Dysphania multifida</i> (JM93: <i>Chenopodium</i>)	Cutleaf Goosefoot	Chenopodi-	- - -	- ZA	Disturbed areas (Jul–Oct)	
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	- - -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)	
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	- - -	- NP	Dry slopes, ridges (Jun–Dec)	
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Onagr-	- - -	- NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)	
<input type="radio"/> <i>Epilobium minutum</i>	Chaparral Willowherb	Onagr-	- - -	- NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)	
<input type="radio"/> <i>Epilobium</i> sp.	Willowherb	Onagr-	- - -	- NA	--	
<input type="radio"/> <i>Erigeron sumatrensis</i> (JM93: <i>Conyza bilbaoana</i>)	Tropical Horseweed	Aster-	- - -	- ZA	Disturbed sites (All year)	
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>caninum</i>	Tiburon Buckwheat	Polygon-	- 1B.2 *A1	- NA	Serpentine (May–Oct)	
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>luteolum</i>	Golden-carpet Wild Buckwheat	Polygon-	- - A2	- NA	Common. Serpentine (Jul–Nov)	
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	- - -	- NP	--	
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Polygon-	- - A1	- NP	Common. Sand or gravel (May–Oct)	
<input type="radio"/> <i>Eriogonum</i> sp.	Wild Buckwheat	Polygon-	- - -	- NAP	--	
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Erodium brachycarpum</i>	Southern European Filaree	Gerani-	- - -	- ZA	Dry, open or disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	- - -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)	
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	- - -	- ZA	Open, disturbed sites (Feb–Sep)	
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	- - -	- NP	Grassy, open areas (Feb–Sep)	
<input type="radio"/> <i>Euphorbia pepylus</i>	Petty Spurge	Euphorbi-	- - -	- ZA	Common. Disturbed areas (Feb–Aug)	
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i>)	Broadleaf Aster	Aster-	- - -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)	
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	- - -	H ZP	Roadsides, disturbed sites (May–Sep)	
<input type="radio"/> <i>Fragaria vesca</i>	Wood Strawberry	Ros-	- - -	- NP	Gen partial shade in forest (Jan–Jul)	
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i>)	Checker Lily	Lili-	- - -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)	
<input type="radio"/> <i>Galium andrewsii</i> subsp. <i>gatense</i>	Phlox-leaf Serpentine Bedstraw	Rubi-	- 4.2 *A2	- NP	Dry, rocky places in serpentine soil, chaparral or open oak/pine woodland (Apr–Jun)	

Herbaceous	Grouped by Growth Form					Herbaceous
Galium	Sorted Alphabetically by Scientific Name					Galium
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	- - -	- NA	Grassy, ± shady places (Apr–Jun)	
<input type="radio"/> <i>Galium californicum subsp. californicum</i>	California Bedstraw	Rubi-	- - -	- NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)	
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	- - -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)	
<input type="radio"/> <i>Galium porrigens var. porrigens</i>	Climbing Bedstraw	Rubi-	- - -	- NP	Among shrubs in chaparral, forest (May–Aug)	
<input type="radio"/> <i>Galium triflorum</i>	Sweet-scent Bedstraw	Rubi-	- - B	- NP	Damp, shady forest (May–Jul)	
<input type="radio"/> <i>Gamochaeta ustulata (JM93: Gnaphalium purpureum)</i>	Purple Cudweed	Aster-	- - -	- NA	Dunes, bluffs, fields, disturbed sites (Apr–Jul)	
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	- - -	M ZA	Open, disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	- - -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)	
<input type="radio"/> <i>Geranium robertianum</i>	Herb Robert	Gerani-	- - -	- ZAB	Open to shaded sites (Apr–Sep)	
<input type="radio"/> <i>Gilia achilleifolia subsp. achilleifolia</i>	California Gilia	Polemoni-	- - B	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)	
<input type="radio"/> <i>Gilia sp.</i>	Gilia	Polemoni-	- - -	- NA	--	
<input type="radio"/> <i>Gnaphalium or Pseudognaphalium sp. (JM93: Gnaphalium sp.)</i>	Everlasting Cudweed	Aster-	- - -	-	--	
<input type="radio"/> <i>Grindelia camporum (JM93: var. camporum, G. hirsutula var. davyi)</i>	Great Valley Gumplant	Aster-	- - -	- NP	Sandy or saline bottomland, roadsides (May–Nov)	
<input type="radio"/> <i>Grindelia hirsutula (JM93: var. hirsutula)</i>	Hairy Gumplant	Aster-	- - C	- NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)	
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	- - -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)	
<input type="radio"/> <i>Helminthotheca echioides (JM93: Picris)</i>	Bristly Ox-tongue	Aster-	- - -	L ZAB	Common. Disturbed areas (All year)	
<input type="radio"/> <i>Hemizonia congesta subsp. ?</i>	Yellow Hayfield Tarweed	Aster-	- - -	- NA	--	
<input type="radio"/> <i>Hemizonia congesta subsp. luzulifolia</i>	Hayfield Tarweed	Aster-	- - -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)	
<input type="radio"/> <i>Heracleum maximum (JM93: H. lanatum)</i>	Cow Parsnip	Api-	- - -	- NP	Moist places, wooded or open (Apr–Jul)	
<input type="radio"/> <i>Hesperevax sparsiflora var. sparsiflora</i>	Erect Hesperevax	Aster-	- - -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)	
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	- - -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)	
<input type="radio"/> <i>Heterotheca sessiliflora subsp. bolanderi</i>	Bolander Goldenaster	Aster-	- - C	- NP	Dunes, headlands, grassy coastal slopes (Jun–Sep)	
<input type="radio"/> <i>Heterotheca sessiliflora subsp. echioides</i>	Bristly Goldenaster	Aster-	- - -	- NP	Grassland, scrub, woodland, open forest, disturbed sites (Jul–Oct)	

Herbaceous <i>Heuchera</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Heuchera</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Heuchera micrantha</i>	Small-flower Alumroot	Saxifrag-	- - C	- NP	Moist, rocky banks and cliffs (Apr–Jul)
<input type="radio"/> <i>Hieracium albiflorum</i>	White Hawkweed	Aster-	- - B	- NP	Forest (May–Sep)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	- - -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Hypericum perforatum</i>	Klamathweed	Hyperic-	- - -	M ZP	Open, disturbed areas in many plant communities (May–Aug)
<input type="radio"/> <i>Hypericum scouleri</i> (JM93: <i>H. formosum</i> var. <i>scouleri</i>)	Scouler St. John's Wort	Hyperic-	- - A1	- NP	
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	- - -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	- - -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Iris douglasiana</i>	Douglas Iris	Irid-	- - A2	- NP	Common. Grassy places, esp near coast (May–Jul)
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	- - -	- ZA	Roadsides, grassland (Jul–Nov)
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	- - -	- ZA	Abundant. Disturbed places (May–Oct)
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	- - -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: <i>subsp. ramosissima</i>)	Common Hare-leaf	Aster-	- - -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lasthenia californica subsp. californica</i> (JM93: <i>no subsp.</i>)	California Goldfields	Aster-	- - -	- NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Fab-	- - -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Lepidium nitidum</i> (JM93: <i>var. nitidum, oreganum</i>)	Threadleaf Peppergrass	Brassic-	- - -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)
<input type="radio"/> <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	- - -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i>)	Pinklobe Leptosiphon	Polemoni-	- - C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i>)	Bicolor Leptosiphon	Polemoni-	- - -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	- - -	- NP	Open, grassy slopes (Mar–Apr)
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	- - -	- NP	Shaded slopes (Feb–Jun)
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i>)	California Fluffweed	Aster-	- - -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Aster-	- - -	- ZA	Bare or grassy openings, burns (Mar–Jul)
<input type="radio"/> <i>Lomatium dasycarpum subsp. dasycarpum</i>	Woolly Fruited Desertparsley	Api-	- - -	- NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)

Herbaceous <i>Lomatium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Lomatium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Api-	- - -	- NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium sp.</i>	Biscuit Root	Api-	- - -	- NP	--	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	- - -	- NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Lotus corniculatus</i>	Bird's-foot Trefoil	Fab-	- - -	- ZP	Open, disturbed areas (Jun–Sep)	
<input type="radio"/> <i>Lunaria annua</i>	Money Plant	Brassic-	- - -	- ZA	Disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Lupinus adsurgens</i>	Drew's Silky Lupine	Fab-	- - -	- NP	Dry slopes, montane forest (May–Jul)	
<input type="radio"/> <i>Lupinus affinis</i>	Sky Lupine	Fab-	- - A1	- NA	Uncommon. Open areas (Mar–May)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus formosus var. formosus</i>	Summer Lupine	Fab-	- - -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	
<input type="radio"/> <i>Lupinus microcarpus var. microcarpus</i>	Chick Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	- - -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	- - -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Madia exigua</i>	Thread-stem Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites, in sandy or clayey soils, including serpentine (Apr–Jul)	
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	- - -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	- - -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum racemosum (JM93: Smilacina racemosa)</i>	Fat False Solomon's Seal	Rusc-	- - -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum (JM93: Smilacina stellata)</i>	Starry False Solomon's Seal	Rusc-	- - -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
<input type="radio"/> <i>Malva sp.</i>	Mallow	Malv-	- - -	-	--	
<input type="radio"/> <i>Marah fabacea (JM93: M. fabaceus)</i>	California Man-root	Cucurbit-	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marah oregana (JM93: M. oreganus)</i>	Coast Man-root	Cucurbit-	- - C	- NP	Shrubby or open areas, forest edges (Mar–May)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea (JM93: Chamomilla suaveolens)</i>	Pineapple Weed	Aster-	- - -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	

Herbaceous <i>Medicago</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Medicago</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Medicago arabica</i>	Spotted Bur Clover	Fab-	- - -	- ZA	Disturbed and agricultural areas, fields, woodland (Mar–Jun)
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Fab-	- - -	- ZAB	Open, disturbed areas (Apr–Oct)
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i>)	California Saxifrage	Saxifrag-	- - -	- NP	Moist, shady places (Feb–May(Jun))
<input type="radio"/> <i>Micropus californicus</i> var. ?	Slender Cottonweed	Aster-	- - -	- NA	--
<input type="radio"/> <i>Microseris acuminata</i>	Sierra Foothill Silverpuffs	Aster-	- - B	- NA	Grassy, open rocky or clay soil (Apr–Jun)
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	- - -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)
<input type="radio"/> <i>Microseris</i> sp.	Silverpuffs	Aster-	- - -	- NA	--
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i>)	Slender Annual Phlox	Polemoni-	- - -	- NA	Dry to moist areas (Mar–Aug)
<input type="radio"/> <i>Mimulus cardinalis</i>	Scarlet Monkeyflower	Phrym-	- - C	- NP	Moist to wet places along streams, seepage areas (May–Sep)
<input type="radio"/> <i>Mimulus douglasii</i>	Mickey Mouse Monkeyflower	Phrym-	- - A2	- NA	Bare clay, serpentine or granitic soils; generally along upper banks of small creeks (Feb–Apr)
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	- - -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)
<input type="radio"/> <i>Mimulus</i> sp.	Monkey Flower	Phrym-	- - -	- NAP	
<input type="radio"/> <i>Minuartia douglasii</i>	Douglas' Stitchwort	Caryophyll-	- - B	- NA	Dry, rocky, slopes, flats in chaparral, oak and pine woodland, often serpentine (Spring–early summer)
<input type="radio"/> <i>Monardella douglasii</i> (JM93: subsp. <i>douglasii</i>)	Douglas Coyotemint	Lami-	- - A2	- NA	Grassland, oak woodland, chaparral, serpentine outcrops (May–Nov)
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>globosa</i> , <i>M. antonina</i> subsp. <i>antonina</i>)	Coyote-mint	Lami-	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)
<input type="radio"/> <i>Myosotis latifolia</i>	Broadleaved Forget-me-not	Boragin-	- - -	L ZP	Moist, disturbed, shady places (Feb–Jul)
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i>)	Water Cress	Brassic-	- - -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Polemoni-	- - C	- NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	- - -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	- - -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Api-	- - -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)

Herbaceous <i>Oxalis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Oxalis</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Oxalis corniculata</i>	Creeping Wood Sorrel	Oxalid-	- - -	- ZP	Disturbed areas (Most of Year)	
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	- - -	M ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)	
<input type="radio"/> <i>Oxalis pilosa</i> (JM93: <i>O. albicans</i> subsp. <i>pilosa</i>)	Hairy Wood Sorrel	Oxalid-	- - A2	- NP	Coastal grassland, scrub, chaparral (Feb–Sep)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Orobanch-	- - -	- NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	- - -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Perideridia oregana</i>	Oregon Yampah	Api-	- - A2	- NP	Open flats or slopes, pine/oak woodland (Jul–Aug)	
<input type="radio"/> <i>Perideridia</i> sp.	Yampah	Api-	- - -	- NP	-	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	- - -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	- - -	- NA	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Phacelia divaricata</i>	Divaricate Phacelia	Boragin-	- - A2	- NA	Open areas, chaparral, woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Phacelia heterophylla</i> subsp. <i>virgata</i>	Varileaf Phacelia	Boragin-	- - -	- NBP	Slopes, flats, roadsides (May–Jul)	
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	- - -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Phacelia nemoralis</i> subsp. <i>nemoralis</i>	Bristly Phacelia	Boragin-	- - B	- NBP	Moist slopes, streambanks, mixed-evergreen forest (Apr–Jul)	
<input type="radio"/> <i>Piperia elegans</i> subsp. <i>elegans</i> (JM93: no subsp.)	Elegant Rein Orchid	Orchid-	- - B	- NP	Generally dry, open sites, scrub, conifer forest (May–Sep)	
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Boragin-	- - -	- NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	- - -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	- - -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	- - -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	- - -	- NA	Open grassland, sandy soil, burns (Mar–May)	
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	- - -	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	- - -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Polygala californica</i>	California Milkwort	Polygal-	- - A1	- NP	Coastal prairie and forest, chaparral, occasionally on serpentine (Apr–Jul)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i>)	Knotweed	Polygon-	- - -	- ZA	Disturbed places (May–Nov)	
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i>)	Hooker Fairy Bells	Lili-	- - C	- NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)	

Herbaceous <i>Pseudognaphalium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Pseudognaphalium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i>)	California Everlasting	Aster-	- - -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)
<input type="radio"/> <i>Pseudognaphalium microcephalum</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>microcephalum</i>)	White Everlasting	Aster-	- - B	- NB	Grassy hillsides, gravelly canyon bottoms, chaparral, coastal-sage scrub (Jun–Aug)
<input type="radio"/> <i>Pseudognaphalium ramosissimum</i> (JM93: <i>Gnaphalium</i>)	Pink Everlasting	Aster-	- - -	- NB	Dry, open slopes, woodland, sandy fields, dunes (Jul–Sep)
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i>)	Cotton-batting Plant	Aster-	- - -	- NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)
<input type="radio"/> <i>Psilocarphus tenellus</i> (JM93: var. <i>tenellus</i>)	Slender Woolly-marbles	Aster-	- - -	- NA	Common. Dry, seasonally moist slopes, flats, burns, trails, rarely vernal pools (Mar–Jul)
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	- - -	- NA	Common. Sand or gravel (Mar–Jul)
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	- - -	- NP	Grassland, open woodland (Mar–Aug)
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	- - -	- NA	Grassland, open woodland (Mar–May)
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	- - -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	- - -	L ZA	Disturbed areas, fields (May–Jul)
<input type="radio"/> <i>Rigiopappus leptocladus</i>	Bristle-head	Aster-	- - B	- NA	Grassy sites, openings among shrubs or trees (Apr–Jul)
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	- - -	M ZP	± Disturbed, often acidic places (Apr–Jul)
<input type="radio"/> <i>Rumex conglomeratus</i>	Green Dock	Polygon-	- - -	- ZP	Common. Moist places (May–Aug)
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	- - -	L ZP	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Rumex obtusifolius</i>	Bitter Dock	Polygon-	- - -	- ZP	Moist places (May–Sep)
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	- - -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)
<input type="radio"/> <i>Rumex transitorius</i> (JM93: <i>R. salicifolius</i> var. <i>transitorius</i>)	Ovate Willow Dock	Polygon-	- - -	- NP	Coastal dunes, wet margins of meadows (Apr–Jun)
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	- - -	- NP	Woodland (May–Sep)
<input type="radio"/> <i>Sagina decumbens</i> subsp. <i>occidentalis</i>	Western Pearlwort	Caryophyll-	- - C	- NA	Dry streams, chaparral, grassy areas, rock outcrops, vernal pools (Spring–early summer)
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	- - -	- NP	Open grassland or pine/oak woodland (Apr–May)
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	- - -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	- - -	- NP	Open slopes, ravines, woodland (Mar–May)
<input type="radio"/> <i>Sanicula laciniata</i>	Fringe-leaf / Coast Sanicle	Api-	- - A2	- NP	Coastal, open or shrubby slopes, woodland (Mar–May)

Herbaceous <i>Sanicula</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Sanicula</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Sanicula tuberosa</i>	Turkey Pea Sanicle	Api-	- - C	- NP	Open gravelly meadows, chaparral, woodland, pine forest (Mar–Jul)
<input type="radio"/> <i>Scabiosa atropurpurea</i>	Pincushion Flower	Dipsac-	- - -	- ZA	Disturbed areas (Mar–Nov)
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	- - -	- ZA	Grassy slopes, roadsides (Apr–Jun)
<input type="radio"/> <i>Scrophularia californica</i> (JM93: <i>subsp. californica, floribunda</i>)	California Figwort	Scrophulari-	- - -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	- - C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	- - -	- ZA	Common. Disturbed areas (Feb–Jul)
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	- - -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)
<input type="radio"/> <i>Sidalcea malviflora subsp. ?</i>	Common Checkerbloom	Malv-	- - -	- NP	- -
<input type="radio"/> <i>Sidalcea malviflora subsp. laciniata</i>	Geranium-leaved Checkerbloom	Malv-	- - C	- NP	Grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	- - -	- ZA	Fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Silene laciniata subsp. californica</i> (JM93: <i>S. californica</i>)	California Pink	Caryophyll-	- - B	- NP	Chaparral, oak woodland, conifer forest, serpentine or not (Spring–summer)
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	- - -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	- - -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	- - -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)
<input type="radio"/> <i>Sisyrinchium californicum</i>	Golden-eyed-grass	Irid-	- - A1	- NP	Gen moist places near coast (Spring–late summer)
<input type="radio"/> <i>Solanum americanum</i>	Small-flowered Nightshade	Solan-	- - -	- NAP	Open, often disturbed places (Apr–Nov)
<input type="radio"/> <i>Solidago elongata</i> (JM93: <i>S. canadensis subsp. elongata</i>)	West Coast Canada Goldenrod	Aster-	- - -	- NP	Meadows, thickets (May–Oct)
<input type="radio"/> <i>Solidago velutina subsp. californica</i> (JM93: <i>S. californica</i>)	California Goldenrod	Aster-	- - -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	- - -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)
<input type="radio"/> <i>Sonchus asper subsp. asper</i>	Prickly Sow Thistle	Aster-	- - -	- ZA	Common. Slightly moist disturbed sites, along streams (All year)
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	- - -	- ZA	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Spergula arvensis</i> (JM93: <i>subsp. arvensis</i>)	Stickwort	Caryophyll-	- - -	- ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)
<input type="radio"/> <i>Spergularia bocconi</i>	Boccone's Sand-spurry	Caryophyll-	- - -	- ZA	Salt marshes, alkaline areas, sandy soils (Spring)
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	- - -	- ZAP	Forest, meadows, mud flats, disturbed

Herbaceous <i>Stachys</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Stachys</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i>)	Common Rigid Hedge-nettle	Lami-	- - -	- NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	- - -	- ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)
<input type="radio"/> <i>Stephanomeria virgata</i> subsp. <i>pleurocarpa</i>	Twiggy Wreath Plant	Aster-	- - C	- NA	Chaparral openings, grassland (Jun–Nov)
<input type="radio"/> <i>Symphotrichum chilense</i> (JM93: <i>Aster chilensis</i>)	Common California Wild Aster	Aster-	- - -	- NP	Grassland, salt marshes, disturbed places (Jun–Oct)
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	- - -	- ZBP	Abundant. Esp disturbed areas (All year)
<input type="radio"/> <i>Taraxia ovata</i> (JM93: <i>Camissonia</i>)	Golden Eggs Suncup	Onagr-	- - C	- NP	Grassy fields, gen clay soil (Mar–Jun)
<input type="radio"/> <i>Tellima grandiflora</i>	Fringe Cups	Saxifrag-	- - B	- NP	Moist slopes (Apr–Jul)
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	- - -	- NP	Moist, open to shaded places, woodland, forest (Mar–Jun)
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	- - -	- NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)
<input type="radio"/> <i>Tolmiea diplomenziesii</i> (JM93: <i>T. menziesii</i>)	Pig-a-back Plant	Saxifrag-	- - A1	- NP	Moist streambanks (May–Aug)
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	- - -	M ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	- - -	- ZA	Disturbed places (Apr–Jun)
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i>)	Common Star Lily	Melanthi-	- - -	- NP	Grassy or wooded slopes, outcrops (Feb–Jun)
<input type="radio"/> <i>Tradescantia fluminensis</i>	Spiderwort	Commelin-	- - -	- ZP	Shaded woodland, streambanks (Apr–Jul)
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	- - -	- ZBP	Common. Disturbed places (Mar–Nov)
<input type="radio"/> <i>Trientalis latifolia</i>	Starflower	Myrsin-	- - C	- NP	Shaded places, esp woodland (Apr–Jul)
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Fab-	- - -	- NA	Open, grassy areas, forest (Apr–Jun)
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	- - -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Fab-	- - -	- NA	Grassland, coastal woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Fab-	- - -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	- - -	- ZA	Agricultural, disturbed areas, lawns (Spring)
<input type="radio"/> <i>Trifolium fucatum</i>	Bull Clover	Fab-	- - C	- NA	Locally abundant. Moist, open grassland, marshes, roadsides, occasionally saline or serpentine soils (Apr–Jun)
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i>)	Pinpoint Clover	Fab-	- - -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	- - -	M ZA	Disturbed areas, roadsides (Apr–May)
<input type="radio"/> <i>Trifolium incarnatum</i>	Crimson Clover	Fab-	- - -	- ZA	Uncommon. Disturbed areas (May–Aug)
<input type="radio"/> <i>Trifolium macraei</i>	Macrae's Clover	Fab-	- - A2	- NA	Disturbed areas, dunes (Mar–May)

Herbaceous <i>Trifolium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Trifolium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Fab-	- - -	- NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium sp.</i>	Clover	Fab-	- - -	- -		
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	- - -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	- - -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)	
<input type="radio"/> <i>Trillium ovatum subsp. ovatum</i>	Western White Trillium	Melanthi-	- - A2	- NP	Redwood, mixed-evergreen forest on moist wooded slopes (Feb–Apr)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	- - -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Themid-	- - -	- NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	- - -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Brassic-	- - B	- NP	Grassy banks, open fields, roadsides, pastures (Mar–May)	
<input type="radio"/> <i>Turritis glabra (JM93: Arabis, var. glabra)</i>	Tower Mustard	Brassic-	- - C	- NP	Open fields, meadows, slopes (Apr–Jul)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	- - -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urtica dioica subsp. holosericea</i>	Hoary Nettle	Urtic-	- - -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	- - -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbena lasiostachys var. lasiostachys</i>	Western Vervain	Verben-	- - -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	- - -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	- - -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia gigantea</i>	Giant Vetch	Fab-	- - C	- NP	Coastal shrub, coastal forest, chaparral (Mar–Aug)	
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	- - -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	- - -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	- - -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)	
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Apocyn-	- - -	M ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))	

Herbaceous <i>Viola</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Viola</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Viola adunca</i> var. <i>adunca</i> (JM93: no var.)	Western Dog Violet	Viol-	- - A1	- NP	Vernally moist meadows, damp streambanks, meadow edges in conifer forest, generally shade (Apr–Aug)
<input type="radio"/> <i>Viola glabella</i>	Stream / Smooth Yellow Violet	Viol-	- - A2	- NP	Moist to wet generally shady places in forest, streambanks, etc (Mar–Aug)
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	- - C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)
<input type="radio"/> <i>Viola sempervirens</i>	Redwood / Evergreen Violet	Viol-	- - A1	- NP	Shady areas in coastal forest (Jan–Jul)
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	- - -	- NP	Grassland (Apr–Aug)
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	- - C	- NP	Gen shady sites (Mar–Jun)
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	- - -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))
<input type="radio"/> <i>Zantedeschia aethiopica</i>	Calla-lily	Ar-	- - -	L ZP	Disturbed areas, near former habitations (Mar–Jun)
<input type="radio"/> <i>Zeltnera muehlenbergii</i> (JM93: <i>Centaurium</i>)	Monterey Centaury	Gentian-	- - -	- NA	Moist coastal bluffs, forest openings (Jun–Aug)

Woody Acacia	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Woody Acacia
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
○ <i>Acacia dealbata</i>	Silver Wattle	Fab-	- - -	M ZP	Locally common. Disturbed areas, often roadsides (Feb–Apr)	
○ <i>Acacia melanoxylon</i>	Blackwood Acacia	Fab-	- - -	L ZP	Uncommon. Disturbed areas (Feb–Mar)	
○ <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	- - -	- NP	Common. Streambanks, canyons (Mar–Jun)	
○ <i>Acer negundo</i> (JM93: var. <i>californicum</i>)	Box Elder	Sapind-	- - -	- NP	Streamsides, bottomland. (Mar–Apr)	
○ <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	- - -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
○ <i>Aesculus californica</i>	California Buckeye	Sapind-	- - -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
○ <i>Alnus rhombifolia</i>	White Alder	Betul-	- - -	- NP	Along permanent streams (Apr–Jun)	
○ <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	- - C	- NP	Conifer, oak forests (Mar–May)	
○ <i>Arctostaphylos crustacea</i> subsp. <i>crustacea</i> (JM93: <i>A. tomentosa</i> subsp. <i>crustacea</i>)	Brittle-leaf Manzanita	Eric-	- - C	- NP	Chaparral, conifer forest (Feb–Apr)	
○ <i>Arctostaphylos glandulosa</i> subsp. <i>glandulosa</i>	Eastwood Manzanita	Eric-	- - B	- NP	Chaparral, conifer forest (Jan–Apr)	
○ <i>Arctostaphylos pallida</i>	Pallid Manzanita	Eric-	FT/CE 1B.1 *A1	- NP	Siliceous shales, slopes, ridges, chaparral (Jan–Mar)	
○ <i>Artemisia californica</i>	California Sagebrush	Aster-	- - -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
○ <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	- - -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
○ <i>Berberis pinnata</i> subsp. <i>pinnata</i>	Shinyleaf Oregon-grape	Berberid-	- - C	- NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
○ <i>Buddleja davidii</i>	Butterfly Bush	Scrophulari-	- - -	- ZP	Disturbed, generally ± damp soil (May–Sep)	
○ <i>Ceanothus oliganthus</i> var. <i>sorediatus</i>	Jim Brush	Rhamn-	- - -	- NP	Slopes, ridges, chaparral, conifer forest (Jan–May)	
○ <i>Ceanothus thrysiflorus</i> var. <i>thrysiflorus</i> (JM93: no var.)	Blue Blossom	Rhamn-	- - -	- NP	Bluffs, slopes, canyons, chaparral, coastal scrub, closed-cone-pine forest (Mar–Jun)	
○ <i>Chrysolepis chrysophylla</i> var. <i>minor</i>	Golden Chinquapin	Fag-	- - A2	- NP	Conifer forest, closed-cone-pine forest, chaparral (Jun–Sep)	
○ <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	- - -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
○ <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	- - C	- NP	Along streams, wet places (Jun–Sep)	
○ <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Corn-	- - C	- NP	Generally moist places (May–Jul)	
○ <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Corn-	- - C	- NP	Generally moist habitats (May–Jul)	
○ <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i>)	California Hazelnut	Betul-	- - -	- NP	Common. Many habitats, esp moist, shady places (Jan–Mar)	
○ <i>Cotoneaster pannosus</i> (JM93: <i>C. pannosa</i>)	Silverleaf Cotoneaster	Ros-	- - -	M ZP	Disturbed places, mixed-evergreen forest (May–Jul)	
○ <i>Crataegus monogyna</i>	Hawthorn	Ros-	- - -	L ZP	Naturalized in scattered places (May–Aug)	

Woody	Grouped by Growth Form						Woody
Dirca	Sorted Alphabetically by Scientific Name						Dirca
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Dirca occidentalis</i>	Western Leatherwood	Thymelae-	- 1B.2 *A2	-	NP	Gen n facing slopes, mixed-evergreen forest to chaparral, gen fog belt (Nov–Mar)	
<input type="radio"/> <i>Ericameria arborescens</i>	Golden-fleece	Aster-	- - A2	-	NP	Woodland, open forest, chaparral, especially after fire (Aug–Nov)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>foliolosum</i>	Leafy California Buckwheat	Polygon-	- - A2	-	NP	Gravel (All year)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	- - -	-	NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Eucalyptus camaldulensis</i>	River Red Gum	Myrt-	- - -	L	ZP	Common. Disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	- - -	M	ZP	Common. Disturbed areas (Oct–Jan)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i>)	California Coffee Berry	Rhamn-	- - -	-	NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Garrya elliptica</i>	Coast Silk Tassel	Garry-	- - B	-	NP	Seacliffs, sand dunes, chaparral, foothill-pine woodland (Jan–Mar)	
<input type="radio"/> <i>Gaultheria shallon</i>	Salal	Eric-	- - A1	-	NP	Moist forest margins (Apr–Jul)	
<input type="radio"/> <i>Genista monspessulana</i>	French Broom	Fab-	- - -	H	ZP	Common. Disturbed places. (Mar–Jun)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	- - C	-	NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Hedera helix</i> (JM93: subsp. <i>helix</i>)	English Ivy	Arali-	- - -	H	ZP	Woodland, open, disturbed areas (Aug–Nov)	
<input type="radio"/> <i>Hesperocyparis macrocarpa</i> (JM93: <i>Cupressus</i>)	Monterey Cypress	Cupress-	- - -	N	NP	Closed-cone-pine/cypress forests -	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	- - -	-	NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	- - -	-	NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Ilex aquifolium</i>	English Holly	Aquifoli-	- - -	M	ZP	Cool, wooded areas (May-Jun)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i>)	N. California Black Walnut	Jugland-	- 1B.1 *A2	-	NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Caprifoli-	- - -	-	NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera involucrata</i> var. <i>ledebourii</i>	Black Twinberry Honeysuckle	Caprifoli-	- - B	-	NP	Moist places (May–Jul)	
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	- - -	-	NP	--	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>albifrons</i>	Bush Lupine	Fab-	- - -	-	NP	Common. Chaparral, foothill woodland (Mar–Jun)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	- - -	-	NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Morella californica</i> (JM93: <i>Myrica</i>)	California Bayberry	Myric-	- - A2	-	NP	Coastal dunes and scrub, closed-cone pine, redwood forest (Mar–Apr)	

Woody	Grouped by Growth Form					Woody
Oemleria	Sorted Alphabetically by Scientific Name					Oemleria
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	- - -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Physocarpus capitatus</i>	Pacific Ninebark	Ros-	- - C	- NP	Moist banks, n-facing slopes, mixed-conifer forest (May–Jul)	
<input type="radio"/> <i>Pinus attenuata</i>	Knobcone Pine	Pin-	- - A2	- NP	Closed-cone-pine forest, chaparral -	
<input type="radio"/> <i>Pinus coulteri</i>	Coulter Pine	Pin-	- - A2	- NP	Chaparral, lower mixed-conifer, mixed-hardwood forests -	
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Pin-	- - -	- NP	Closed-cone-pine forest, oak woodland -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	- - B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Populus fremontii subsp. fremontii</i>	Fremont Cottonwood	Salic-	- - -	- NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)	
<input type="radio"/> <i>Populus trichocarpa (JM93: P. balsamifera subsp. trichocarpa)</i>	Black Cottonwood	Salic-	- - C	- NP	Alluvial bottomland, streamsides (Feb–Apr)	
<input type="radio"/> <i>Prunus cerasifera</i>	Cherry Plum	Ros-	- - -	L ZP	Roadsides, streambanks, chaparral as waif (Feb–Mar)	
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Ros-	- - C	- NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Prunus ilicifolia subsp. ilicifolia</i>	Holly-leaved Cherry	Ros-	- - C	- NP	Canyons, slopes, scrubland, woodland (Apr–May)	
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Ros-	- - C	- NP	Mixed-evergreen or conifer forest (Mar–May)	
<input type="radio"/> <i>Pyracantha angustifolia</i>	Woolly Firethorn	Ros-	- - -	L ZP	Disturbed areas, fencerows, abandoned fields, roadsides (Feb–Jun)	
<input type="radio"/> <i>Quercus agrifolia var. agrifolia</i>	Coast Live Oak	Fag-	- - A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	- - -	- NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus wislizeni var. frutescens</i>	Dwarf Interior Live Oak	Fag-	- - C	- NP	Valleys, chaparral (Mar–May)	
<input type="radio"/> <i>Quercus wislizeni var. wislizeni</i>	Interior Live Oak	Fag-	- - -	- NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Ribes californicum var. californicum</i>	Hillside Gooseberry	Grossulari-	- - -	- NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes divaricatum var. pubiflorum</i>	Straggle Gooseberry	Grossulari-	- - B	- NP	Uncommon. Coastal bluffs, forest edges (Mar–May)	
<input type="radio"/> <i>Ribes menziesii var. menziesii (JM93: no var.)</i>	Canyon Gooseberry	Grossulari-	- - -	- NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes sanguineum var. glutinosum</i>	Red-flowering Currant	Grossulari-	- - C	- NP	Many habitats (Feb–Apr)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Fab-	- - -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	

Woody	Grouped by Growth Form						Woody
Rosa	Sorted Alphabetically by Scientific Name						Rosa
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)	
<input type="checkbox"/> <i>Rosa californica</i>	California Rose	Ros-	- - -	-	NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="checkbox"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	- - -	-	NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="checkbox"/> <i>Rosa spithamea</i>	Coast Ground Rose	Ros-	- - C	-	NP	Open forest, chaparral, especially after fire (Apr–Aug)	
<input type="checkbox"/> <i>Rubus armeniicus</i> (JM93: <i>R. discolor</i>)	Himalayan Blackberry	Ros-	- - -	H	ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="checkbox"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	- - -	-	NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="checkbox"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	- - -	-	NP	Open, disturbed areas (Mar–Jul)	
<input type="checkbox"/> <i>Salix laevigata</i>	Red Willow	Salic-	- - -	-	NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)	
<input type="checkbox"/> <i>Salix lasiandra</i> var. <i>lasiandra</i> (JM93: <i>S. lucida</i> subsp. <i>Lasiandra</i>)	Pacific Willow	Salic-	- - -	-	NP	Wet meadows, lakeshores, riverbanks (May–Jun)	
<input type="checkbox"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	- - -	-	NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="checkbox"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i>)	Blue Elderberry	Adox-	- - -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="checkbox"/> <i>Sambucus racemosa</i> var. <i>racemosa</i>	Red Elderberry	Adox-	- - A1	-	NP	Moist places (May–Jul)	
<input type="checkbox"/> <i>Sequoia sempervirens</i>	Redwood	Cupress-	- - B	-	NP	Redwood forest -	
<input type="checkbox"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	- - -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="checkbox"/> <i>Spartium junceum</i>	Spanish Broom	Fab-	- - -	H	ZP	Disturbed areas (Apr–Jun)	
<input type="checkbox"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	- - -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="checkbox"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	- - -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="checkbox"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	- - -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="checkbox"/> <i>Umbellularia californica</i>	California Bay	Laur-	- - -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)	
<input type="checkbox"/> <i>Vaccinium ovatum</i>	California Huckleberry	Eric-	- - B	-	NP	Edges, clearings in conifer forest (Mar–May)	
<input type="checkbox"/> <i>Vitis californica</i>	California Wild Grape	Vit-	- - -	-	NP	Streamsides, springs, canyons (May–Jun)	

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex deweyana ssp. *leptopoda* (*C. leptopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thyrsoiflorus (var. *thyrsoiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria* *amphibia*)

Polygonum amphibium var. *stipulaceum* (*Persicaria* *amphibia*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)