

# Sunol Regional Wilderness

## Checklist of Wild Plants

### Sorted Alphabetically by Growth Form, Scientific Name



This is a comprehensive list of the wild plants reported to be found in Sunol Regional Wilderness. The plants are sorted alphabetically by growth form, then by scientific name. This list includes the common name, family, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix.

Column Heading	Description
✓	Checklist column for marking off the plants you observe
<b>Scientific Name</b>	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> ( <a href="http://ucjeps.berkeley.edu/IJM.html">ucjeps.berkeley.edu/IJM.html</a> )
<b>(JM93 if different)</b>	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
<b>Common Name</b>	According to <i>JM2</i> and other references (not standardized)
<b>Family</b>	Scientific family name according to <i>JM2</i> , abbreviated by replacing the “aceae” ending with “-” (ie. Asteraceae = Aster-)
<b>Status</b>	Special status rating (if any), listed in 3 categories, divided by vertical bars (‘ ’): <b>Federal/California (Fed./Calif.)   California Native Plant Society (CNPS)   East Bay chapter of the CNPS (EBCNPS)</b>  <b>Fed./Calif.:</b> FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare  <b>CNPS</b> (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California  <b>EBCNPS</b> (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
<b>Inv</b>	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
<b>OL</b>	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
<b>Habitat (Bloom)</b>	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, [wlegard@ebparks.org](mailto:wlegard@ebparks.org)). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

# Sunol Regional Wilderness Checklist of Wild Plants

Ferns & Horsetails	Grouped by Growth Form					Ferns & Horsetails
<i>Adiantum</i>	Sorted Alphabetically by Scientific Name					<i>Adiantum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Pterid-	-   -   -	- NP	Shaded hillsides, moist woodland -	
<input type="radio"/> <i>Azolla filiculoides</i>	Mosquito Fern	Azoll-	-   -   -	- NP	Common. Ponds, slow streams -	
<input type="radio"/> <i>Cheilanthes covillei</i>	Coville's Lip Fern	Pterid-	-   -   A2	- NP	Crevices, bases of rocks, sun or shade -	
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Dryopterid-	-   -   -	- NP	Locally common. Open, wooded slopes, caves -	
<input type="radio"/> <i>Equisetum arvense</i>	Common Horsetail	Equiset-	-   -   -	- NP	Streambanks, wet meadows, springs, other wet, shaded places -	
<input type="radio"/> <i>Equisetum hyemale subsp. affine</i>	Common Scouring Rush	Equiset-	-   -   -	- NP	Streams, moist, sandy, gravelly areas -	
<input type="radio"/> <i>Equisetum laevigatum</i>	Smooth Scouring Rush	Equiset-	-   -   B	- NP	Moist, sandy or gravelly areas -	
<input type="radio"/> <i>Equisetum telmateia subsp. braunii</i>	Giant Horsetail	Equiset-	-   -   -	- NP	Streambanks, roadside ditches, seepage areas -	
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Pterid-	-   -   -	- NP	Generally rocky or dry areas -	
<input type="radio"/> <i>Pellaea mucronata var. mucronata</i>	Bird's-foot Fern	Pterid-	-   -   C	- NP	Rocky or dry areas -	
<input type="radio"/> <i>Pentagramma triangularis subsp. triangularis</i>	Goldenback Fern	Pterid-	-   -   -	- NP	Gen shaded, sometimes rocky or wooded areas -	
<input type="radio"/> <i>Polypodium calirhiza</i>	Polypody Fern	Polypodi-	-   -   -	- NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="radio"/> <i>Pteridium aquilinum var. pubescens</i>	Bracken Fern	Dennstaedti-	-   -   -	- NP	Pastures, woodland, meadows, hillsides, partial to full sun -	
<input type="radio"/> <i>Selaginella bigelovii</i>	Spike-moss	Selaginell-	-   -   A2	- NP	Open sites, mineral soil on rock outcrops, amid shrubs -	
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Blechn-	-   -   -	- NP	Near streams, springs, seeps -	

Grasses & Grass-like <i>Agrostis</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Agrostis</i>	
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Agrostis avenacea</i>	Pacific Bent Grass	Po-	-   -   -	L	ZP	Open, often disturbed place (Jun-Jul)
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Po-	-   -   -	-	NP	Common. Open meadows, woodland, forest, subalpine (Jun-Aug)
<input type="radio"/> <i>Aira caryophylla</i>	Silver Hair Grass	Po-	-   -   -	-	ZA	Sandy soils, open or disturbed sites (Apr-Jun)
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Po-	-   -   -	M	ZA	Disturbed sites (Mar-Jun)
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Po-	-   -   -	M	ZA	Disturbed sites (Apr-Jun)
<input type="radio"/> <i>Brachypodium distachyon</i>	Purple False-brome	Po-	-   -   -	M	ZA	Disturbed areas, dry slopes (Apr-Jul)
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Po-	-   -   -	-	ZA	Shaded or moist, open sites (Apr-Jul)
<input type="radio"/> <i>Bromus carinatus</i> var. <i>carinatus</i>	California Brome	Po-	-   -   -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Po-	-   -   -	M	ZA	Open, gen disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Po-	-   -   -	L	ZA	Fields, disturbed areas (Apr-Jul)
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Po-	-   -   -	-	NP	Shrubland, conifer forest, shaded streambanks, roadsides (May-Jul)
<input type="radio"/> <i>Bromus madritensis</i> subsp. <i>rubens</i>	Red Brome	Po-	-   -   -	H	ZA	Disturbed areas, roadsides (Mar-Jun)
<input type="radio"/> <i>Bromus sterilis</i>	Poverty Brome	Po-	-   -   -	-	ZA	Open, disturbed areas (Mar-Jun)
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Cyper-	-   -   B	-	NP	Seasonally wet places (May-Aug)
<input type="radio"/> <i>Carex nebrascensis</i>	Nebraska Sedge	Cyper-	-   -   B	-	NP	Meadows, swamps (Jun-Sep)
<input type="radio"/> <i>Carex nudata</i>	Torrent Sedge	Cyper-	-   -   A2	-	NP	Rocky or sandy streambeds below high-water mark (Apr-Jul)
<input type="radio"/> <i>Carex</i> sp.	Sedge	Cyper-	-   -   -	-	-	-
<input type="radio"/> <i>Crypsis schoenoides</i>	Swamp Prickle Grass	Po-	-   -   -	-	ZA	Wet places (Jun-Oct)
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Po-	-   -   -	M	ZP	Disturbed sites (Jun-Aug)
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Po-	-   -   -	M	ZA	Open, disturbed sites (May-Jul)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Cyper-	-   -   -	-	NP	Vernal pools, streambanks (May-Nov)
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Po-	-   -   -	L	ZP	Disturbed, often moist sites (May-Aug)
<input type="radio"/> <i>Echinochloa crus-galli</i>	Barnyard Grass	Po-	-   -   -	-	ZA	Gen wet, disturbed sites, fields, roadsides (Jun-Oct)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Cyper-	-   -   -	-	NP	Common. Fresh to brackish wetland (Spring-summer)
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Po-	-   -   -	-	NP	Open areas, chaparral, woodland, forest (Jun-Aug)
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Po-	-   -   C	-	NP	Open, sandy to rocky areas (May-Jul)
<input type="radio"/> <i>Elymus stebbinsii</i>	Parish's Wheat Grass	Po-	-   -   A2	-	NP	Dry slopes, chaparral, conifer forest (Jun-Jul)
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i> )	Beardless Wild Rye	Po-	-   -   -	-	NP	Dry to moist, often saline, meadows (Jun-Jul)
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Po-	-   -   -	M	ZP	Disturbed places (May-Jun)

Grasses & Grass-like <i>Festuca</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Festuca</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i> )	Brome Fescue	Po-	-   -   -	- ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Po-	-   -   C	- NP	Dry, open forest, moist streambanks, chaparral (May–Jun)
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Po-	-   -   C	- NP	Dry, open or shady places (Jul–Sep)
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i> )	Hairy Fescue	Po-	-   -   -	- NA	Disturbed, open, gen sandy soils (Apr–Jun)
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i> )	Rattail Sixweeks Grass	Po-	-   -   -	- ZA	Common. Generally open places, sandy soils (Feb–May)
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i> )	Rye Grass	Po-	-   -   -	- ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Po-	-   -   -	- NA	--
<input type="radio"/> <i>Festuca</i> sp. (perennial)	Perennial Fescue	Po-	-   -   -	-	--
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>californicum</i>	California Barley	Po-	-   -   B	- NP	Meadows, pastures, streambanks (Mar–Jul)
<input type="radio"/> <i>Hordeum jubatum</i> subsp. <i>jubatum</i> (JM93: no subsp.)	Foxtail Barley	Po-	-   -   B	- NAP	Roadsides, disturbed areas, meadows, marshes (May–Jul)
<input type="radio"/> <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Po-	-   -   -	- ZA	Dry to moist, disturbed sites (Apr–Jun)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites. Common (Feb–May)
<input type="radio"/> <i>Hordeum murinum</i> subsp. <i>murinum</i>	Wall Barley	Po-	-   -   -	- ZA	Moist, gen disturbed sites (Feb–May)
<input type="radio"/> <i>Juncus balticus</i> subsp. <i>ater</i> (JM93: no subsp.)	Baltic Rush	Junc-	-   -   -	- NP	Moist to ± dry sites (Jul–Nov)
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Junc-	-   -   -	- NP	Marshy places, creeks, seeps (Jun–Oct)
<input type="radio"/> <i>Juncus phaeocephalus</i> var. <i>paniculatus</i>	Panicled Brownhead Rush	Junc-	-   -   C	- NP	Wet places, coastal, inland (Jun–Sep)
<input type="radio"/> <i>Juncus phaeocephalus</i> var. <i>phaeocephalus</i>	Brownheaded Rush	Junc-	-   -   B	- NP	Coastal meadows, dune hollows, marsh edges (Jun–Aug)
<input type="radio"/> <i>Juncus</i> sp.	Rush	Junc-	-   -   -	-	--
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Junc-	-   -   -	- NP	Wet places (Jul–Oct)
<input type="radio"/> <i>Koeleria gerardii</i> (JM93: var. <i>hindsii</i> )	Annual June Grass	Po-	-   -   -	- ZA	Open, disturbed sites (Apr–Jul)
<input type="radio"/> <i>Koeleria macrantha</i>	June Grass	Po-	-   -   C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)
<input type="radio"/> <i>Lamarckia aurea</i>	Goldentop	Po-	-   -   -	- ZA	Open ground, moist seeps, rocky hillsides, sandy soil (Feb–May)
<input type="radio"/> <i>Leptochloa fusca</i> subsp. <i>fascicularis</i> (JM93: <i>L. fascicularis</i> )	Bearded Sprangletop	Po-	-   -   A2	- NA	Marshes, wetlands, often wet disturbed areas, often associated with agriculture (Jun–Oct)
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Junc-	-   -   -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)

Grasses & Grass-like <i>Melica</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Grasses & Grass-like <i>Melica</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Melica bulbosa</i>	Oniongrass	Po-	-   -   A1?	- NP	Dry rocky slopes, conifer forest (Jul–Aug)
<input type="radio"/> <i>Melica californica</i>	California Melic	Po-	-   -   C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Melica imperfecta</i>	Little California Melica	Po-	-   -   -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)
<input type="radio"/> <i>Melica torreyana</i>	Torrey's Melic	Po-	-   -   -	- NP	Chaparral, conifer forest (Mar–Jun)
<input type="radio"/> <i>Paspalum dilatatum</i>	Dallis Grass	Po-	-   -   -	- ZP	Disturbed areas (May–Nov)
<input type="radio"/> <i>Paspalum distichum</i>	Knot Grass	Po-	-   -   -	- ZP	Edges of lakes, ponds, rice fields, wet roadside ditches (Jun–Oct)
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Po-	-   -   -	- ZA	Abundant. Disturbed moist ground (Feb–Sep)
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Po-	-   -   -	- NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Po-	-   -   -	- ZP	Common. Streambanks (May–Aug)
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Po-	-   -   -	L ZA	Moist places, along streams (Apr–Aug)
<input type="radio"/> <i>Polypogon viridis</i> (JM93: <i>Agrostis</i> )	Water Beard Grass	Po-	-   -   -	- ZP	Common. Disturbed areas, wet areas, ponds, streambanks (May–Jun)
<input type="radio"/> <i>Stipa cernua</i> (JM93: <i>Nassella</i> )	Nodding Needle Grass	Po-	-   -   C	- NP	Grassland, chaparral, juniper woodland (Feb–Jul)
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i> )	Foothill Needle Grass	Po-	-   -   C	- NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)
<input type="radio"/> <i>Stipa miliacea</i> var. <i>miliacea</i> (JM93: <i>Piptatherum miliaceum</i> )	Smilo Grass	Po-	-   -   -	L ZP	Salt marshes, streambanks, chaparral, open woodland, disturbed (Mar–Sep)
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i> )	Purple Needle Grass	Po-	-   -   C	- NP	Oak woodland, chaparral, grassland (Mar–Jun)
<input type="radio"/> <i>Triglochin scilloides</i> (JM93: <i>Lilaea</i> )	Flowering-quillwort	Juncagin-	-   -   C	- NA	Vernal pools, streams, ponds, lake margins (Mar–Oct)
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Typh-	-   -   -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Typh-	-   -   -	- NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)

Herbaceous <i>Acanthomintha</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Acanthomintha</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acanthomintha lanceolata</i>	Santa Clara Thornmint	Lami-	-   4.2   *A2	- NP	Arid, rocky slopes, openings in scrub or woodland, gen on serpentine (Mar–Jun)	
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Aster-	-   -   -	- NP	Many habitats (Apr–Sep)	
<input type="radio"/> <i>Achyrrachaena mollis</i>	Blow Wives	Aster-	-   -   -	- NA	Common. Grassy sites, often clay soils (Mar–Jun)	
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i> )	Spanish Clover	Fab-	-   -   -	- NA		
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i> )	Colchita	Fab-	-   -   -	- NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i> )	Deerweed	Fab-	-   -   -	- NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon</i> or <i>Hosackia</i> sp. (JM93: <i>Lotus</i> sp.)	Lotus	Fab-	-   -   -	- -		
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i> )	Small-flower Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Acmispon strigosus</i> (JM93: <i>Lotus</i> )	Hairy Lotus	Fab-	-   -   A2	- NP	Coastal scrub, chaparral, foothills, deserts, roadsides, other disturbed area (Mar–Jun)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i> )	California Lotus	Fab-	-   -   -	- NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Aster-	-   -   -	- NP	Grassland, scrub, woodland (Apr–Jul)	
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Aster-	-   -   -	- NA	Many open habitats (May–Jun)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Alli-	-   -   -	- NP	Common. Grassy slopes (Apr–May)	
<input type="radio"/> <i>Amaranthus albus</i>	Tumbleweed	Amaranth-	-   -   -	- ZA	Disturbed areas, roadsides, riverbanks, sandy places, agricultural fields (Jun–Oct)	
<input type="radio"/> <i>Amaranthus blitoides</i>	Procumbent Pigweed	Amaranth-	-   -   C	- NA	Disturbed areas, roadsides, agricultural fields, sandy soil (Jul–Nov)	
<input type="radio"/> <i>Amaranthus</i> sp.	Amaranth	Amaranth-	-   -   -	- -		
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i> )	Common Fiddleneck	Boragin-	-   -   -	- NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Amsinckia menziesii</i> (JM93: var. <i>menziesii</i> )	Small-flowered Fiddleneck	Boragin-	-   -   -	- NA	Shade-tolerant, open, disturbed areas at forest/woodland edges (May–Jul)	
<input type="radio"/> <i>Amsinckia tessellata</i> var. <i>tessellata</i>	Desert Fiddleneck	Boragin-	-   -   B	- NA	Rocky or sandy soils (Feb–Jun)	
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Myrsin-	-   -   -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Anaphalis margaritacea</i>	Pearly Everlasting	Aster-	-   -   -	- NP	Woodland, disturbed places (Jul–Oct)	

Herbaceous <i>Ancistrocarphus</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Ancistrocarphus</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Ancistrocarphus filagineus</i>	Woolly Fishhooks	Aster-	-   -   B	- NA	Bare or grassy, often serpentine or clay, drainages, road beds, burns, vernal moist sites (Mar–Jun)
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Api-	-   -   -	- NP	Generally wooded areas (Jun–Aug)
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Aster-	-   -   -	- ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Api-	-   -   -	- ZA	Generally shady places (Apr–Jun)
<input type="radio"/> <i>Antirrhinum multiflorum</i>	Mayflower Snapdragon	Plantagin-	-   -   A1	- NA	Rocky or disturbed areas, burns (Apr–Aug)
<input type="radio"/> <i>Antirrhinum vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i> (JM93: <i>A. vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i> )	Wiry Snapdragon	Plantagin-	-   -   B	- NA	Disturbed areas (Jun–Aug)
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Ros-	-   -   -	- NA	Seasonally moist grassland, chaparral, woodland (Mar–May)
<input type="radio"/> <i>Apocynum cannabinum</i>	Indian Hemp	Apocyn-	-   -   A2	- NP	Moist places, near streams, springs, or as weed in orchards (Apr–Oct)
<input type="radio"/> <i>Aquilegia eximia</i>	Serpentine Columbine	Ranuncul-	-   -   A2	- NP	Generally serpentine seeps, occasionally moist ravines, mixed-evergreen or conifer forests (May–Oct)
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Ranuncul-	-   -   C	- NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Aster-	-   -   -	- NP	Common. Open to shady areas, often in drainages (May–Nov)
<input type="radio"/> <i>Artemisia dracunculus</i>	Wild Tarragon	Aster-	-   -   A1	- NP	Common. Many habitats, especially disturbed sites (Aug–Oct)
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Apocyn-	-   -   B	- NP	Flats, grassy or brushy hillsides (Apr–Jul)
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Apocyn-	-   -   -	- NP	Dry ground, valleys, foothills (May–Oct)
<input type="radio"/> <i>Astragalus didymocarpus</i> var. <i>didymocarpus</i>	Two-seed Milkvetch	Fab-	-   -   A1	- NA	Grassy areas (Feb–May)
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvetch	Fab-	-   -   -	- NA	Open, grassy areas, scrub (Mar–Jul)
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Brassic-	-   -   -	- NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Brassic-	-   -   -	- NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Orobanch-	-   -   -	L ZA	Disturbed grassland. (Apr–Jun)
<input type="radio"/> <i>Berula erecta</i>	Cutleaf Water-parsnip	Api-	-   -   B	- NP	Marshy areas, streams (Jul–Oct)
<input type="radio"/> <i>Boechera breweri</i> subsp. <i>breweri</i> (JM93: <i>Arabis breweri</i> var. <i>breweri</i> )	Brewer's Rockcress	Brassic-	-   -   A1	- NP	Rocky outcrops, ledges, talus Mar–Jul
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Api-	-   -   -	- NA	Shade of trees, rocks, shrubs (Mar–Apr)

Herbaceous <i>Brassica</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Brassica</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Brassic-	-   -   -	M ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Brassic-	-   -   -	L ZA	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Themid-	-   -   -	- NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Monti-	-   -   -	- NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Lili-	-   -   -	- NP	Common. Shady to open woodland, scrub (Apr–Jun)	
<input type="radio"/> <i>Calochortus argillosus</i>	Clay Mariposa Lily	Lili-	-   -   -	- NP	Hard clay from volcanic or metamorphic rocks (Apr–Jun)	
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Lili-	-   -   C	- NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)	
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Lili-	-   -   -	- NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Lili-	-   -   B	- NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)	
<input type="radio"/> <i>Calycadenia truncata</i>	Rosinweed	Aster-	-   -   A2	- NA	Common. Dry, open hillsides, rocky ridges, talus, grassland, openings in foothill woodland, scrub (May–Oct)	
<input type="radio"/> <i>Calystegia malacophylla subsp. pedicellata</i>	Woolly Morning-glory	Convolvul-	-   -   A2	- NP	Dry slopes, chaparral (Apr–Jul)	
<input type="radio"/> <i>Calystegia occidentalis subsp. occidentalis</i>	Western Morning-glory	Convolvul-	-   -   -	- NP	Dry slopes, chaparral, pine woodland (May–Jul)	
<input type="radio"/> <i>Calystegia purpurata subsp. purpurata</i>	Climbing Morning-glory	Convolvul-	-   -   -	- NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia sp.</i>	Morning-glory	Convolvul-	-   -   -	- NP	--	
<input type="radio"/> <i>Calystegia subacaulis subsp. subacaulis</i>	Shortstem Morning-glory	Convolvul-	-   -   -	- NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Camissonia contorta</i>	Contorted Suncup	Onagr-	-   -   B	- NA	Sandy soil, slopes, flats, often disturbed, grassland, chaparral, pinyon/juniper woodland (May–Ju)	
<input type="radio"/> <i>Camissoniopsis intermedia (JM93: Camissonia)</i>	Intermediate Suncup	Onagr-	-   -   A2	- NA	Shrubby slopes, especially burns (Mar–May)	
<input type="radio"/> <i>Campanula exigua</i>	Chaparral / Rock Harebell	Campanul-	-   1B.2   *A2	- NA	Talus slopes, generally serpentine soil (May–Jun)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Brassic-	-   -   -	- ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Brassic-	-   -   -	- NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Brassic-	-   -   -	- NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Carduus pycnocephalus subsp. pycnocephalus (JM93: no subsp.)</i>	Italian Thistle	Aster-	-   -   -	M ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Castilleja affinis subsp. affinis</i>	Common Indian Paintbrush	Orobanch-	-   -   -	- NP	Chaparral, coastal scrub (Mar–Jun)	



Herbaceous <i>Castilleja</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Castilleja</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Castilleja exserta</i> subsp. <i>exserta</i>	Purple Owl's-clover	Orobanch-	-   -   -	- NA	Open fields, grassland (Mar–May)
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Orobanch-	-   -   -	- NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)
<input type="radio"/> <i>Caulanthus lasiophyllus</i> (JM93: <i>Guillenia lasiophylla</i> )	California Mustard	Brassic-	-   -   -	- NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Aster-	-   -   -	M ZAB	Pastures, disturbed places (Apr–Nov)
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Aster-	-   -   -	M ZA	Disturbed fields, open woodland (Apr–Jul)
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Aster-	-   -   -	H ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)
<input type="radio"/> <i>Centromadia fitchii</i> (JM93: <i>Hemizonia</i> )	Fitch Spikeweed	Aster-	-   -   -	- NP	Grassland, ± alkaline flats, vernal pools, woodland, disturbed sites, serpentine (May–Nov)
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Caryophyll-	-   -   -	- ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)
<input type="radio"/> <i>Chamaesyce maculata</i>	Spotted Spurge	Euphorbi-	-   -   -	- ZA	Disturbed places (Apr–Oct)
<input type="radio"/> <i>Chamaesyce serpyllifolia</i> subsp. <i>serpyllifolia</i>	Thyme-leafed Spurge	Euphorbi-	-   -   B	- NA	Common. Dry habitats (Aug–Oct)
<input type="radio"/> <i>Chenopodium californicum</i>	California Goosefoot	Chenopodi-	-   -   -	- NP	Generally open sites, sandy to clay soils (Mar–Sep)
<input type="radio"/> <i>Chenopodium murale</i>	Wall Goosefoot	Chenopodi-	-   -   -	- ZA	Common. Disturbed areas, fields (All year)
<input type="radio"/> <i>Chlorogalum pomeridianum</i> var. <i>pomeridianum</i>	Wavyleaf Soap Plant	Agav-	-   -   -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)
<input type="radio"/> <i>Chorizanthe membranacea</i>	Pink Spineflower	Polygon-	-   -   A2	- NA	Common. Sand, gravel or rocks (Apr–Jul)
<input type="radio"/> <i>Cicuta douglasii</i>	Western Water-hemlock	Api-	-   -   B	- NP	Wet places, gen aquatic (Jun–Sep)
<input type="radio"/> <i>Cirsium cymosum</i> var. <i>cymosum</i> (JM93: no var.)	Peregrine Thistle	Aster-	-   -   B	- NP	Scrubland, woodland, open forest, meadows, occ serpentine (Jun–Jul)
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Aster-	-   -   -	- NB	Disturbed areas, grassland, woodland (May–Jul)
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Aster-	-   -   -	M ZB	Common. Disturbed areas (May–Oct)
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>concinna</i>	Red Ribbons	Onagr-	-   -   C	- NA	Mixed-evergreen forest, woodland, coastal scrub (Apr–Jul)
<input type="radio"/> <i>Clarkia gracilis</i> subsp. <i>gracilis</i>	Herald-of-summer	Onagr-	-   -   B	- NA	Common. Openings in woodland, forest (Apr–Jul)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. ?	Winecup Clarkia	Onagr-	-   -   -	- NA	--
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Onagr-	-   -   -	- NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Onagr-	-   -   -	- NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Onagr-	-   -   -	- NA	--
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Onagr-	-   -   -	- NA	Common. Woodland (Apr–Sep)

Herbaceous <i>Claytonia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Claytonia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i> )	Small-flower Claytonia	Monti-	-   -   -	- NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Monti-	-   -   -	- NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)	
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i> )	Yerba Buena	Lami-	-   -   -	- NP	Shady places, chaparral, woodland (Apr–Sep)	
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. ?	White Chinese Houses	Plantagin-	-   -   -	- NA	-	
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Plantagin-	-   -   -	- NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)	
<input type="radio"/> <i>Collinsia sparsiflora</i> var. ?	Few-flowered Collinsia	Plantagin-	-   -   -	- NA	-	
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Plantagin-	-   -   C	- NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)	
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Api-	-   -   -	M ZB	Common. Moist, esp disturbed places (Apr–Jul)	
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Convolvul-	-   -   -	- ZP	Roadsides, open areas in many pl communities (Mar–Oct)	
<input type="radio"/> <i>Cordylanthus rigidus</i> subsp. <i>rigidus</i>	Stiff Bird's Beak	Orobanch-	-   -   A1	- NA	Open foothill woodland, chaparral margins, conifer forest (Jul–Sep)	
<input type="radio"/> <i>Corethrogyne filaginifolia</i> (JM93: <i>Lessingia</i> , var. <i>filaginifolia</i> )	Common California-aster	Aster-	-   -   C	- NP	Coastal scrub, chaparral, grassland, foothill woodland, forest (Jul–Nov)	
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Aster-	-   -   -	L ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)	
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Crassul-	-   -   -	- NA	Open areas (Feb–May)	
<input type="radio"/> <i>Crepis vesicaria</i> subsp. <i>taraxacifolia</i>	Dandelion-leaf Hawksbeard	Aster-	-   -   -	- ZAB	Sandy clearings, hillsides, disturbed places (Feb–Oct)	
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i> )	Turkey-mullein	Euphorbi-	-   -   -	- NA	Dry, open, often disturbed areas (May–Oct)	
<input type="radio"/> <i>Cryptantha flaccida</i>	Weak-stemmed Cryptantha	Boragin-	-   -   -	- NA	Common; semi-barren, gravelly, loose soils, rocky sites, washes, slopes, ridges (Apr–Jul)	
<input type="radio"/> <i>Cuscuta californica</i> var. <i>californica</i>	California Dodder	Convolvul-	-   -   A2	- NA	On herbs, shrubs, roadsides, chaparral, grassland, yellow-pine forest (May–Sep)	
<input type="radio"/> <i>Cuscuta subinclusa</i>	Canyon Dodder	Convolvul-	-   -   B	- NP	Generally on herbs, shrubs, in forests near streams, river canyon bottoms, salt marshes (Mar–Oct(Dec?))	
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Boragin-	-   -   -	- NP	Chaparral, woodland (Feb–May)	
<input type="radio"/> <i>Datisca glomerata</i>	Durango Root	Datisc-	-   -   A2	- NP	Dry streambeds or washes (May–Jul)	
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Api-	-   -   -	- NA	Rocky or sandy places (Apr–Jun)	
<input type="radio"/> <i>Delphinium californicum</i> subsp. <i>californicum</i>	California Larkspur	Ranuncul-	-   -   A2	- NP	Generally slopes in dense chaparral, w side of coast ranges (Apr–Jun)	

Herbaceous <i>Delphinium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Delphinium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Ranuncul-	-   -   C	- NP	Open coastal grassland, chaparral (Mar–May)	
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>pallescens</i>	Pale-flowered Western Larkspur	Ranuncul-	-   -   C	- NP	Oak woodland, e slope coast ranges (Mar–May)	
<input type="radio"/> <i>Delphinium</i> sp.	Larkspur	Ranuncul-	-   -   -	- NP	- -	
<input type="radio"/> <i>Delphinium variegatum</i> subsp. <i>variegatum</i>	Royal Larkspur	Ranuncul-	-   -   C	- NP	Grassland, open oak woodland (Mar–May)	
<input type="radio"/> <i>Dichelostemma capitatum</i> subsp. <i>capitatum</i>	Blue Dicks	Themid-	-   -   -	- NP	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Themid-	-   -   -	- NP	Open woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Dichelostemma</i> sp.	Blue Dicks/ookow	Themid-	-   -   -	- NP	- -	
<input type="radio"/> <i>Dodecatheon clevelandii</i> subsp. <i>patulum</i>	Padre Shooting Star	Primul-	-   -   C	- NP	Moist places, often on serpentine or in ± alkaline sites (Mar–May)	
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Primul-	-   -   -	- NP	Gen in shady sites (Mar–Jul)	
<input type="radio"/> <i>Draba cuneifolia</i> (JM93: var. <i>integrifolia</i> )	Wedge-leaf Whitlow Grass	Brassic-	-   -   A1	- NA	Open or disturbed places (May)	
<input type="radio"/> <i>Draba verna</i>	Spring Whitlow Grass	Brassic-	-   -   B	- NA	Open or disturbed areas (Feb–May)	
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i> )	Sticky Cinquefoil	Ros-	-   -   -	- NP	Gen ± shady or moist areas (May–Jul)	
<input type="radio"/> <i>Dudleya cymosa</i> subsp. <i>paniculata</i>	Common / Hot Rock Dudleya	Crassul-	-   -   B	- NP	Uncommon. Rocky outcrops, canyons (May–Jun)	
<input type="radio"/> <i>Dysphania ambrosioides</i> (JM93: <i>Chenopodium</i> )	Mexican Tea	Chenopodi-	-   -   -	- ZAP	Disturbed places (Jul–Sep)	
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Onagr-	-   -   -	- NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)	
<input type="radio"/> <i>Epilobium campestre</i> (JM93: <i>E. pygmaeum</i> )	Smooth Willowherb	Onagr-	-   -   B	- NA	Vernal pools, clay mud flats (May–Sep)	
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Onagr-	-   -   -	- NP	Dry slopes, ridges (Jun–Dec)	
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Onagr-	-   -   -	- NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)	
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>watsonii</i>	Watson Willowherb	Onagr-	-   -   B	- NP	Moist coastal bluffs, streamsides, ± disturbed sites (May–Oct)	
<input type="radio"/> <i>Epilobium densiflorum</i>	Dense-flower Willowherb	Onagr-	-   -   C	- NA	Streambanks, outwashes, seasonal moist flats (May–Oct)	
<input type="radio"/> <i>Epilobium minutum</i>	Chaparral Willowherb	Onagr-	-   -   -	- NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)	
<input type="radio"/> <i>Erigeron canadensis</i> (JM93: <i>Conyza</i> )	Horseweed	Aster-	-   -   -	- ZA	Disturbed places (All year)	
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Aster-	-   -   -	- NP	Grassy dunes, chaparral, oak woodland (May–Oct)	
<input type="radio"/> <i>Erigeron philadelphicus</i> var. <i>philadelphicus</i> (JM93: no var.)	Philadelphia Fleabane	Aster-	-   -   C	- NBP	Streamsides, other moist habitats (May–Jun)	
<input type="radio"/> <i>Erigeron</i> sp. (JM93: <i>Conyza</i> )	Fleabane Daisy	Aster-	-   -   -	-	- -	
<input type="radio"/> <i>Eriogonum angulosum</i>	Angle-stem Wild Buckwheat	Polygon-	-   -   A2	- NA	Common. Clay (All year)	

Herbaceous <i>Eriogonum</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Eriogonum</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Eriogonum gracile</i> var. <i>gracile</i>	Slender Woolly Wild Buckwheat	Polygon-	-   -   B	- NA	Common. Sand (All year)	
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Polygon-	-   -   -	- NP	--	
<input type="radio"/> <i>Eriogonum roseum</i>	Wand Wild Buckwheat	Polygon-	-   -   B	- NA	Common. Sand or gravel (May–Nov)	
<input type="radio"/> <i>Eriogonum wrightii</i> var. <i>trachygonum</i>	Rough-node Bastard-sage	Polygon-	-   -   A2	- NP	Common. Gravel (Jul–Oct)	
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Erodium brachycarpum</i>	Southern European Filaree	Gerani-	-   -   -	- ZA	Dry, open or disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Gerani-	-   -   -	L ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)	
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Gerani-	-   -   -	- ZA	Open, disturbed sites (Feb–Sep)	
<input type="radio"/> <i>Erysimum capitatum</i> var. <i>capitatum</i> (JM93: var. <i>capitatum</i> , var. <i>angustatum</i> )	Western Wallflower	Brassic-	-   -   C	- NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)	
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Papaver-	-   -   -	- NP	Grassy, open areas (Feb–Sep)	
<input type="radio"/> <i>Eucrypta chrysanthemifolia</i> var. <i>chrysanthemifolia</i>	Common Eucrypta	Boragin-	-   -   B	- NA	Roadsides, burns, coastal bluffs, ravines (Mar–Jun)	
<input type="radio"/> <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i> )	Broadleaf Aster	Aster-	-   -   -	- NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)	
<input type="radio"/> <i>Euthamia occidentalis</i>	Western Goldenrod	Aster-	-   -   -	- NP	Marshes, streambanks, meadows (Jul–Nov)	
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Api-	-   -   -	H ZP	Roadsides, disturbed sites (May–Sep)	
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i> )	Checker Lily	Lili-	-   -   -	- NP	Common. Oak or pine scrub, grassland (Mar–Jun)	
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Rubi-	-   -   -	- NA	Grassy, ± shady places (Apr–Jun)	
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Rubi-	-   -   -	- ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)	
<input type="radio"/> <i>Galium parisiense</i>	Wall Bedstraw	Rubi-	-   -   -	- ZA	Warm, dry, generally rocky soil to moist areas, coastal-sage scrub, chaparral, grassy hillsides with oaks, roadsides (Apr–Aug)	
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Rubi-	-   -   -	- NP	Among shrubs in chaparral, forest (May–Aug)	
<input type="radio"/> <i>Galium</i> sp.	Bedstraw	Rubi-	-   -   -	-	--	
<input type="radio"/> <i>Galium trifidum</i> subsp. <i>columbianum</i> (JM93: var. <i>pacificum</i> )	Trifid Bedstraw	Rubi-	-   -   A2	- NP	Wet places, yellow-pine forest (Jul–Aug)	
<input type="radio"/> <i>Gamochaeta ustulata</i> (JM93: <i>Gnaphalium purpureum</i> )	Purple Cudweed	Aster-	-   -   -	- NA	Dunes, bluffs, fields, disturbed sites (Apr–Jul)	
<input type="radio"/> <i>Geranium carolinianum</i>	Carolina Geranium	Gerani-	-   -   -	- NA	Open to shaded sites, grassland, scrub, forest (Feb–Aug)	
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Gerani-	-   -   -	M ZA	Open, disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Gerani-	-   -   -	- ZA	Open to shaded sites, disturbed ground (Feb–Aug)	
<input type="radio"/> <i>Geranium robertianum</i>	Herb Robert	Gerani-	-   -   -	- ZAB	Open to shaded sites (Apr–Sep)	

Herbaceous <i>Gilia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Gilia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>achilleifolia</i>	California Gilia	Polemoni-	-   -   B	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>multicaulis</i>	Many-stem California Gilia	Polemoni-	-   -   -	- NA	Open or shaded, gen grassy places, sandy or rocky soil (Feb–Jun)
<input type="radio"/> <i>Gilia capitata</i> subsp. <i>staminea</i>	Pale Gilia	Polemoni-	-   -   B	- NA	Sandhills, flats (Mar–May)
<input type="radio"/> <i>Gilia tricolor</i> subsp. ?	Bird's-eye Gilia	Polemoni-	-   -   -	- NA	--
<input type="radio"/> <i>Gilia tricolor</i> subsp. <i>tricolor</i>	Bird's-eye Gilia	Polemoni-	-   -   B	- NA	Open, grassland, hills, valleys (Mar–May)
<input type="radio"/> <i>Glinus lotoides</i>	Damascisa	Mollugin-	-   -   -	- ZA	Uncommon. Moist or seasonally dry margins of wetlands (Jun–Nov)
<input type="radio"/> <i>Glycyrrhiza lepidota</i>	Wild Licorice	Fab-	-   -   C	- NP	In colonies, moist, gen open, disturbed areas incl streambanks, roadsides, alkaline soils or not (May–Jul)
<input type="radio"/> <i>Gnaphalium</i> or <i>Pseudognaphalium</i> sp. (JM93: <i>Gnaphalium</i> sp.)	Everlasting Cudweed	Aster-	-   -   -	- --	
<input type="radio"/> <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i> )	Great Valley Gumplant	Aster-	-   -   -	- NP	Sandy or saline bottomland, roadsides (May–Nov)
<input type="radio"/> <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i> )	Hairy Gumplant	Aster-	-   -   C	- NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)
<input type="radio"/> <i>Grindelia</i> sp.	Gumplant	Aster-	-   -   -	- NP	--
<input type="radio"/> <i>Helenium puberulum</i>	Rosilla	Aster-	-   -   -	- NB	Streambanks, seepage areas, lake margins (Jun–Aug)
<input type="radio"/> <i>Helianthella castanea</i>	Diablo Helianthella	Aster-	-   1B.2   *A2	- NP	Open, grassy sites (Apr–Jun)
<input type="radio"/> <i>Helianthus californicus</i>	California Sunflower	Aster-	-   -   C	- NP	Springs, marshes, streambanks, canyons (Jul–Oct)
<input type="radio"/> <i>Heliotropium curassavicum</i> var. <i>oculatum</i> (JM93: no var.)	Seaside Heliotrope	Boragin-	-   -   -	- NP	Moist to dry, saline to alkaline soils, gen near water (Feb–Oct)
<input type="radio"/> <i>Helminthotheca echioides</i> (JM93: <i>Picris</i> )	Bristly Ox-tongue	Aster-	-   -   -	L ZAB	Common. Disturbed areas (All year)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>lutescens</i> (JM93: subsp. <i>congesta</i> )	Yellow Hayfield Tarweed	Aster-	-   -   C	- NA	Grassland, barrens, openings in chaparral and woodland, often on serpentine (Apr–Dec)
<input type="radio"/> <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Aster-	-   -   -	- NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i> )	Cow Parsnip	Api-	-   -   -	- NP	Moist places, wooded or open (Apr–Jul)
<input type="radio"/> <i>Herniaria hirsuta</i> var. <i>cinerea</i> (JM93: subsp.)	Ash-color Rupturewort	Caryophyll-	-   -   -	- ZA	Disturbed areas, alkaline hills, clay flats (Spring–fall)
<input type="radio"/> <i>Hesperivax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperivax	Aster-	-   -   -	- NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)

Herbaceous <i>Hesperocnide</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Hesperocnide</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Urtic-	-   -   -	- NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)
<input type="radio"/> <i>Heterotheca oregona var. rudis</i>	Inland Oregon Goldenaster	Aster-	-   -   A1	- NP	Seasonally dry streambed (Jul–Oct)
<input type="radio"/> <i>Hieracium albiflorum</i>	White Hawkweed	Aster-	-   -   B	- NP	Forest (May–Sep)
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Brassic-	-   -   -	M ZBP	Disturbed areas (Apr–Oct)
<input type="radio"/> <i>Hoita macrostachya</i>	Leather Root	Fab-	-   -   B	- NP	Streamsides, marshes, spring-moist places (May–Aug)
<input type="radio"/> <i>Holocarpha heermannii</i>	Heermann Tarplant	Aster-	-   -   -	- NA	Grassland (May–Nov)
<input type="radio"/> <i>Holocarpha virgata subsp. virgata</i>	Sticky Tarplant	Aster-	-   -   -	- NA	Grassland (May–Nov)
<input type="radio"/> <i>Horkelia sp.</i>	Horkelia	Ros-	-   -   -	- NP	- -
<input type="radio"/> <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Aster-	-   -   -	L ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)
<input type="radio"/> <i>Hypochaeris radicata</i>	Rough Cat's-ear	Aster-	-   -   -	M ZP	Disturbed areas, grassland, open woodland (Apr–Jul)
<input type="radio"/> <i>Kickxia elatine</i>	Sharp Point Fluvellin	Plantagin-	-   -   -	- ZA	Disturbed, open places (Apr–Oct)
<input type="radio"/> <i>Lactuca saligna</i>	Willow Lettuce	Aster-	-   -   -	- ZA	Roadsides, grassland (Jul–Nov)
<input type="radio"/> <i>Lactuca serriola</i>	Prickly Lettuce	Aster-	-   -   -	- ZA	Abundant. Disturbed places (May–Oct)
<input type="radio"/> <i>Lactuca virosa</i>	Tall Wild Lettuce	Aster-	-   -   -	- ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)
<input type="radio"/> <i>Lagophylla ramosissima (JM93: subsp. ramosissima)</i>	Common Hare-leaf	Aster-	-   -   -	- NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lamium amplexicaule</i>	Clasping Henbit	Lami-	-   -   -	- ZAB	Disturbed sites, cult or abandoned fields (Apr–Sep)
<input type="radio"/> <i>Lasthenia californica subsp. californica (JM93: no subsp.)</i>	California Goldfields	Aster-	-   -   -	- NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lasthenia glaberrima</i>	Vernal Pool Goldfields	Aster-	-   -   B	- NA	
<input type="radio"/> <i>Lathyrus jepsonii var. californicus</i>	Bluff Pea	Fab-	-   -   B	- NP	Forest, open areas (Apr–Aug)
<input type="radio"/> <i>Lathyrus vestitus var. vestitus</i>	Pale Purple Pacific Pea	Fab-	-   -   -	- NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)
<input type="radio"/> <i>Layia chrysanthemoides</i>	Chrysanthemum Tidytops	Aster-	-   -   A2	- NA	Grassy or open heavy soil, sometimes ± alkaline (Mar–Jun)
<input type="radio"/> <i>Layia gaillardiioides</i>	Yellow Woodland Tidytops	Aster-	-   -   A2	- NA	Open or semi-shaded slopes, in sandy or clayey soil (including serpentine) (Mar–Aug)
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Aster-	-   -   C	- NA	Many habitats (Feb–Jul)
<input type="radio"/> <i>Lepidium nitidum (JM93: var. nitidum, oreganum)</i>	Threadleaf Peppergrass	Brassic-	-   -   -	- NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)

Herbaceous <i>Lepidium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Lepidium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Lepidium strictum</i>	Prostrate Peppergrass	Brassic-	-   -   -	- ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)
<input type="radio"/> <i>Leptosiphon ambiguus</i> (JM93: <i>Linanthus</i> )	Serpentine Leptosiphon	Polemoni-	-   4.2   *A2	- NA	Grassy areas gen serpentine (Apr–May)
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i> )	Pinklobe Leptosiphon	Polemoni-	-   -   C	- NA	Open or shaded areas in woodland, chaparral (Apr–Jun)
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i> )	Bicolor Leptosiphon	Polemoni-	-   -   -	- NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. ? (JM93: no subsp.)	Evening Snow	Polemoni-	-   -   A2	- NA	--
<input type="radio"/> <i>Linum bienne</i>	Narrow-leaf Flax	Lin-	-   -   -	- ZP	Garden escape; grassland, woodland, disturbed places, esp coastal (Mar–Jun)
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Saxifrag-	-   -   -	- NP	Open, grassy slopes (Mar–Apr)
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Saxifrag-	-   -   -	- NP	Shaded slopes (Feb–Jun)
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i> )	California Fluffweed	Aster-	-   -   -	- NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i> )	Daggerleaf Cottonrose	Aster-	-   -   -	- ZA	Bare or grassy openings, burns (Mar–Jul)
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Api-	-   -   B	- NP	Woodland, brushy slopes (Apr–Jun)
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Api-	-   -   -	- NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)
<input type="radio"/> <i>Lomatium nudicaule</i>	Swollen-stem Biscuit Root	Api-	-   -   -	- NP	Rocky slopes, flats, gen pine woodland (Apr–Jun)
<input type="radio"/> <i>Lomatium</i> sp.	Biscuit Root	Api-	-   -   -	- NP	--
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Api-	-   -   -	- NP	Open grassy slopes, meadows, woodland (Feb–May)
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas (Mar–Jun)
<input type="radio"/> <i>Lupinus</i> sp.	Lupine	Fab-	-   -   -	- NAP	--
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Fab-	-   -   -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)
<input type="radio"/> <i>Lycopus</i> sp.	Water Horehound / Bugleweed	Lami-	-   -   -	- NP	--
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Lythr-	-   -   -	- ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)
<input type="radio"/> <i>Madia anomala</i>	Plump-seed Tarweed	Aster-	-   -   A1	- NA	Openings in woodland or chaparral (Apr–Jun)

Herbaceous	Grouped by Growth Form					Herbaceous
<i>Madia</i>	Sorted Alphabetically by Scientific Name					<i>Madia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Aster-	-   -   -	- NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)	
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Aster-	-   -   -	- NA	Grassy, open, or disturbed sites (May–Oct)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i> )	Starry False Solomon's Seal	Rusc-	-   -   -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
<input type="radio"/> <i>Malva parviflora</i>	Cheeseweed	Malv-	-   -   -	- ZA	Common. Disturbed places (Mar–May)	
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i> )	California Man-root	Cucurbit-	-   -   -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Lami-	-   -   -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i> )	Pineapple Weed	Aster-	-   -   -	- ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Fab-	-   -   -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i> )	Sourclover	Fab-	-   -   -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Melissa officinalis</i>	Lemon Balm	Lami-	-   -   -	- ZP	Moist sites, meadows, fields (Jun–Sep)	
<input type="radio"/> <i>Mentha pulegium</i>	Pennyroyal	Lami-	-   -   -	M ZP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha spicata</i> (JM93: var. <i>spicata</i> )	Spearmint	Lami-	-   -   -	- ZP	Moist places, marshes, lakeshores, fields (Jul–Oct)	
<input type="radio"/> <i>Mentzelia affinis</i>	Hydra Blazing Star	Loas-	-   -   A2	- NA	Rocky or gray-white soils in grassland, woodland, creosote-bush scrub (Apr–May)	
<input type="radio"/> <i>Mentzelia lindleyi</i>	Lindley Blazing Star	Loas-	-   -   A2	- NA	Rocky, open slopes, coastal-sage scrub, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Micranthes californica</i> (JM93: <i>Saxifraga</i> )	California Saxifrage	Saxifrag-	-   -   -	- NP	Moist, shady places (Feb–May(Jun))	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Aster-	-   -   -	- NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Aster-	-   -   -	- NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris elegans</i>	Nodding Silverpuffs	Aster-	-   -   A2	- NA	Generally inland clay grassland, often near vernal pools (Apr–Jun)	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i> )	Slender Annual Phlox	Polemoni-	-   -   -	- NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Mimulus cardinalis</i>	Scarlet Monkeyflower	Phrym-	-   -   C	- NP	Moist to wet places along streams, seepage areas (May–Sep)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Phrym-	-   -   -	- NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Mimulus latidens</i>	Broadtooth Monkeyflower	Phrym-	-   -   A2	- NA	Vernally wet depressions (Apr–Jun)	
<input type="radio"/> <i>Mimulus pilosus</i>	Downy Monkeyflower	Phrym-	-   -   A2	- NA	Moist, sandy areas, especially by small streams, disturbed areas (Apr–Aug)	



Herbaceous <i>Minuartia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Minuartia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Minuartia douglasii</i>	Douglas' Stitchwort	Caryophyll-	-   -   B	- NA	Dry, rocky, slopes, flats in chaparral, oak and pine woodland, often serpentine (Spring–early summer)	
<input type="radio"/> <i>Monardella villosa</i> subsp. <i>villosa</i> (JM93: <i>M. villosa</i> subsp. <i>villosa</i> , <i>M. antonina</i> subsp. <i>antonina</i> )	Coyote-mint	Lami-	-   -   -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Monolopia major</i>	Cupped Monolopia	Aster-	-   -   B	- NA	Grassland, bare clay (Feb–Jul)	
<input type="radio"/> <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i> )	Water Cress	Brassic-	-   -   -	- NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Polemoni-	-   -   C	- NA	Open, slopes, gravel, clay (May–Jul)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Polemoni-	-   -   -	- NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Boragin-	-   -   -	- NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)	
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Boragin-	-   -   -	- NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)	
<input type="radio"/> <i>Nemophila parviflora</i> var. <i>parviflora</i>	Small-flower Nemophila	Boragin-	-   -   C	- NA	Woodland, forest, roadsides, slopes (Mar–Jul)	
<input type="radio"/> <i>Nicotiana quadrivalvis</i>	Indian Tobacco	Solan-	-   -   B	- NA	Open, well-drained washes, slopes (May–Oct)	
<input type="radio"/> <i>Orobanche fasciculata</i>	Clustered Broomrape	Orobanch-	-   -   B	- NP	Dry, gen bare places. Root parasite on various shrubs (Apr–Jul)	
<input type="radio"/> <i>Orobanche vallicola</i>	Sharp-lobe Broom-rape	Orobanch-	-   -   A2	- NP	Riparian woodland, on <i>Sambucus nigra</i> (Jul–Sep )	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i> )	Sweet-cicely	Api-	-   -   -	- NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza brachypoda</i>	California Sweet Cicely	Api-	-   -   B	- NP	Moist ravines, conifer forest, woodland (Mar–May)	
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Oxalid-	-   -   -	M ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)	
<input type="radio"/> <i>Packera breweri</i> (JM93: <i>Senecio</i> )	Brewer's Ragwort	Aster-	-   -   B	- NP	Common. Seasonally damp grassland, oak savanna, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Papaver californicum</i>	Fire Poppy	Papaver-	-   -   A2	- NA	Burns, disturbed areas, open woodland (Apr–May)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i> )	Wind Poppy	Papaver-	-   -   -	- NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Pectocarya penicillata</i>	Northern Pectocarya	Boragin-	-   -   A1	- NA	Disturbed sites, roadsides (Feb–May)	
<input type="radio"/> <i>Pectocarya pusilla</i>	Little Pectocarya	Boragin-	-   -   B	- NA	Dry, semi-barren sites in grassland, chaparral, woodland, roadsides (Mar–Jun)	
<input type="radio"/> <i>Penstemon centranthifolius</i>	Scarlet Bugler	Plantagin-	-   -   A2	- NP	Dry, open chaparral or oak woodland (Apr–Jul)	
<input type="radio"/> <i>Penstemon heterophyllus</i> var. <i>heterophyllus</i>	Foothill Penstemon	Plantagin-	-   -   B	- NP	Grassland, chaparral, forest openings (Apr–Jul)	
<input type="radio"/> <i>Perideridia californica</i>	California Yampah	Api-	-   -   A2	- NP	Damp soil by streams (Apr–May)	

Herbaceous <i>Perideridia</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Perideridia</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Api-	-   -   -	- NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Persicaria punctata</i> (JM93: <i>Polygonum punctatum</i> )	Water Smartweed	Polygon-	-   -   C	- NAP	Shallow water, shores, marshes, floodplain forest (Jun–Nov)	
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Boragin-	-   -   -	- NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)	
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Boragin-	-   -   -	- NA	Common. Clay to rocky soils, slopes (Mar–May)	
<input type="radio"/> <i>Phacelia imbricata</i> subsp. <i>imbricata</i>	Rock Phacelia	Boragin-	-   -   -	- NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Phacelia malvifolia</i>	Stinging Phacelia	Boragin-	-   -   A2	- NA	Sandy or gravelly soils, scrub, conifer forest (Apr–Jul)	
<input type="radio"/> <i>Phacelia rattanii</i>	Rattan Phacelia	Boragin-	-   -   A1	- NA	Shaded crevices, steep slopes (May–Jul)	
<input type="radio"/> <i>Pholistoma auritum</i> var. <i>auritum</i>	Fiesta Flower	Boragin-	-   -   B	- NA	Ocean bluffs, talus slopes, woodland, streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Pholistoma membranaceum</i>	White Fiesta Flower	Boragin-	-   -   -	- NA	Beaches, bluffs, ravines, wooded slopes, desert washes (Feb–May)	
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>tomentosum</i> (JM93: <i>P. villosum</i> )	Oak Mistletoe	Visc-	-   -   -	- NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)	
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i> )	Lemon Verbena	Verben-	-   -   -	- NP	Wet places, pond margins (May–Nov)	
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Boragin-	-   -   -	- NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)	
<input type="radio"/> <i>Plagiobothrys humistratus</i>	Low Popcornflower	Boragin-	-   -   B	- NA	Vernal pools, wet places, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Boragin-	-   -   -	- NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys</i> sp.	Popcornflower	Boragin-	-   -   -	-	- -	
<input type="radio"/> <i>Plagiobothrys trachycarpus</i>	Rough-nutlet Popcornflower	Boragin-	-   -   B	- NA	Shallow vernal pools, wet places in grassland, scrub, chaparral, woodland (Mar–May)	
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Plantagin-	-   -   -	- NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)	
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Plantagin-	-   -   -	L ZA	Common. Disturbed areas (Apr–Aug)	
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Plantagin-	-   -   -	- ZAP	Disturbed areas (Apr–Sep)	
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Papaver-	-   -   -	- NA	Open grassland, sandy soil, burns (Mar–May)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: subsp. <i>ciliosa</i> , <i>insignis</i> )	Longspur Plectritis	Valerian-	-   -   B	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Valerian-	-   -   -	- NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Lami-	-   -   -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i> )	Knotweed	Polygon-	-   -   -	- ZA	Disturbed places (May–Nov)	

Herbaceous <i>Potamogeton</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name				Herbaceous <i>Potamogeton</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)
<input type="radio"/> <i>Potamogeton</i> or <i>Stuckenia</i> sp. (JM93: <i>Potamogeton</i> sp.)	Pondweed	Potamogeton-	-   -   -	- NP	--
<input type="radio"/> <i>Prunella vulgaris</i> var. ?	European Selfheal	Lami-	-   -   -	- ZP	--
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i> )	California Everlasting	Aster-	-   -   -	- NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i> )	Weedy Cudweed	Aster-	-   -   -	- ZA	Disturbed sites, fields, streambeds (Apr–Aug)
<input type="radio"/> <i>Pseudognaphalium microcephalum</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>microcephalum</i> )	White Everlasting	Aster-	-   -   B	- NB	Grassy hillsides, gravelly canyon bottoms, chaparral, coastal-sage scrub (Jun–Aug)
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i> )	Cotton-batting Plant	Aster-	-   -   -	- NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Polygon-	-   -   -	- NA	Common. Sand or gravel (Mar–Jul)
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Aster-	-   -   B	- NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)
<input type="radio"/> <i>Ranunculus aquatilis</i> var. <i>diffusus</i> (JM93: var. <i>capillaceus</i> )	Thread-leaf Water Buttercup	Ranuncul-	-   -   -	- NP	Ponds, lakes, streams (Mar–Sep)
<input type="radio"/> <i>Ranunculus californicus</i> var. <i>californicus</i> (JM93: no var.)	California Buttercup	Ranuncul-	-   -   -	- NP	Grassland, open woodland (Mar–Aug)
<input type="radio"/> <i>Ranunculus canus</i> var. <i>canus</i> (JM93: no var.)	Sacramento Valley Buttercup	Ranuncul-	-   -   B	- NP	Grassland, open oak woodlan (Mar–Jul)
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Ranuncul-	-   -   -	- NA	Grassland, open woodland (Mar–May)
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Ranuncul-	-   -   -	- ZA	Stream-banks, drainages, low meadows (Apr–Jun)
<input type="radio"/> <i>Raphanus sativus</i>	Radish	Brassic-	-   -   -	L ZA	Disturbed areas, fields (May–Jul)
<input type="radio"/> <i>Rigiopappus leptocladus</i>	Bristle-head	Aster-	-   -   B	- NA	Grassy sites, openings among shrubs or trees (Apr–Jul)
<input type="radio"/> <i>Rorippa curvisiliqua</i>	Yellow Water Cress	Brassic-	-   -   A2	- NA	Uncommon. Streambanks, marshy ground, seepage areas, mud flats, meadows (May–Oct)
<input type="radio"/> <i>Rumex acetosella</i>	Sheep Sorrel	Polygon-	-   -   -	M ZP	± Disturbed, often acidic places (Apr–Jul)
<input type="radio"/> <i>Rumex conglomeratus</i>	Green Dock	Polygon-	-   -   -	- ZP	Common. Moist places (May–Aug)
<input type="radio"/> <i>Rumex crispus</i>	Curly Dock	Polygon-	-   -   -	L ZP	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Rumex fueginus</i> (JM93: <i>R. maritimus</i> )	Golden Dock	Polygon-	-   -   A2	- NA	Riparian, disturbed places, shores, marshes, bogs, wet meadows (May–Aug)
<input type="radio"/> <i>Rumex pulcher</i>	Fiddle Dock	Polygon-	-   -   -	- ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)
<input type="radio"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i> )	Thin-leaf Willow Dock	Polygon-	-   -   B	- NP	Wet places, margins, rocky slopes (May–Jul)
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Fab-	-   -   -	- NP	Woodland (May–Sep)

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b>Sagina</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Sagina</b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Caryophyll-	-   -   -	- NA	Sandy disturbed areas, river bars, streamsides (Spring–early summer)	
<input type="radio"/> <i>Salvia columbariae</i>	Chia	Lami-	-   -   B	- NA	Dry, disturbed sites, chaparral, coastal-sage scrub (Mar–Jun)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Api-	-   -   -	- NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Api-	-   -   -	- NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Api-	-   -   -	- NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Api-	-   -   -	- ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Scrophularia californica</i> (JM93: <i>subsp. californica, floribunda</i> )	California Figwort	Scrophulari-	-   -   -	- NP	Common; damp places, chaparral, roadsides (Mar–Jul)	
<input type="radio"/> <i>Scutellaria siphocampyloides</i>	Grayleaf Skullcap	Lami-	-   -   A1	- NP	Open sites, seeps, dry stream beds, scrub, woodland (May–Jul)	
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberos Skullcap	Lami-	-   -   C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sedum spathulifolium</i>	Broadleaf Stonecrop	Crassul-	-   -   B	- NP	Outcrops, often in shade (Apr–Aug)	
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Aster-	-   -   C	- NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Aster-	-   -   -	- ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Rubi-	-   -   -	- ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Sidalcea malviflora subsp. laciniata</i>	Geranium-leaved Checkerbloom	Malv-	-   -   C	- NP	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Caryophyll-	-   -   -	- ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silene laciniata subsp. californica</i> (JM93: <i>S. californica</i> )	California Pink	Caryophyll-	-   -   B	- NP	Chaparral, oak woodland, conifer forest, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Silene verecunda</i> (JM93: <i>subsp. platyota</i> )	Cuyamaca Catchfly	Caryophyll-	-   -   A2	- NP		
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Aster-	-   -   -	L ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Sisymbrium officinale</i>	Hedge Mustard	Brassic-	-   -   -	- ZA	Disturbed areas, fields, pastures (Apr–Sep)	
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Irid-	-   -   -	- NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
<input type="radio"/> <i>Solanum americanum</i>	Small-flowered Nightshade	Solan-	-   -   -	- NAP	Open, often disturbed places (Apr–Nov)	
<input type="radio"/> <i>Solidago elongata</i> (JM93: <i>S. canadensis subsp. elongata</i> )	West Coast Canada Goldenrod	Aster-	-   -   -	- NP	Meadows, thickets (May–Oct)	
<input type="radio"/> <i>Solidago velutina subsp. californica</i> (JM93: <i>S. californica</i> )	California Goldenrod	Aster-	-   -   -	- NP	Woodland margins, grassland, disturbed soils (May–Nov)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Aster-	-   -   -	- ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	

Herbaceous <b>Sonchus</b>	<b>Grouped by Growth Form</b>					Herbaceous <b>Sonchus</b>
<b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Aster-	-   -   -	-	ZA	Common. Slightly moist disturbed sites, along streams (All year)
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Aster-	-   -   -	-	ZA	Abundant. Disturbed places (All year)
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Caryophyll-	-   -   -	-	ZAP	Forest, meadows, mud flats, disturbed
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Lami-	-   -   C	-	NP	Streambanks, springs, pine/oak forest (Jun–Oct)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i> )	Common Rigid Hedge-nettle	Lami-	-   -   -	-	NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Stebbinsoseris heterocarpa</i>	Grassland Silverpuffs	Aster-	-   -   B	-	NA	Open, sometimes disturbed sites, rocky to clay soils, generally inland (Apr–Jun)
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Caryophyll-	-   -   -	-	ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Caryophyll-	-   -   C	-	NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)
<input type="radio"/> <i>Stephanomeria virgata</i> subsp. <i>pleurocarpa</i>	Twiggy Wreath Plant	Aster-	-   -   C	-	NA	Chaparral openings, grassland (Jun–Nov)
<input type="radio"/> <i>Streptanthus glandulosus</i> subsp. <i>glandulosus</i> (JM93: incl. <i>S. albidus</i> subsp. <i>peramoenus</i> )	Bristly Jewel Flower	Brassic-	-   -   B	-	NA	Serpentine, bare slopes, chaparral & woodland openings (Apr–Jul)
<input type="radio"/> <i>Symphyotrichum chilense</i> (JM93: <i>Aster chilensis</i> )	Common California Wild Aster	Aster-	-   -   -	-	NP	Grassland, salt marshes, disturbed places (Jun–Oct)
<input type="radio"/> <i>Symphyotrichum subulatum</i> var. <i>parviflorum</i> (JM93: <i>Aster subulatus</i> var. <i>ligulatus</i> )	Annual Saltmarsh Aster	Aster-	-   -   -	-	NA	Marshes, disturbed places (Jul–Oct)
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Aster-	-   -   -	-	ZBP	Abundant. Esp disturbed areas (All year)
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Api-	-   -   C	-	NP	Chaparral, pine/oak woodland (Mar–May)
<input type="radio"/> <i>Thalictrum fendleri</i> var. <i>polycarpum</i>	Foothill Meadow-rue	Ranuncul-	-   -   -	-	NP	Moist, open to shaded places, woodland, forest (Mar–Jun)
<input type="radio"/> <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Brassic-	-   -   -	-	NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)
<input type="radio"/> <i>Thysanocarpus laciniatus</i> var. <i>laciniatus</i> (JM93: no var.)	Narrowleaf Fringepod	Brassic-	-   -   B	-	NA	Oak woodland, rocky ridges, slopes, chaparral, washes (Mar–May)
<input type="radio"/> <i>Tonella tenella</i>	Small-flowered Tonella	Plantagin-	-   -   A2	-	NA	Moist, shaded places in chaparral, oak and mixed woodland (Mar–Jun)
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Api-	-   -   -	M	ZA	Disturbed places (Apr–Jul)
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Api-	-   -   -	-	ZA	Disturbed places (Apr–Jun)
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Aster-	-   -   -	-	ZBP	Common. Disturbed places (Mar–Nov)
<input type="radio"/> <i>Trichostema lanceolatum</i>	Vinegar Weed	Lami-	-   -   -	-	NA	Dry, open, gen disturbed habitats (Jun–Nov)
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decepiens</i>	Deceiving Clover	Fab-	-   -   -	-	NA	Open, grassy areas, forest (Apr–Jun)

Herbaceous <i>Trifolium</i>	Grouped by Growth Form Sorted Alphabetically by Scientific Name					Herbaceous <i>Trifolium</i>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Fab-	-   -   -	- NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Fab-	-   -   -	- NA	Grassland, coastal woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Fab-	-   -   -	- NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Fab-	-   -   -	- ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium fucatum</i>	Bull Clover	Fab-	-   -   C	- NA	Locally abundant. Moist, open grassland, marshes, roadsides, occasionally saline or serpentine soils (Apr–Jun)	
<input type="radio"/> <i>Trifolium glomeratum</i>	Clustered Clover	Fab-	-   -   -	- ZA	Uncommon. Disturbed areas (Mar–May)	
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i> )	Pinpoint Clover	Fab-	-   -   -	- NA	Open, disturbed places, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Fab-	-   -   -	M ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Fab-	-   -   -	- NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium obtusiflorum</i>	Clammy Clover	Fab-	-   -   B	- NA	Moist disturbed areas, gravel bars, marshes (Apr–Jul)	
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Fab-	-   -   C	- NA	Woody or shrubby slopes, roadsides (Mar–Jun)	
<input type="radio"/> <i>Trifolium olivaceum</i> (JM93: <i>T. albopurpureum</i> var. <i>olivaceum</i> )	Olive Clover	Fab-	-   -   B	- NA	Grassy slopes, valley meadows, disturbed areas (Apr–May)	
<input type="radio"/> <i>Trifolium</i> sp.	Clover	Fab-	-   -   -	- -		
<input type="radio"/> <i>Trifolium variegatum</i> var. ? (JM93: no var.)	White-tip Clover	Fab-	-   -   -	- NA	- -	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Fab-	-   -   -	- NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Melanthi-	-   -   -	- NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)	
<input type="radio"/> <i>Triphysaria eriantha</i> subsp. <i>eriantha</i>	Yellow Johnny-tuck	Orobanch-	-   -   -	- NA	Grassland, foothills (Mar–May)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Orobanch-	-   -   -	- NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Themid-	-   -   -	- NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Brassic-	-   -   B	- NP	Grassy banks, open fields, roadsides, pastures (Mar–May)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Aster-	-   -   -	- NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urtica dioica</i> subsp. <i>holosericea</i>	Hoary Nettle	Urtic-	-   -   -	- NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	

<b>Herbaceous</b>	<b>Grouped by Growth Form</b>					<b>Herbaceous</b>
<b><i>Urtica</i></b>	<b>Sorted Alphabetically by Scientific Name</b>					<b><i>Urtica</i></b>
<b>✓ Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv OL</b>	<b>Habitat (Bloom)</b>	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Urtic-	-   -   -	- ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
<input type="radio"/> <i>Verbascum thapsus</i>	Woolly Mullein	Scrophulari-	-   -   -	L ZB	Roadsides, streambanks, disturbed areas (May–Sep)	
<input type="radio"/> <i>Verbena lasiostachys var. scabrida</i>	Robust Vervain	Verben-	-   -   -	- NP	Open, dry to wet places (May–Sep)	
<input type="radio"/> <i>Veronica americana</i>	American Brooklime	Plantagin-	-   -   C	- NP	Common. Moist to wet soil, springs, slow streams, meadows, lakeshores (May–Aug)	
<input type="radio"/> <i>Veronica arvensis</i>	Common Speedwell	Plantagin-	-   -   -	- ZA	Meadows (Apr–Jul)	
<input type="radio"/> <i>Veronica peregrina subsp. xalapensis</i>	Purslane Speedwell	Plantagin-	-   -   -	- NA	Moist places (Apr–Aug)	
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Plantagin-	-   -   -	- ZA	Wet, disturbed areas, fields (Feb–May)	
<input type="radio"/> <i>Vicia americana subsp. americana (JM93: var. americana)</i>	American Vetch	Fab-	-   -   -	- NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. nigra</i>	Narrow-leaved Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa subsp. sativa</i>	Spring Vetch	Fab-	-   -   -	- ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. varia</i>	Sparsely Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa subsp. villosa</i>	Hairy Vetch	Fab-	-   -   -	- ZAB	Grassland, roadside, disturbed areas (May–Jul)	
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Viol-	-   -   C	- NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)	
<input type="radio"/> <i>Viola purpurea subsp. purpurea</i>	Mountain Violet	Viol-	-   -   A2	- NP	In openings or beneath shrubs, generally in yellow-pine forest or higher (Mar–Jul)	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Aster-	-   -   -	- NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Aster-	-   -   C	- NP	Gen shady sites (Mar–Jun)	
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Aster-	-   -   -	- NP	Open grassland, woodland, scrub (Mar–May(Aug))	
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Aster-	-   -   -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Xanthium strumarium</i>	Cocklebur	Aster-	-   -   -	- NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Api-	-   -   C	- NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Zeltnera davyi (JM93: Centaurium)</i>	Davy's Centaury	Gentian-	-   -   B	- NA	Moist coastal bluffs, dunes, open forest (May–Aug)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Acer</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Acer</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Sapind-	-   -   -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Ros-	-   -   -	- NP	Dry slopes, ridges, chaparral (May–Jun)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Sapind-	-   -   -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Betul-	-   -   -	- NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Ros-	-   -   C	- NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr–Jun)	
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Eric-	-   -   C	- NP	Conifer, oak forests (Mar–May)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Eric-	-   -   C	- NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Aster-	-   -   -	- NP	Coastal scrub, chaparral, open woodland (Aug–Nov)	
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Aster-	-   -   -	- NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)	
<input type="radio"/> <i>Baccharis salicifolia</i> subsp. <i>salicifolia</i> (JM93: no subsp.)	Mule Fat	Aster-	-   -   -	- NP	Riparian woodland, canyon bottoms, disturbed sites, often forming thickets (All year)	
<input type="radio"/> <i>Brickellia californica</i>	California Brickellbush	Aster-	-   -   B	- NP	Dry, rocky hillsides, canyons, sea bluffs (Jul–Dec)	
<input type="radio"/> <i>Cercis occidentalis</i>	Western Redbud	Fab-	-   -   A1	- NP	Dry, shrubby slopes, canyons, streambanks, chaparral, foothill woodland, yellow-pine forest (Mar–May)	
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Ros-	-   -   C	- NP	Dry, rocky slopes, chaparral (Mar–May)	
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Ranuncul-	-   -   -	- NP	Hillsides, chaparral, open woodland (Jan–Jun)	
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Ranuncul-	-   -   C	- NP	Along streams, wet places (Jun–Sep)	
<input type="radio"/> <i>Ericameria nauseosa</i> var. <i>mohavensis</i> (JM93: <i>Chrysothamnus nauseosus</i> subsp. <i>mohavensis</i> )	Mojave Rabbitbrush	Aster-	-   -   A2	- NP	Common. Dry scrub (Aug–Oct)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>foliolosum</i>	Leafy California Buckwheat	Polygon-	-   -   A2	- NP	Gravel (All year)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	Mojave Desert Calif. Buckwheat	Polygon-	-   -   A1	- NP	Sand, gravel or rocks (All year)	
<input type="radio"/> <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Aster-	-   -   -	- NP	Many dry habitats (Apr–Aug)	
<input type="radio"/> <i>Eriophyllum jepsonii</i>	Jepson's Woolly Sunflower	Aster-	-   4.3   *A2	- NP	Dry oak woodland (Apr–Jun)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Myrt-	-   -   -	M ZP	Common. Disturbed areas (Oct–Jan)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>californica</i> (JM93: <i>Rhamnus</i> )	California Coffee Berry	Rhamn-	-   -   -	- NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Frangula californica</i> subsp. <i>tomentella</i> (JM93: <i>Rhamnus tomentella</i> subsp. <i>tomentella</i> )	Hoary Coffee Berry	Rhamn-	-   -   B	- NP	Chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Fraxinus latifolia</i>	Oregon Ash	Ole-	-   -   B	- NP	Canyons, streambanks, woodland (Mar–May)	



<b>Woody</b> <b>Gutierrezia</b>	<b>Grouped by Growth Form</b> <b>Sorted Alphabetically by Scientific Name</b>					<b>Woody</b> <b>Gutierrezia</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Gutierrezia californica</i>	California Matchweed	Aster-	-   -   C	- NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Ros-	-   -   -	- NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)	
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Ros-	-   -   -	- NP	Moist woodland edges, rocky slopes (May–Aug)	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i> )	N. California Black Walnut	Jugland-	-   1B.1   *A2	- NP	Along streams, disturbed slopes (Apr–May)	
<input type="radio"/> <i>Keckiella corymbosa</i>	Redwood / Red Rock Penstemon	Plantagin-	-   -   A1	- NP	Rocky slopes in conifer or hardwood forests, (chaparral) (Jun–Oct)	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i> )	Hairy Vine Honeysuckle	Caprifoli-	-   -   -	- NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera subspicata</i> var. <i>denudata</i>	Southern Honeysuckle	Caprifoli-	-   -   B	- NP	Chaparral slopes (Jun–Jul)	
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Fab-	-   -   -	- NP	- -	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>collinus</i>	Bay Area Silver Lupine	Fab-	-   -   -	- NP	Cliffs, forest openings (Mar–Jun)	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Phrym-	-   -   -	- NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Nicotiana glauca</i>	Tree Tobacco	Solan-	-   -   -	M ZP	Open, disturbed flats or slopes (Apr–Aug)	
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Ros-	-   -   -	- NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)	
<input type="radio"/> <i>Physocarpus capitatus</i>	Pacific Ninebark	Ros-	-   -   C	- NP	Moist banks, n-facing slopes, mixed-conifer forest (May–Jul)	
<input type="radio"/> <i>Pinus coulteri</i>	Coulter Pine	Pin-	-   -   A2	- NP	Chaparral, lower mixed-conifer, mixed-hardwood forests -	
<input type="radio"/> <i>Pinus sabiniana</i>	Gray Pine	Pin-	-   -   -	- NP	Foothill woodland, n oak woodland, chaparral, infertile soils in mixed-conifer and hardwood forests -	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Platan-	-   -   B	- NP	Common. Streamsides, canyons, arroyos (Feb–Apr)	
<input type="radio"/> <i>Populus fremontii</i> subsp. <i>fremontii</i>	Fremont Cottonwood	Salic-	-   -   -	- NP	Scattered. Alluvial bottomland, streamsides (Mar–Apr)	
<input type="radio"/> <i>Prunus ilicifolia</i> subsp. <i>ilicifolia</i>	Holly-leaved Cherry	Ros-	-   -   C	- NP	Canyons, slopes, scrubland, woodland (Apr–May)	
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Ros-	-   -   -	- NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Fag-	-   -   A2	- NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Fag-	-   -   C	- NP	Dry slopes, chaparral (Feb–Apr)	

<b>Woody</b>	<b>Grouped by Growth Form</b>					<b>Woody</b>
<b>Quercus</b>	<b>Sorted Alphabetically by Scientific Name</b>					<b>Quercus</b>
✓ Scientific Name (JM93 if different)	Common Name	Family	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Fag-	-   -   -	-	NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Fag-	-   -   -	-	NP	Dry slopes, interior foothills, woodland (Apr–May)
<input type="radio"/> <i>Quercus durata</i> var. <i>durata</i>	Leather Oak	Fag-	-   -   A2	-	NP	Chaparral, generally serpentine (Apr–May)
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Fag-	-   -   -	-	NP	Slopes, valleys, woodland, conifer forest (Apr–May)
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Fag-	-   -   B	-	NP	Slopes, valleys, savanna (Mar–Apr)
<input type="radio"/> <i>Quercus palmeri</i>	Palmer Oak	Fag-	-   -   A2	-	NP	Uncommon. Rocky slopes, flats (Apr–May)
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Fag-	-   -   -	-	NP	Interior canyons, slopes, pine/oak woodland (Mar–May)
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Rhamn-	-   -   C	-	NP	Chaparral, montane forest (Mar–Jun)
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Grossulari-	-   -   -	-	NP	Forest openings, woodland (Feb–Mar)
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Grossulari-	-   -   -	-	NP	Common. Forest openings, chaparral (Feb–Apr)
<input type="radio"/> <i>Ribes sanguineum</i> var. <i>glutinosum</i>	Red-flowering Currant	Grossulari-	-   -   C	-	NP	Many habitats (Feb–Apr)
<input type="radio"/> <i>Rosa californica</i>	California Rose	Ros-	-   -   -	-	NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Ros-	-   -   -	-	NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)
<input type="radio"/> <i>Rubus armeniacus</i> (JM93: <i>R. discolor</i> )	Himalayan Blackberry	Ros-	-   -   -	H	ZP	Common. Disturbed areas, roadsides (Mar–Jun)
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Ros-	-   -   -	-	NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Ros-	-   -   -	-	NP	Open, disturbed areas (Mar–Jul)
<input type="radio"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Salic-	-   -   -	-	NP	Common. Floodplains, sandy gravel (Apr–May)
<input type="radio"/> <i>Salix laevigata</i>	Red Willow	Salic-	-   -   -	-	NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Salic-	-   -   -	-	NP	Common. Shores, marshes, meadows, etc (Jan–Jun)
<input type="radio"/> <i>Salix</i> sp.	Willow	Salic-	-   -   -	-	?P	- -
<input type="radio"/> <i>Sambucus nigra</i> subsp. <i>caerulea</i> (JM93: <i>S. mexicana</i> )	Blue Elderberry	Adox-	-   -   -	-	NP	Common. Streambanks, open places in forest (Mar–Sep)
<input type="radio"/> <i>Schinus molle</i>	Pepper Tree	Anacardi-	-   -   -	L	ZP	Washes, slopes, abandoned fields (Jun–Aug)
<input type="radio"/> <i>Senecio flaccidus</i> var. <i>douglasii</i>	Douglas' Threadleaf Ragwort	Aster-	-   -   A2	-	NP	Generally disturbed dry, open, sandy or rocky sites (Jun–Oct)
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Solan-	-   -   -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)

<b>Woody</b>	<b>Grouped by Growth Form</b>						<b>Woody</b>
<b><i>Symphoricarpos</i></b>	<b>Sorted Alphabetically by Scientific Name</b>						<b><i>Symphoricarpos</i></b>
<input checked="" type="checkbox"/> <b>Scientific Name (JM93 if different)</b>	<b>Common Name</b>	<b>Family</b>	<b>Status</b>	<b>Inv</b>	<b>OL</b>	<b>Habitat (Bloom)</b>	
<input type="checkbox"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Caprifoli-	-   -   -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="checkbox"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Caprifoli-	-   -   -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
<input type="checkbox"/> <i>Tamarix parviflora</i>	Smallflower Tamarisk	Tamaric-	-   -   -	H	ZP	Common. Washes, streambanks, slopes, roadsides (Mar–Apr)	
<input type="checkbox"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Anacardi-	-   -   -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
<input type="checkbox"/> <i>Umbellularia californica</i>	California Bay	Laur-	-   -   -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)	
<input type="checkbox"/> <i>Vitis californica</i>	California Wild Grape	Vit-	-   -   -	-	NP	Streamsides, springs, canyons (May–Jun)	

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Acer negundo* var. *californicum* (no var.)

*Adenostoma fasciculatum* (var. *fasciculatum*)

*Agoseris grandiflora* (var. *grandiflora*)

*Agoseris heterophylla* (var. *cryptopleura*)

*Agrostis viridis* (*Polypogon*)

*Alisma plantago-aquatica* (*A. triviale*)

*Amsinckia menziesii* var. *intermedia* (*A. intermedia*)

*Amsinckia menziesii* var. *menziesii* (no var.)

*Antirrhinum vexillo-calyculatum* ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

*Arabis breweri* var. *breweri* (*Boechera breweri* subsp. *breweri*)

*Arabis glabra* var. *glabra* (*Turritis glabra*)

*Arctostaphylos tomentosa* ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

*Aster chilensis* (*Symphyotrichum chilense*)

*Aster lentus* (*Symphyotrichum lentum*)

*Aster radulinus* (*Eurybia radulina*)

*Aster subulatus* var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

*Atriplex joaquiniana* (*A. joaquinana*)

*Atriplex lentiformis* ssp. *lentiformis* (no subsp.)

*Atriplex triangularis* (*A. prostrata*)

*Azolla mexicana* (*A. microphylla*)

*Baccharis douglasii* (*B. glutinosa*)

*Baccharis pilularis* (subsp. *consanguinea*)

*Baccharis salicifolia* (subsp. *salicifolia*)

*Balsamorhiza macrolepis* var. *macrolepis* (no var.)

*Blepharizonia plumosa* ssp. *plumosa* (no subsp.)

*Blepharizonia plumosa* ssp. *viscida* (*Blepharizonia laxa*)

*Boschniakia strobilacea* (*Kopsiopsis*)

*Bromus catharticus* (var. *catharticus*)

*Camissonia boothii* ssp. *decorticans* (*Eremothera*)

*Camissonia graciliflora* (*Tetrapteron graciliflorum*)

*Camissonia intermedia* (*Camissoniopsis*)

*Camissonia micrantha* (*Camissoniopsis*)

*Camissonia ovata* (*Taraxia*)

*Cardaria draba* (*Lepidium*)

*Carduus pycnocephalus* (subsp. *pycnocephalus*)

*Carex deweyana* ssp. *leptopoda* (*C. leptopoda*)

*Carex lanuginosa* (*C. pellita*)

*Carex ovalis* (*C. leporina*)

*Ceanothus thyrsoiflorus* (var. *thyrsoiflorus*)

*Centaurium davyi* (*Zeltnera*)

*Centaurium muehlenbergii* (*Zeltnera*)

*Cephalanthus occidentalis* var. *californicus* (no var.)

*Chamomilla suaveolens* (*Matricaria discoidea*)

*Chenopodium ambrosioides* (*Dysphania*)

*Chenopodium macrospermum* var. *halophilum* (no var.)

*Chenopodium multifidum* (*Dysphania multifida*)

*Chenopodium rubrum* (var. *rubrum*)

*Chrysothamnus nauseosus* ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

*Cirsium cymosum* (var. *cymosum*)

*Claytonia parviflora* ssp. *parviflora* (subsp. *utahensis*)

*Collinsia bartsifolia* var. *davidsonii* (var. *stricta*)

*Collinsia heterophylla* (var. *heterophylla*)

*Conyza bilbaoana* (*Erigeron sumatrensis*)

*Conyza bonariensis* (*Erigeron*)

*Conyza canadensis* (*Erigeron*)

*Conyza coulteri* (*Laennecia*)

*Corallorhiza maculata* (var. *maculata*)

*Cordylanthus mollis* ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

*Coronopus didymus* (*Lepidium didymum*)

*Corylus cornuta* var. *californica* (subsp. *californica*)

*Cotoneaster pannosa* (*C. pannosus*)

*Cryptantha decipiens* (*C. rattanii*)

*Cryptantha intermedia* (var. *intermedia*)

*Cryptantha muricata* (var. *muricata*)

*Cryptantha torreyana* (var. *pumila*)

*Cupressus macrocarpa* (*Hesperocyparis*)

*Cuscuta salina* var. *major* (*C. pacifica* var. *pacifica*)

*Cynara cardunculus* (subsp. *flavescens*)

*Danthonia californica* var. *californica* (no var.)

*Dicentra chrysantha* (*Ehrendorferia*)

*Disporum hookeri* (*Prosartes*)

*Disporum smithii* (*Prosartes*)

*Draba cuneifolia* var. *integrifolia* (no var.)

*Duchesnea indica* (no var.)

*Epilobium pygmaeum* (*E. campestre*)

*Eragrostis curvula* var. *curvula* (no var.)

*Erechtites glomerata* (*Senecio glomeratus*)

*Erechtites minima* (*Senecio minimus*)

*Eremocarpus setigerus* (*Croton*)

*Erigeron philadelphicus* (var. *philadelphicus*)

*Eriogonum umbellatum* var. *bahiiforme* (var. *smallianum*)

*Eriophyllum lanatum* var. *achillaeoides* (var. *achilleoides*)

*Eryngium aristulatum* var. *aristulatum* (*E. jepsonii*)

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

*Erysimum capitatum* var. *angustatum* (var. *capitatum*)

*Filago californica* (*Logfia filaginoides*)

*Filago gallica* (*Logfia*)

*Fritillaria affinis* var. *affinis* (no var.)

*Galium trifidum* var. *pacificum* (subsp. *columbianum*)

*Gastridium ventricosum* (*G. phleoides*)

*Gnaphalium bicolor* (*Pseudognaphalium biolettii*)

*Gnaphalium californicum* (*Pseudognaphalium*)

*Gnaphalium canescens* ssp. *beneolens*  
(*Pseudognaphalium beneolens*)

*Gnaphalium canescens* ssp. *microcephalum*  
(*Pseudognaphalium microcephalum*)

*Gnaphalium luteo-album* (*Pseudognaphalium luteoalbum*)

*Gnaphalium purpureum* (*Gamochaeta ustulata*)

*Gnaphalium ramosissimum* (*Pseudognaphalium*)

*Gnaphalium stramineum* (*Pseudognaphalium*)

*Grindelia camporum* var. *camporum* (no var.)

*Grindelia hirsutula* var. *davyi* (*G. camporum*)

*Grindelia hirsutula* var. *hirsutula* (no var.)

*Guillenia flavescens* (*Caulanthus*)

*Guillenia lasiophylla* (*Caulanthus lasiophyllus*)

*Hedera helix* ssp. *helix* (no subsp.)

*Heliotropium curassavicum* (var. *oculatum*)

*Hemizonia congesta* ssp. *congesta* (subsp. *lutescens*)

*Hemizonia corymbosa* ssp. *corymbosa* (*Deinandra corymbosa*)

*Hemizonia fitchii* (*Centromadia*)

*Hemizonia lobbii* (*Deinandra*)

*Hemizonia parryi* ssp. *congdonii* (*Centromadia*)

*Hemizonia parryi* ssp. *parryi* (*Centromadia*)

*Hemizonia pungens* ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

*Hemizonia pungens* ssp. *pungens* (*Centromadia*)

*Heracleum lanatum* (*H. maximum*)

*Herniaria hirsuta* ssp. *cinerea* (var. *cinerea*)

*Holodiscus discolor* (var. *discolor*)

*Hordeum jubatum* (subsp. *jubatum*)

*Horkelia californica* ssp. *californica* (var. *californica*)

*Horkelia californica* ssp. *dissita* (var. *elata*)

*Horkelia californica* ssp. *frondosa* (var. *frondosa*)

*Isopyrum occidentale* (*Enemion*)

*Iva axillaris* ssp. *robustior* (no subsp.)

*Juglans californica* var. *hindsii* (*J. hindsii*)

*Juncus balticus* (subsp. *ater*)

*Juncus effusus* var. *pacificus* (subsp. *pacificus*)

*Juncus lesueurii* (*J. lescurii*)

*Koeleria phleoides* (*K. gerardii*)

*Lagophylla ramosissima* ssp. *ramosissima* (no subsp.)

*Lasthenia californica* (subsp. *californica*)

*Lavatera cretica* (*Malva pseudolavatera*)

*Lepidium dictyotum* var. *dictyotum* (no var.)

*Lepidium latipes* var. *latipes* (no var.)

*Lepidium nitidum* var. *nitidum* (no var.)

*Lepidium nitidum* var. *oreganum* (no var.)

*Leptochloa fascicularis* (*L. fusca* subsp. *fascicularis*)

*Lessingia filaginifolia* var. *californica* (*Corethrogyne filaginifolia*)

*Lessingia filaginifolia* var. *filaginifolia*  
(*Corethrogyne filaginifolia*)

*Lewisia rediviva* (var. *rediviva*)

*Leymus condensatus* (*Elymus*)

*Leymus triticoides* (*Elymus*)

*Lilaea scilloides* (*Triglochin*)

*Limosella subulata* (*Limosella australis*)

*Linanthus acicularis* (*Leptosiphon*)

*Linanthus ambiguus* (*Leptosiphon*)

*Linanthus androsaceus* (*Leptosiphon*)

*Linanthus bicolor* (*Leptosiphon*)

*Linanthus ciliatus* (*Leptosiphon*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus dichotomus* (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

*Linanthus parviflorus* (*Leptosiphon*)

*Linaria canadensis* (*Nuttallanthus texanus*)

*Lolium multiflorum* (*Festuca perennis*)

*Lolium perenne* (*Festuca perennis*)

*Lolium temulentum* (*Festuca temulenta*)

*Lonicera hispidula* var. *vacillans* (no var.)

*Lotus humistratus* (*Acmispon brachycarpus*)

*Lotus micranthus* (*Acmispon parviflorus*)

*Lotus purshianus* var. *purshianus* (*Acmispon americanus* var. *americanus*)

*Lotus scoparius* var. *scoparius* (*Acmispon glaber* var. *glaber*)

*Lotus strigosus* (*Acmispon*)

*Lotus wrangelianus* (*Acmispon*)

*Luzula comosa* (var. *comosa*)

*Madia madioides* (*Anisocarpus*)

*Malacothamnus fasciculatus* (var. *nuttallii*)

*Marah fabaceus* (*M. fabacea*)

*Marah oreganus* (*M. oregana*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Melilotus alba* (*M. albus*)

*Melilotus indica* (*M. indicus*)

*Mentha spicata* var. *spicata* (no var.)

*Mimulus aurantiacus* (var. *aurantiacus*)

*Mirabilis californica* (*M. laevis* var. *crassifolia*)

*Monardella antonina* ssp. *antonina* (*M. villosa* subsp. *villosa*)

*Monardella douglasii* ssp. *douglasii* (no subsp.)

*Monardella villosa* ssp. *globosa* (subsp. *villosa*)

*Montia fontana* ssp. *chondrosperma* (no subsp.)

*Myrica californica* (*Morella*)

*Nassella cernua* (*Stipa*)

*Nassella lepida* (*Stipa*)

*Nassella pulchra* (*Stipa*)

*Osmorhiza chilensis* (*O. berteroi*)

*Oxalis albicans* ssp. *pilosa* (*O. pilosa*)

*Parvisedum pentandrum* (*Sedella pentandra*)

*Phacelia ramosissima* var. *latifolia* (no var.)

*Phacelia ramosissima* var. *ramosissima* (no var.)

*Phlox gracilis* (*Microsteris*)

*Phoradendron macrophyllum* (*P. serotinum* subsp. *macrophyllum*)

*Phoradendron villosum* (*P. serotinum* subsp. *tomentosum*)

*Phyla nodiflora* var. *incisa* (no var.)

*Phyla nodiflora* var. *nodiflora* (no var.)

*Picris echioides* (*Helminthotheca*)

*Piperia elegans* (subsp. *elegans*)

*Piptatherum miliaceum* (*Stipa miliacea* var. *miliacea*)

*Plagiobothrys canescens* (var. *canescens*)

*Plagiobothrys fulvus* (var. *campestris*)

*Plectritis brachystemon* (*P. congesta* subsp. *brachystemon*)

*Plectritis ciliosa* ssp. *ciliosa* (no subsp.)

*Plectritis ciliosa* ssp. *insignis* (no subsp.)

*Plectritis congesta* (subsp. *congesta*)

*Pleuropogon californicus* (var. *californicus*)

*Pluchea odorata* (var. *odorata*)

*Poa bulbosa* (subsp. *vivipara*)

*Polygonum amphibium* var. *emersum* (*Persicaria amphibia*)

*Polygonum amphibium* var. *stipulaceum* (*Persicaria amphibia*)

*Polygonum arenastrum* (*P. aviculare* subsp. *depressum*)

*Polygonum hydropiperoides* (*Persicaria*)

*Polygonum lapathifolium* (*Persicaria lapathifolia*)

*Polygonum persicaria* (*Persicaria maculosa*)

*Polygonum punctatum* (*Persicaria punctata*)

*Populus balsamifera* ssp. *trichocarpa* (*P. trichocarpa*)

*Potamogeton pectinatus* (*Stuckenia pectinata*)

*Potentilla glandulosa* ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

*Psilocarphus tenellus* var. *tenellus* (no var.)

*Ranunculus aquatilis* var. *capillaceus* (var. *diffusus*)

*Ranunculus californicus* (var. *californicus*)

*Ranunculus canus* (var. *canus*)

*Ranunculus occidentalis* (var. *occidentalis*)

*Ranunculus sceleratus* (var. *sceleratus*)

*Rhamnus californica* ssp. *californica* (*Frangula*)

*Rhamnus tomentella* ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

*Rhus trilobata* (*R. aromatica*)

*Ribes menziesii* (var. *menziesii*)

*Rorippa nasturtium-aquaticum* (*Nasturtium officinale*)

*Rorippa palustris* var. *occidentalis* (subsp. *palustris*)

*Rosa gymnocarpa* (var. *gymnocarpa*)

*Rubus discolor* (*R. armeniacus*)

*Rumex maritimus* (*R. fueginus*)

*Rumex salicifolius* var. *crassus* (*R. crassus*)

*Rumex salicifolius* var. *salicifolius* (no var.)

*Rumex salicifolius* var. *transitorius* (*R. transitorius*)

*Salicornia europaea* (*S. depressa*)

*Salicornia subterminalis* (*Arthrocnemum subterminale*)

*Salicornia virginica* (*S. pacifica*)

*Salix exigua* (var. *hindsiana*)

*Salix lucida* ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

*Sambucus mexicana* (*S. nigra* subsp. *caerulea*)

*Satureja douglasii* (*Clinopodium*)

*Saxifraga californica* (*Micranthes*)

*Scirpus acutus* var. *occidentalis* (*Schoenoplectus*)

*Scirpus americanus* (*Schoenoplectus*)

*Scirpus californicus* (*Schoenoplectus*)

*Scirpus cernuus* (*Isolepis*)

*Scirpus maritimus* (*Bolboschoenus maritimus* subsp. *paludosus*)

*Scirpus robustus* (*Bolboschoenus*)

*Scrophularia californica* ssp. *californica* (no subsp.)

*Scrophularia californica* ssp. *floribunda* (no subsp.)

*Senecio breweri* (*Packera*)

*Senecio mikanioides* (*Delairea odorata*)

*Silene californica* (*S. laciniata* subsp. *californica*)

---

## East Bay Regional Park District

### Wild Plant Name Changes

#### JM93 Scientific Name (JM2 name changes)

---

*Smilacina racemosa* (*Maianthemum racemosum*)

*Smilacina stellata* (*Maianthemum stellatum*)

*Solidago californica* (*S. velutina* subsp. *californica*)

*Solidago canadensis* ssp. *elongata* (*S. elongata*)

*Sparganium erectum* ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

*Sparganium eurycarpum* ssp. *eurycarpum* (var. *eurycarpum*)

*Spergula arvensis* ssp. *arvensis* (no subsp.)

*Spergularia media* ( var. *media*)

*Stachys ajugoides* var. *ajugoides* (no var.)

*Stachys ajugoides* var. *rigida* (*Stachys rigida* var. *quercetorum*)

*Streptanthus albidus* ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

*Stylomecon heterophylla* (*Papaver heterophyllum*)

*Taeniatherum caput-medusae* (*Elymus*)

*Thysanocarpus laciniatus* (var. *laciniatus*)

*Tolmiea menziesii* (*T. diplomenziesii*)

*Trifolium albopurpureum* var. *albopurpureum* (no var.)

*Trifolium albopurpureum* var. *dichotomum* (*T. dichotomum*)

*Trifolium albopurpureum* var. *olivaceum* (*T. olivaceum*)

*Trifolium barbigerum* var. *andrewsii* (*T. grayi*)

*Trifolium barbigerum* var. *barbigerum* (no var.)

*Trifolium gracilentum* var. *gracilentum* (no var.)

*Vicia americana* var. *americana* (subsp. *americana*)

*Vicia ludoviciana* var. *ludoviciana* (subsp. *ludoviciana*)

*Viola adunca* (var. *adunca*)

*Vulpia bromoides* (*Festuca*)

*Vulpia microstachys* var. *ciliata* (*Festuca microstachys*)

*Vulpia microstachys* var. *confusa* (*Festuca microstachys*)

*Vulpia microstachys* var. *microstachys* (*Festuca microstachys*)

*Vulpia microstachys* var. *pauciflora* (*Festuca microstachys*)

*Vulpia myuros* var. *hirsuta* (*Festuca myuros*)

*Vulpia myuros* var. *myuros* (*Festuca myuros*)

*Vulpia octoflora* var. *hirtella* (*Festuca octoflora*)

*Vulpia octoflora* var. *octoflora* (*Festuca octoflora*)

*Zigadenus fremontii* (*Toxicoscordion*)