

East Bay Regional Park District

Checklist of Wild Plants

Sorted Alphabetically by Family, Scientific Name



This is a comprehensive list of the wild plants reported to be found in the East Bay Regional Park District. The plants are sorted alphabetically by family, then by scientific name. This list includes the common name, growth form, status, invasiveness rating, origin, longevity, habitat, and bloom dates. EBRPD plant names that have changed since the 1993 Jepson Manual are listed alphabetically in an appendix, as well as changes in families for specific genera.

Column Heading	Description
	✓ Checklist column for marking off the plants you observe
Scientific Name	According to <i>The Jepson Manual: Vascular Plants of California, Second Edition (JM2)</i> and <i>eFlora</i> (ucjeps.berkeley.edu/IJM.html)
(JM93 if different)	If the scientific name used in the 1993 edition of <i>The Jepson Manual (JM93)</i> is different, the change is noted as (JM93: xxx)
Common Name	According to <i>JM2</i> and other references (not standardized)
Form	Growth Form: Fern-like (Ferns & Horsetails), Grass-like (Grasses & Grass-like), Herbaceous, and Woody
Status	Special status rating (if any), listed in 3 categories, divided by vertical bars (' '): Federal/California (Fed./Calif.) California Native Plant Society (CNPS) East Bay chapter of the CNPS (EBCNPS) Fed./Calif.: FE = Fed. Endangered, FT = Fed. Threatened, CE = Calif. Endangered, CR = Calif. Rare CNPS (online as of 2012-01-23): 1B = Rare, threatened or endangered in Calif, 3 = Review List, 4 = Watch List; 0.1 = Seriously endangered in California, 0.2 = Fairly endangered in California EBCNPS (online as of 2012-01-23): *A = Statewide listed rare; A1 = 2 East Bay regions or less; A1x = extirpated; A2 = 3-5 regions; B = 6-9 regions; C = watch list
Inv	California Invasive Plant Council Inventory (Cal-IPCI) Invasiveness rating: H = High, L = Limited, M = Moderate, N = Native
OL	Origin and Longevity. Origin: N = Native, Z = naturalized or W = Waif; followed by Longevity: A = Annual, B = Biennial, P = Perennial, or a combination (ie. ZAB = naturalized annual or biennial)
Habitat (Bloom)	Habitat description (Bloom date range) according to <i>JM2</i> and other sources

This document is a compilation of field surveys submitted by various sources. Readers are encouraged to report any corrections (including probable misidentifications), additions, suggested improvements, or any other feedback by emailing the District Botanist (Wilde Legard, wlegard@ebparks.org). The latest version of this checklist can be found at: <http://www.ebparks.org/stewardship/plants/checklist>.

East Bay Regional Park District

Checklist of Wild Plants

Adoxaceae		Sorted Alphabetically by Family, Scientific Name				Adoxaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
Adoxaceae						Adoxaceae
<input type="radio"/> <i>Sambucus nigra subsp. caerulea</i> (JM93: <i>S. mexicana</i>)	Blue Elderberry	Woody	- - -	- NP	Common. Streambanks, open places in forest (Mar–Sep)	
<input type="radio"/> <i>Sambucus racemosa var. racemosa</i>	Red Elderberry	Woody	- - A1	- NP	Moist places (May–Jul)	
<input type="radio"/> <i>Viburnum ellipticum</i>	Oval-leaved Viburnum	Woody	- 2.3 *A2	- NP	Chaparral, yellow-pine forest, generally n-facing slopes (Jun–Aug)	
Agavaceae						Agavaceae
<input type="radio"/> <i>Chlorogalum pomeridianum var. pomeridianum</i>	Wavyleaf Soap Plant	Herbaceous	- - -	- NP	Common. Open grassland, chaparral, woodland (May–Aug)	
Aizoaceae						Aizoaceae
<input type="radio"/> <i>Carpobrotus chilensis</i>	Sea Fig	Woody	- - -	M ZP	Common on coastal sandy shores (All year)	
<input type="radio"/> <i>Carpobrotus edulis</i>	Freeway Iceplant	Woody	- - -	H ZP	Common. Many coastal habitats, esp. sand (All year)	
<input type="radio"/> <i>Carpobrotus sp.</i>	Hottentot / Sea Fig	Woody	- - -	- ZP	- -	
<input type="radio"/> <i>Tetragonia tetragonioides</i>	New Zealand Spinach	Herbaceous	- - -	L ZA	Common. Sand dunes, bluffs, margins of coastal wetlands (Apr–Sep)	
<input type="radio"/> <i>Trianthema portulacastrum</i>	Horse-purslane	Herbaceous	- - A1	- NA	Uncommon. Moist or seasonally dry wetlands, disturbed areas (Jun–Nov)	
Alismataceae						Alismataceae
<input type="radio"/> <i>Alisma triviale</i> (JM93: <i>A. plantago-aquatica</i>)	Northern Water Plantain	Herbaceous	- - B	- NP	Ponds (Spring–fall)	
<input type="radio"/> <i>Echinodorus berteroi</i>	Upright Burhead	Herbaceous	- - A2	- NA	Ponds, ditches (Mid–summer–fall)	
<input type="radio"/> <i>Sagittaria latifolia</i>	Wappato / Tule Potato / Arrowhead	Herbaceous	- - A1	- NP	Ponds, slow streams (Jul–Aug)	
Alliaceae						Alliaceae
<input type="radio"/> <i>Allium amplexans</i>	Narrowleaf Onion	Herbaceous	- - A2	- NP	Clays including serpentine, open or wooded places (Apr–Jul)	
<input type="radio"/> <i>Allium bolanderi var. bolanderi</i>	Bolander's Onion	Herbaceous	- - A1	- NP	Uncommon. Rocky clays including serpentine (May–Aug)	
<input type="radio"/> <i>Allium falcifolium</i>	Sickle Leaf Onion	Herbaceous	- - A2	- NP	Common. Heavy clays including serpentine (Apr–Jun)	
<input type="radio"/> <i>Allium serra</i>	Jeweled Onion	Herbaceous	- - -	- NP	Common. Grassy slopes (Apr–May)	

Alliaceae		Sorted Alphabetically by Family, Scientific Name					Alliaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Allium triquetrum</i>	Three Cornered Leek	Herbaceous	- - -	-	ZP	Locally common. Shady ± disturbed places (Mar–Apr)	
<input type="radio"/> <i>Allium unifolium</i>	One-leaf Onion	Herbaceous	- - C	-	NP	Uncommon. Moist clay or serpentine, especially grassy streambanks (May–Jun)	
Amaranthaceae							Amaranthaceae
<input type="radio"/> <i>Amaranthus albus</i>	Tumbleweed	Herbaceous	- - -	-	ZA	Disturbed areas, roadsides, riverbanks, sandy places, agricultural fields (Jun–Oct)	
<input type="radio"/> <i>Amaranthus blitoides</i>	Procumbent Pigweed	Herbaceous	- - C	-	NA	Disturbed areas, roadsides, agricultural fields, sandy soil (Jul–Nov)	
<input type="radio"/> <i>Amaranthus californicus</i>	California Amaranth	Herbaceous	- - A2	-	NA	Seasonally moist flats, lake margins, disturbed areas (Jul–Oct)	
<input type="radio"/> <i>Amaranthus deflexus</i>	Low Amaranth	Herbaceous	- - -	-	ZA	Railroad right-of-ways, disturbed areas (May–Nov)	
<input type="radio"/> <i>Amaranthus powellii</i>	Powell's Amaranth	Herbaceous	- - A2	-	NA	Agricultural fields, disturbed areas (Jun–Oct)	
<input type="radio"/> <i>Amaranthus retroflexus</i>	Redroot Pigweed	Herbaceous	- - -	-	ZA	Common. Wet fields, roadside ditches, waste places, agricultural fields (Jun–Nov)	
<input type="radio"/> <i>Amaranthus sp.</i>	Amaranth	Herbaceous	- - -	-	- -		
Anacardiaceae							Anacardiaceae
<input type="radio"/> <i>Rhus aromatica</i> (JM93: <i>R. trilobata</i>)	Skunk Bush	Woody	- - C	-	NP	Slopes, washes, scrub (Mar–May)	
<input type="radio"/> <i>Schinus molle</i>	Pepper Tree	Woody	- - -	L	ZP	Washes, slopes, abandoned fields (Jun–Aug)	
<input type="radio"/> <i>Toxicodendron diversilobum</i>	Western Poison Oak	Woody	- - -	-	NP	Canyons, slopes, chaparral, coastal scrub, oak woodland (Apr–Jun)	
Apiaceae							Apiaceae
<input type="radio"/> <i>Angelica tomentosa</i>	California Wood Angelica	Herbaceous	- - -	-	NP	Generally wooded areas (Jun–Aug)	
<input type="radio"/> <i>Anthriscus caucalis</i>	Bur-chervil	Herbaceous	- - -	-	ZA	Generally shady places (Apr–Jun)	
<input type="radio"/> <i>Apiastrum angustifolium</i>	Wild Celery	Herbaceous	- - B	-	NA	Chaparral, coastal scrub (Mar–Apr)	
<input type="radio"/> <i>Apium graveolens</i>	Celery	Herbaceous	- - -	-	ZA	Wet places (May–Jul)	
<input type="radio"/> <i>Berula erecta</i>	Cutleaf Water-parsnip	Herbaceous	- - B	-	NP	Marshy areas, streams (Jul–Oct)	
<input type="radio"/> <i>Bowlesia incana</i>	Bowlesia	Herbaceous	- - -	-	NA	Shade of trees, rocks, shrubs (Mar–Apr)	
<input type="radio"/> <i>Cicuta douglasii</i>	Western Water-hemlock	Herbaceous	- - B	-	NP	Wet places, gen aquatic (Jun–Sep)	
<input type="radio"/> <i>Cicuta maculata</i> var. <i>bolanderi</i>	Bolander Water Hemlock	Herbaceous	- 2.1 *A1	-	NP	Coastal wetlands (Jul–Sep)	
<input type="radio"/> <i>Conium maculatum</i>	Poison Hemlock	Herbaceous	- - -	M	ZB	Common. Moist, esp disturbed places (Apr–Jul)	
<input type="radio"/> <i>Daucus pusillus</i>	Rattlesnake Weed	Herbaceous	- - -	-	NA	Rocky or sandy places (Apr–Jun)	
<input type="radio"/> <i>Eryngium articulatum</i>	Beethistle	Herbaceous	- - A1	-	NP	Seasonally wet areas on lake and stream margins, marshes (Jun–Aug)	

Apiaceae		Sorted Alphabetically by Family, Scientific Name					Apiaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Eryngium jepsonii</i> (JM93: <i>E. aristulatum</i> var. <i>aristulatum</i>)	Button Celery	Herbaceous	- - -	-	NP	Moist clay soil, vernal pools, lake shores, drying lakes, wet depressions (Apr–Aug)	
<input type="radio"/> <i>Eryngium</i> sp.	Eryngo	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Foeniculum vulgare</i>	Fennel	Herbaceous	- - -	H	ZP	Roadsides, disturbed sites (May–Sep)	
<input type="radio"/> <i>Heracleum maximum</i> (JM93: <i>H. lanatum</i>)	Cow Parsnip	Herbaceous	- - -	-	NP	Moist places, wooded or open (Apr–Jul)	
<input type="radio"/> <i>Lilaeopsis masonii</i>	Mason's Lilaeopsis	Herbaceous	CR 1B.1 *A2	-	NP	Intertidal marshes, streambanks. (Jun–Aug)	
<input type="radio"/> <i>Lomatium californicum</i>	Celery Weed	Herbaceous	- - B	-	NP	Woodland, brushy slopes (Apr–Jun)	
<input type="radio"/> <i>Lomatium caruifolium</i> var. <i>caruifolium</i>	Alkali Desertparsley	Herbaceous	- - B	-	NP	Wet, clay depressions, open grassland (Mar–May)	
<input type="radio"/> <i>Lomatium dasycarpum</i> subsp. <i>dasycarpum</i>	Woolly Fruited Desertparsley	Herbaceous	- - -	-	NP	Rocky (gen serpentine), chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Lomatium macrocarpum</i>	Sheep Biscuit Root	Herbaceous	- - -	-	NP	Generally serpentine rocky slopes in chaparral or woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium nudicaule</i>	Swollen-stem Biscuit Root	Herbaceous	- - -	-	NP	Rocky slopes, flats, gen pine woodland (Apr–Jun)	
<input type="radio"/> <i>Lomatium</i> sp.	Biscuit Root	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Lomatium utriculatum</i>	Bladder Parsnip	Herbaceous	- - -	-	NP	Open grassy slopes, meadows, woodland (Feb–May)	
<input type="radio"/> <i>Oenanthe sarmentosa</i>	Water Parsley	Herbaceous	- - -	-	NP	Streams, marshes, ponds, gen aquatic (Jun–Oct)	
<input type="radio"/> <i>Osmorhiza berteroi</i> (JM93: <i>O. chilensis</i>)	Sweet-cicely	Herbaceous	- - -	-	NP	Conifer forest, woodland, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Osmorhiza brachypoda</i>	California Sweet Cicely	Herbaceous	- - B	-	NP	Moist ravines, conifer forest, woodland (Mar–May)	
<input type="radio"/> <i>Perideridia californica</i>	California Yampah	Herbaceous	- - A2	-	NP	Damp soil by streams (Apr–May)	
<input type="radio"/> <i>Perideridia kelloggii</i>	Kellogg Yampah	Herbaceous	- - -	-	NP	Open grassland, serpentine outcrops (Jul–Aug)	
<input type="radio"/> <i>Perideridia oregana</i>	Oregon Yampah	Herbaceous	- - A2	-	NP	Open flats or slopes, pine/oak woodland (Jul–Aug)	
<input type="radio"/> <i>Perideridia</i> sp.	Yampah	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Sanicula arctopoides</i>	Footsteps-of-spring	Herbaceous	- - A1x	-	NP	Open coastal bluffs, headlands, dunes (Feb–May)	
<input type="radio"/> <i>Sanicula bipinnata</i>	Poison Sanicle	Herbaceous	- - -	-	NP	Open grassland or pine/oak woodland (Apr–May)	
<input type="radio"/> <i>Sanicula bipinnatifida</i>	Purple Sanicle	Herbaceous	- - -	-	NP	Open grassland, gen on serpentine, or pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Sanicula crassicaulis</i>	Pacific Woodland Sanicle	Herbaceous	- - -	-	NP	Open slopes, ravines, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula laciniata</i>	Fringe-leaf / Coast Sanicle	Herbaceous	- - A2	-	NP	Coastal, open or shrubby slopes, woodland (Mar–May)	
<input type="radio"/> <i>Sanicula tuberosa</i>	Turkey Pea Sanicle	Herbaceous	- - C	-	NP	Open gravelly meadows, chaparral, woodland, pine forest (Mar–Jul)	
<input type="radio"/> <i>Scandix pecten-veneris</i>	Venus' Needle	Herbaceous	- - -	-	ZA	Grassy slopes, roadsides (Apr–Jun)	
<input type="radio"/> <i>Sium suave</i>	Hemlock Waterparsnip	Herbaceous	- - A1	-	NP	Swamps, marshes, streambanks (Jul–Aug)	

Apiaceae		Sorted Alphabetically by Family, Scientific Name					Apiaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Tauschia hartwegii</i>	Hartweg's Tauschia	Herbaceous	- - C	-	NP	Chaparral, pine/oak woodland (Mar–May)	
<input type="radio"/> <i>Torilis arvensis</i>	Tall Sock-destroyer	Herbaceous	- - -	M	ZA	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Torilis nodosa</i>	Short Sock-destroyer	Herbaceous	- - -	-	ZA	Disturbed places (Apr–Jun)	
<input type="radio"/> <i>Yabea microcarpa</i>	California Hedge Parsley	Herbaceous	- - C	-	NA	Grassy slopes, dunes, chaparral, woodland (Apr–Jun)	
Apocynaceae							Apocynaceae
<input type="radio"/> <i>Apocynum androsaemifolium</i>	Bitter Dogbane	Herbaceous	- - B	-	NP	Open slopes, rocky places, with conifers, chaparral (May–Oct)	
<input type="radio"/> <i>Apocynum cannabinum</i>	Indian Hemp	Herbaceous	- - A2	-	NP	Moist places, near streams, springs, or as weed in orchards (Apr–Oct)	
<input type="radio"/> <i>Asclepias californica</i>	California Milkweed	Herbaceous	- - B	-	NP	Flats, grassy or brushy hillsides (Apr–Jul)	
<input type="radio"/> <i>Asclepias cordifolia</i>	Purple Milkweed	Herbaceous	- - A1	-	NP	Rocky slopes, talus, woodland, chaparral, lava flows (May–Jul)	
<input type="radio"/> <i>Asclepias fascicularis</i>	Narrow-leaf Milkweed	Herbaceous	- - -	-	NP	Dry ground, valleys, foothills (May–Oct)	
<input type="radio"/> <i>Asclepias speciosa</i>	Showy Milkweed	Herbaceous	- - A2	-	NP	Many habitats incl fields, roadsides (May–Sep)	
<input type="radio"/> <i>Vinca major</i>	Greater Periwinkle	Herbaceous	- - -	M	ZP	Coastal bluffs, sheltered places, esp along stream beds (Mar–Jun(Jan))	
Aquifoliaceae							Aquifoliaceae
<input type="radio"/> <i>Ilex aquifolium</i>	English Holly	Woody	- - -	M	ZP	Cool, wooded areas (May–Jun)	
Araceae							Araceae
<input type="radio"/> <i>Zantedeschia aethiopica</i>	Calla-lily	Herbaceous	- - -	L	ZP	Disturbed areas, near former habitations (Mar–Jun)	
Araliaceae							Araliaceae
<input type="radio"/> <i>Aralia californica</i>	Elk Clover	Herbaceous	- - B	-	NP	Moist shade, canyons, stream sides (Jun–Aug)	
<input type="radio"/> <i>Hedera helix (JM93: subsp. helix)</i>	English Ivy	Woody	- - -	H	ZP	Woodland, open, disturbed areas (Aug–Nov)	
<input type="radio"/> <i>Hydrocotyle ranunculoides</i>	Floating Marsh Pennywort	Herbaceous	- - A1	-	NP	Lake margins, ponds, slow-moving streams (Mar–Aug)	
<input type="radio"/> <i>Hydrocotyle sp.</i>	Water / Marsh Pennywort	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Hydrocotyle verticillata</i>	Whorled Marsh Pennywort	Herbaceous	- - A2	-	NP	Lake margins, ponds, slow-moving streams, canals, seeps, springs, marshes (Apr–Sep)	
Arecaceae							Arecaceae
<input type="radio"/> <i>Phoenix canariensis</i>	Canary Island Date Palm	Woody	- - -	L	ZP	Uncommon. Near development, disturbed areas (Oct–Apr)	
Aristolochiaceae							Aristolochiaceae
<input type="radio"/> <i>Aristolochia californica</i>	Dutchman's Pipe	Herbaceous	- - C	-	NP	Streamsides, forest, chaparral (Jan–Apr)	

Aristolochiaceae		Sorted Alphabetically by Family, Scientific Name				Aristolochiaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Asarum caudatum</i>	Wild Ginger	Herbaceous	- - A2	-	NP	Moist places in forest (Mar–Aug)
Asteraceae						Asteraceae
<input type="radio"/> <i>Achillea millefolium</i>	Yarrow	Herbaceous	- - -	-	NP	Many habitats (Apr–Sep)
<input type="radio"/> <i>Achyraea mollis</i>	Blow Wives	Herbaceous	- - -	-	NA	Common. Grassy sites, often clay soils (Mar–Jun)
<input type="radio"/> <i>Agoseris grandiflora</i> var. <i>grandiflora</i> (JM93: no var.)	Giant Native Dandelion	Herbaceous	- - -	-	NP	Grassland, scrub, woodland (Apr–Jul)
<input type="radio"/> <i>Agoseris heterophylla</i> var. <i>cryptopleura</i> (JM93: no var.)	Annual Native Dandelion	Herbaceous	- - -	-	NA	Many open habitats (May–Jun)
<input type="radio"/> <i>Agoseris retrorsa</i>	Spearleaf Native Dandelion	Herbaceous	- - B	-	NP	Scrub, oak woodland, conifer forest (Apr–Aug)
<input type="radio"/> <i>Agoseris</i> sp.	Agoseris	Herbaceous	- - -	-	NP	--
<input type="radio"/> <i>Ambrosia chamissonis</i>	Beach Bur-sage	Herbaceous	- - A2	-	NP	Beaches, dunes (Jul–Oct)
<input type="radio"/> <i>Ambrosia psilostachya</i>	Western Ragweed	Herbaceous	- - -	-	NP	Common. Roadsides, dry fields (Jun–Nov)
<input type="radio"/> <i>Anaphalis margaritacea</i>	Pearly Everlasting	Herbaceous	- - -	-	NP	Woodland, disturbed places (Jul–Oct)
<input type="radio"/> <i>Ancistrocarphus filagineus</i>	Woolly Fishhooks	Herbaceous	- - B	-	NA	Bare or grassy, often serpentine or clay, drainages, road beds, burns, vernal moist sites (Mar–Jun)
<input type="radio"/> <i>Anisocarpus madioides</i> (JM93: <i>Madia</i>)	Woodland Tarweed	Herbaceous	- - A2	-	NP	Moist forest, woodland (Apr–Sep)
<input type="radio"/> <i>Anthemis cotula</i>	Mayweed	Herbaceous	- - -	-	ZA	Common. Disturbed areas, fields, coastal dunes, chaparral, oak woodland (Apr–Aug)
<input type="radio"/> <i>Argyranthemum foeniculaceum</i>	Canary Islands Marguerite	Woody	- - -	-	ZP	Escape from cult in disturbed coastal areas (Mar–Aug)
<input type="radio"/> <i>Arnica discoidea</i>	Rayless Arnica	Herbaceous	- - B	-	NP	Chaparral, foothill woodland (May–Jul)
<input type="radio"/> <i>Artemisia californica</i>	California Sagebrush	Woody	- - -	-	NP	Coastal scrub, chaparral, open woodland (Aug–Nov)
<input type="radio"/> <i>Artemisia douglasiana</i>	Mugwort	Herbaceous	- - -	-	NP	Common. Open to shady areas, often in drainages (May–Nov)
<input type="radio"/> <i>Artemisia dracunculus</i>	Wild Tarragon	Herbaceous	- - A1	-	NP	Common. Many habitats, especially disturbed sites (Aug–Oct)
<input type="radio"/> <i>Aster</i> sp.	Aster	Herbaceous	- - -	-	NP	--
<input type="radio"/> <i>Baccharis glutinosa</i> (JM93: <i>B. douglasii</i>)	Marsh Baccharis	Herbaceous	- - -	-	NP	Coastal freshwater and saltwater marshes, streambanks (Jul–Oct)
<input type="radio"/> <i>Baccharis pilularis</i> subsp. <i>consanguinea</i> (JM93: no subsp.)	Coyote Brush	Woody	- - -	-	NP	Coastal bluffs, woodland, grassland, disturbed sites, occ on serpentine (Jul–Dec)
<input type="radio"/> <i>Baccharis salicifolia</i> subsp. <i>salicifolia</i> (JM93: no subsp.)	Mule Fat	Woody	- - -	-	NP	Riparian woodland, canyon bottoms, disturbed sites, often forming thickets (All year)
<input type="radio"/> <i>Balsamorhiza deltoidea</i>	Balsamroot	Herbaceous	- - A2	-	NP	Common. Grassy slopes, open forests, shrubby areas (Apr–Jul)

Asteraceae		Sorted Alphabetically by Family, Scientific Name				Asteraceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Balsamorhiza macrolepis</i> (JM93: var. <i>macrolepis</i>)	Bigscale Balsamroot	Herbaceous	- 1B.2 *A1	-	NP	Open grassy or rocky slopes, valleys (Mar–Jul)	
<input type="radio"/> <i>Bellis perennis</i>	English Daisy	Herbaceous	- - -	-	ZP	Damp, grassy areas (Dec–Sep)	
<input type="radio"/> <i>Bidens frondosa</i>	Sticktight	Herbaceous	- - B	-	NA	Uncommon. Damp soil, esp disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Bidens laevis</i>	Bur-marigold	Herbaceous	- - A1	-	NAP	Freshwater wetlands (Aug–Nov)	
<input type="radio"/> <i>Blennosperma nanum</i> var. <i>nanum</i>	Glue-seed	Herbaceous	- - C	-	NA	Open, grassy areas, often margins of seeps or vernal pools (Jan–May)	
<input type="radio"/> <i>Blepharizonia laxa</i> (JM93: <i>B. plumosa</i> subsp. <i>viscida</i>)	Stickyleaf Big Tarweed	Herbaceous	- - A2	-	NA	Openings in woodland, chaparral, grassland (Jul–Nov)	
<input type="radio"/> <i>Blepharizonia plumosa</i> (JM93: subsp.)	Big Tarplant	Herbaceous	- 1B.1 *A2	-	NA	Dry slopes in grassland (Jul–Nov)	
<input type="radio"/> <i>Brickellia californica</i>	California Brickellbush	Woody	- - B	-	NP	Dry, rocky hillsides, canyons, sea bluffs (Jul–Dec)	
<input type="radio"/> <i>Calycadenia multiglandulosa</i>	Sticky Rosinweed	Herbaceous	- - A1	-	NA	Common. Gen dry, open valleys, hillsides, rocky ridges (May–Oct)	
<input type="radio"/> <i>Calycadenia truncata</i>	Rosinweed	Herbaceous	- - A2	-	NA	Common. Dry, open hillsides, rocky ridges, talus, grassland, openings in foothill woodland, scrub (May–Oct)	
<input type="radio"/> <i>Carduus pycnocephalus</i> subsp. <i>pycnocephalus</i> (JM93: no subsp.)	Italian Thistle	Herbaceous	- - -	M	ZA	Roadsides, pastures, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Carduus tenuiflorus</i>	Slender Flower Thistle	Herbaceous	- - -	L	ZAB	Roadsides, pastures, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Centaurea calcitrapa</i>	Purple Star-thistle	Herbaceous	- - -	M	ZAB	Pastures, disturbed places (Apr–Nov)	
<input type="radio"/> <i>Centaurea melitensis</i>	Tocalote	Herbaceous	- - -	M	ZA	Disturbed fields, open woodland (Apr–Jul)	
<input type="radio"/> <i>Centaurea solstitialis</i>	Yellow Star-thistle	Herbaceous	- - -	H	ZA	Invasive, roadsides, disturbed grassland or woodland (May–Oct)	
<input type="radio"/> <i>Centromadia fitchii</i> (JM93: <i>Hemizonia</i>)	Fitch Spikeweed	Herbaceous	- - -	-	NP	Grassland, ± alkaline flats, vernal pools, woodland, disturbed sites, serpentine (May–Nov)	
<input type="radio"/> <i>Centromadia parryi</i> subsp. <i>congdonii</i> (JM93: <i>Hemizonia</i>)	Congdon's Tarplant	Herbaceous	- 1B.2 *A2	-	NA	Terraces, swales, floodplains, grassland, disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Centromadia parryi</i> subsp. <i>parryi</i> (JM93: <i>Hemizonia</i>)	Parry Spikeweed	Herbaceous	- - -	-	NA	Grassland, coastal salt marshes, alkaline springs, seeps (Jun–Oct)	
<input type="radio"/> <i>Centromadia pungens</i> subsp. <i>pungens</i> (JM93: <i>Hemizonia</i> , subsp. <i>maritima</i> , <i>pungens</i>)	Common Spikeweed	Herbaceous	- - -	-	NA	Grassland, saltbush scrub, disturbed sites (Apr–Nov)	
<input type="radio"/> <i>Chaenactis glabriuscula</i> var. <i>megacephala</i>	Big-headed Yellow Pincushion	Herbaceous	- - A1	-	NA	Dry, often sandy slopes, openings in chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Cichorium intybus</i>	Chicory	Herbaceous	- - -	-	ZP	Common. Roadsides, disturbed places (Apr–Oct)	
<input type="radio"/> <i>Cirsium brevistylum</i>	Indian Thistle	Herbaceous	- - B	-	NA	Moist places (Mar–Aug)	
<input type="radio"/> <i>Cirsium cymosum</i> var. <i>cymosum</i> (JM93: no var.)	Peregrine Thistle	Herbaceous	- - B	-	NP	Scrubland, woodland, open forest, meadows, occ serpentine (Jun–Jul)	

Asteraceae		Sorted Alphabetically by Family, Scientific Name					Asteraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Cirsium occidentale</i> var. <i>venustum</i>	Venus Thistle	Herbaceous	- - -	-	NB	Disturbed areas, grassland, woodland (May–Jul)	
<input type="radio"/> <i>Cirsium quercetorum</i>	Brownie Thistle	Herbaceous	- - A2	-	NP	Open places, grassland, woodland (Apr–Aug)	
<input type="radio"/> <i>Cirsium</i> sp.	Thistle	Herbaceous	- - -	-	-	-	
<input type="radio"/> <i>Cirsium vulgare</i>	Bull Thistle	Herbaceous	- - -	M	ZB	Common. Disturbed areas (May–Oct)	
<input type="radio"/> <i>Corethrogyne filaginifolia</i> (JM93: <i>Lessingia</i> , var. <i>filaginifolia</i>)	Common California-aster	Herbaceous	- - C	-	NP	Coastal scrub, chaparral, grassland, foothill woodland, forest (Jul–Nov)	
<input type="radio"/> <i>Cotula australis</i>	Australian Brass Buttons	Herbaceous	- - -	-	ZP	Common. Disturbed areas (Jan–May)	
<input type="radio"/> <i>Cotula coronopifolia</i>	Brass-buttons	Herbaceous	- - -	L	ZP	Common. Saline and freshwater marshes, mud flats (Mar–Dec)	
<input type="radio"/> <i>Crepis vesicaria</i> subsp. <i>taraxacifolia</i>	Dandelion-leaf Hawksbeard	Herbaceous	- - -	-	ZAB	Sandy clearings, hillsides, disturbed places (Feb–Oct)	
<input type="radio"/> <i>Crocidium multicaule</i>	Spring Gold	Herbaceous	- - A1	-	NA	Sandy soils, grassland, open woodland (Feb–Jun)	
<input type="radio"/> <i>Cynara cardunculus</i> subsp. <i>flavescens</i> (JM93: no subsp.)	Artichoke Thistle	Herbaceous	- - -	M	ZP	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Deinandra corymbosa</i> (JM93: <i>Hemizonia</i> , subsp. <i>corymbosa</i>)	Coast Spikeweed	Herbaceous	- - A2	-	NA	Grassland, openings in scrub or woodland, dunes, disturbed sites (Mar–Nov)	
<input type="radio"/> <i>Deinandra lobbii</i> (JM93: <i>Hemizonia</i>)	Lobb Tarweed	Herbaceous	- - B	-	NA	Grassland, open woodland, sagebrush scrub, disturbed areas (May–Dec)	
<input type="radio"/> <i>Delairea odorata</i> (JM93: <i>Senecio mikanioides</i>)	German Ivy	Herbaceous	- - -	H	ZP	Shady, ± disturbed places, riparian woodland, coastal scrub (Nov–Mar)	
<input type="radio"/> <i>Dittrichia graveolens</i>	Stinkwort	Herbaceous	- - -	M	ZA	Disturbed areas (Sep–Nov)	
<input type="radio"/> <i>Eclipta prostrata</i>	False Daisy	Herbaceous	- - A1	-	NA	Damp places (All year)	
<input type="radio"/> <i>Ericameria arborescens</i>	Golden-fleece	Woody	- - A2	-	NP	Woodland, open forest, chaparral, especially after fire (Aug–Nov)	
<input type="radio"/> <i>Ericameria linearifolia</i>	Interior Goldenbush	Woody	- - -	-	NP	Dry slopes, valleys, foothill and desert woodland, saltbush and creosote-bush scrub (Mar–May)	
<input type="radio"/> <i>Ericameria nauseosa</i> var. <i>mohavensis</i> (JM93: <i>Chysothamnus nauseosus</i> subsp. <i>mohavensis</i>)	Mojave Rabbitbrush	Woody	- - A2	-	NP	Common. Dry scrub (Aug–Oct)	
<input type="radio"/> <i>Erigeron bonariensis</i> (JM93: <i>Conyza</i>)	Flax-leaved Horseweed	Herbaceous	- - -	-	ZA	Disturbed sites (All year)	
<input type="radio"/> <i>Erigeron canadensis</i> (JM93: <i>Conyza</i>)	Horseweed	Herbaceous	- - -	-	ZA	Disturbed places (All year)	
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>foliosus</i>	Leafy Fleabane	Herbaceous	- - -	-	NP	Open, rocky grassland, chaparral, forest (May–Aug)	
<input type="radio"/> <i>Erigeron foliosus</i> var. <i>franciscensis</i>	Francis Leafy Fleabane	Herbaceous	- - -	-	NP	Grassy dunes, chaparral, oak woodland (May–Oct)	
<input type="radio"/> <i>Erigeron petrophilus</i> var. <i>petrophilus</i>	Rock-loving Fleabane	Herbaceous	- - A2	-	NP	Rocky foothills to montane forest, sometimes on serpentine (May–Sep)	
<input type="radio"/> <i>Erigeron philadelphicus</i> var. <i>philadelphicus</i> (JM93: no var.)	Philadelphia Fleabane	Herbaceous	- - C	-	NBP	Streamsides, other moist habitats (May–Jun)	
<input type="radio"/> <i>Erigeron</i> sp. (JM93: <i>Conyza</i>)	Fleabane Daisy	Herbaceous	- - -	-	-	-	

Asteraceae		Sorted Alphabetically by Family, Scientific Name					Asteraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
○ <i>Erigeron sumatrensis</i> (JM93: <i>Conyza bilbaoana</i>)	Tropical Horseweed	Herbaceous	- - -	-	ZA	Disturbed sites (All year)	
○ <i>Eriophyllum confertiflorum</i> var. <i>confertiflorum</i>	Golden-yarrow	Woody	- - -	-	NP	Many dry habitats (Apr–Aug)	
○ <i>Eriophyllum jepsonii</i>	Jepson's Woolly Sunflower	Woody	- 4.3 *A2	-	NP	Dry oak woodland (Apr–Jun)	
○ <i>Eriophyllum lanatum</i> var. <i>achilleoides</i> (JM93: var. <i>achillaeoides</i>)	Common Woolly Sunflower	Herbaceous	- - C	-	NP	Dry, often rocky sites, chaparral, forest (May–Jul)	
○ <i>Eriophyllum staechadifolium</i>	Seaside Woolly Sunflower	Woody	- - A2	-	NP	Dunes, sea bluffs, coastal scrub, canyons (Apr–Sep)	
○ <i>Eurybia radulina</i> (JM93: <i>Aster radulinus</i>)	Broadleaf Aster	Herbaceous	- - -	-	NP	Dry forest, oak/pine woodland, brushy slopes (Jul–Sep)	
○ <i>Euthamia occidentalis</i>	Western Goldenrod	Herbaceous	- - -	-	NP	Marshes, streambanks, meadows (Jul–Nov)	
○ <i>Gamochaeta ustulata</i> (JM93: <i>Gnaphalium purpureum</i>)	Purple Cudweed	Herbaceous	- - -	-	NA	Dunes, bluffs, fields, disturbed sites (Apr–Jul)	
○ <i>Gnaphalium</i> or <i>Pseudognaphalium</i> sp. (JM93: <i>Gnaphalium</i> sp.)	Everlasting Cudweed	Herbaceous	- - -	-	--		
○ <i>Gnaphalium palustre</i>	Lowland Cudweed	Herbaceous	- - -	-	NA	Arroyos, sandy streambeds, pond edges, potholes (May–Oct)	
○ <i>Grindelia camporum</i> (JM93: var. <i>camporum</i> , <i>G. hirsutula</i> var. <i>davyi</i>)	Great Valley Gumplant	Herbaceous	- - -	-	NP	Sandy or saline bottomland, roadsides (May–Nov)	
○ <i>Grindelia hirsutula</i> (JM93: var. <i>hirsutula</i>)	Hairy Gumplant	Herbaceous	- - C	-	NP	Sandy, clay, or serpentine slopes or roadsides (Apr–Jun)	
○ <i>Grindelia</i> sp.	Gumplant	Herbaceous	- - -	-	NP	--	
○ <i>Grindelia stricta</i> var. <i>angustifolia</i>	Marsh Gumplant	Woody	- - C	-	NP	Tidal wetlands (May–Dec)	
○ <i>Grindelia stricta</i> var. <i>platyphylla</i>	Pacific Gumplant	Herbaceous	- - A1	-	NP	Coastal bluffs, dunes (All year)	
○ <i>Gutierrezia californica</i>	California Matchweed	Woody	- - C	-	NP	Grassland, arid woodland and shrubland, serpentine (Jul–Nov)	
○ <i>Helenium bigelovii</i>	Bigelow Sneezeweed	Herbaceous	- - A2	-	NP	Wet meadows, marshes, bogs, fens, streambanks, lake margins (Jul–Aug)	
○ <i>Helenium puberulum</i>	Rosilla	Herbaceous	- - -	-	NB	Streambanks, seepage areas, lake margins (Jun–Aug)	
○ <i>Helianthella castanea</i>	Diablo Helianthella	Herbaceous	- 1B.2 *A2	-	NP	Open, grassy sites (Apr–Jun)	
○ <i>Helianthus annuus</i>	Common Sunflower	Herbaceous	- - -	-	NP	Disturbed areas, scrub, grassland, many other habitats (Jun–Oct)	
○ <i>Helianthus californicus</i>	California Sunflower	Herbaceous	- - C	-	NP	Springs, marshes, streambanks, canyons (Jul–Oct)	
○ <i>Helminthotheca echioides</i> (JM93: <i>Picris</i>)	Bristly Ox-tongue	Herbaceous	- - -	L	ZAB	Common. Disturbed areas (All year)	
○ <i>Hemizonia congesta</i> subsp. ?	Yellow Hayfield Tarweed	Herbaceous	- - -	-	NA	--	

Asteraceae		Sorted Alphabetically by Family, Scientific Name				Asteraceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
○ <i>Hemizonia congesta</i> subsp. <i>lutescens</i> (JM93: subsp. <i>congesta</i>)	Yellow Hayfield Tarweed	Herbaceous	- - C	-	NA	Grassland, barrens, openings in chaparral and woodland, often on serpentine (Apr–Dec)	
○ <i>Hemizonia congesta</i> subsp. <i>luzulifolia</i>	Hayfield Tarweed	Herbaceous	- - -	-	NP	Disturbed, open, or grassy sites, often clayey soils, serpentine (Mar–Dec)	
○ <i>Hemizonia</i> , <i>Centromadia</i> , or <i>Deinandra</i> sp. (JM93: <i>Hemizonia</i> sp.)	Spikeweed	Herbaceous	- - -	-	NAP	--	
○ <i>Hesperevax acaulis</i> var. <i>ambusticola</i>	Dwarf Fire Evax	Herbaceous	- - A2	-	NA	Clearings, barren slopes, burns (Apr–Jun)	
○ <i>Hesperevax sparsiflora</i> var. <i>sparsiflora</i>	Erect Hesperevax	Herbaceous	- - -	-	NA	Common. Open, clay and/or rocky, generally serpentine soil (Mar–Jun)	
○ <i>Heterotheca grandiflora</i>	Telegraph Weed	Herbaceous	- - -	-	NA	Disturbed areas, dry streambeds, sand dunes (Jun–Oct(± all year))	
○ <i>Heterotheca oregona</i> var. <i>rudis</i>	Inland Oregon Goldenaster	Herbaceous	- - A1	-	NP	Seasonally dry streambed (Jul–Oct)	
○ <i>Heterotheca oregona</i> var. <i>scaberrima</i>	Sticky Oregon Goldenaster	Herbaceous	- - A1	-	NP	Seasonally dry streambeds (Jul–Oct)	
○ <i>Heterotheca sessiliflora</i> subsp. ?	Sessileflower Goldenaster	Herbaceous	- - -	-	NP	--	
○ <i>Heterotheca sessiliflora</i> subsp. <i>bolanderi</i>	Bolander Goldenaster	Herbaceous	- - C	-	NP	Dunes, headlands, grassy coastal slopes (Jun–Sep)	
○ <i>Heterotheca sessiliflora</i> subsp. <i>echioides</i>	Bristly Goldenaster	Herbaceous	- - -	-	NP	Grassland, scrub, woodland, open forest, disturbed sites (Jul–Oct)	
○ <i>Heterotheca</i> sp.	Goldenaster	Herbaceous	- - -	-	NP	--	
○ <i>Hieracium albiflorum</i>	White Hawkweed	Herbaceous	- - B	-	NP	Forest (May–Sep)	
○ <i>Holocarpha heermannii</i>	Heermann Tarplant	Herbaceous	- - -	-	NA	Grassland (May–Nov)	
○ <i>Holocarpha macradenia</i>	Santa Cruz Tarplant	Herbaceous	FT/CE 1B.1 *A1	-	NA	Grassy areas, clay soil (Jun–Nov)	
○ <i>Holocarpha obconica</i>	San Joaquin Tarplant	Herbaceous	- - B	-	NA	Grassland (Apr–Nov)	
○ <i>Holocarpha</i> sp.	Tarplant	Herbaceous	- - -	-	NA	--	
○ <i>Holocarpha virgata</i> subsp. <i>virgata</i>	Sticky Tarplant	Herbaceous	- - -	-	NA	Grassland (May–Nov)	
○ <i>Hypochaeris glabra</i>	Smooth Cat's-ear	Herbaceous	- - -	L	ZA	Common. Disturbed areas, grassland, open woodland (Mar–Jun)	
○ <i>Hypochaeris radicata</i>	Rough Cat's-ear	Herbaceous	- - -	M	ZP	Disturbed areas, grassland, open woodland (Apr–Jul)	
○ <i>Iva axillaris</i> (JM93: subsp. <i>robustior</i>)	Poverty Weed	Herbaceous	- - -	-	NP	Seasonally wet, saline habitats, roadsides (Apr–Oct)	
○ <i>Jaumea carnosa</i>	Fleshy Jaumea	Herbaceous	- - C	-	NP	Coastal salt marshes, bases of sea cliffs (Apr–Dec)	
○ <i>Lactuca saligna</i>	Willow Lettuce	Herbaceous	- - -	-	ZA	Roadsides, grassland (Jul–Nov)	
○ <i>Lactuca serriola</i>	Prickly Lettuce	Herbaceous	- - -	-	ZA	Abundant. Disturbed places (May–Oct)	
○ <i>Lactuca</i> sp.	Wild Lettuce	Herbaceous	- - -	-	ZA	--	
○ <i>Lactuca virosa</i>	Tall Wild Lettuce	Herbaceous	- - -	-	ZB	Disturbed, shrubby and wooded slopes (Jun–Aug)	

Asteraceae		Sorted Alphabetically by Family, Scientific Name				Asteraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Laennecia coulteri</i> (JM93: <i>Conyza</i>)	Coulter's Horseweed	Herbaceous	- - A1	-	NA	Disturbed sites, clayey or sandy soils, often seasonally wet, alkaline (May–Nov)
<input type="radio"/> <i>Lagophylla ramosissima</i> (JM93: <i>subsp. ramosissima</i>)	Common Hare-leaf	Herbaceous	- - -	-	NA	Grassland, openings in scrub, woodland, forest (Apr–Oct)
<input type="radio"/> <i>Lasthenia californica subsp. californica</i> (JM93: <i>no subsp.</i>)	California Goldfields	Herbaceous	- - -	-	NA	Many habitats (Feb–Jun)
<input type="radio"/> <i>Lasthenia glabrata subsp. glabrata</i>	Yellow-ray Goldfields	Herbaceous	- - B	-	NA	Saline places, vernal pools (Mar–May)
<input type="radio"/> <i>Lasthenia microglossa</i>	Small-ray Goldfields	Herbaceous	- - B	-	NA	Shaded slopes of woodland, chaparral, desert scrub (Mar–May)
<input type="radio"/> <i>Layia chrysanthemoides</i>	Chrysanthemum Tidytops	Herbaceous	- - A2	-	NA	Grassy or open heavy soil, sometimes ± alkaline (Mar–Jun)
<input type="radio"/> <i>Layia gaillardiioides</i>	Yellow Woodland Tidytops	Herbaceous	- - A2	-	NA	Open or semi-shaded slopes, in sandy or clayey soil (including serpentine) (Mar–Aug)
<input type="radio"/> <i>Layia hieracioides</i>	Tall Tidytops	Herbaceous	- - A2	-	NA	Open, semi-shady, or disturbed sites, in light soil (Apr–Jul)
<input type="radio"/> <i>Layia platyglossa</i>	Tidy-tips	Herbaceous	- - C	-	NA	Many habitats (Feb–Jul)
<input type="radio"/> <i>Lessingia nemaclada</i>	Funnel-flower Lessingia	Herbaceous	- - A1	-	NA	Open fields, roadsides, woodland borders, occasionally on serpentine soil (Jul–Oct)
<input type="radio"/> <i>Lessingia sp.</i>	Lessingia	Herbaceous	- - -	-	NA	--
<input type="radio"/> <i>Logfia filaginoides</i> (JM93: <i>Filago californica</i>)	California Fluffweed	Herbaceous	- - -	-	NA	Common, ± weedy. Bare, rocky, or grassy sites, drainages (Feb–May)
<input type="radio"/> <i>Logfia gallica</i> (JM93: <i>Filago</i>)	Daggerleaf Cottonrose	Herbaceous	- - -	-	ZA	Bare or grassy openings, burns (Mar–Jul)
<input type="radio"/> <i>Madia anomala</i>	Plump-seed Tarweed	Herbaceous	- - A1	-	NA	Openings in woodland or chaparral (Apr–Jun)
<input type="radio"/> <i>Madia exigua</i>	Thread-stem Tarweed	Herbaceous	- - -	-	NA	Grassy, open, or disturbed sites, in sandy or clayey soils, including serpentine (Apr–Jul)
<input type="radio"/> <i>Madia gracilis</i>	Gumweed	Herbaceous	- - -	-	NA	Open, semi-shaded or disturbed sites, many habitats, incl serpentine (Apr–Aug)
<input type="radio"/> <i>Madia or Anisocarpus sp.</i> (JM93: <i>Madia sp.</i>)	Tarweed	Herbaceous	- - -	-	NA	--
<input type="radio"/> <i>Madia sativa</i>	Coast Tarweed	Herbaceous	- - -	-	NA	Grassy, open, or disturbed sites (May–Oct)
<input type="radio"/> <i>Malacothrix coulteri</i>	Snake's Head Dandelion	Herbaceous	- - A1	-	NA	Sandy, open areas, in coastal-sage scrub, grassland, desert (Mar–May)
<input type="radio"/> <i>Malacothrix floccifera</i>	Woolly / Flocked Dandelion	Herbaceous	- - A2	-	NA	In loose soil of open burns, slides, in chaparral, pinyon/juniper woodland, yellow-pine forest (Mar–Nov)
<input type="radio"/> <i>Matricaria discoidea</i> (JM93: <i>Chamomilla suaveolens</i>)	Pineapple Weed	Herbaceous	- - -	-	ZA	Abundant. Disturbed sites, riverbanks (Feb–Aug)
<input type="radio"/> <i>Micropus amphibolus</i>	Mount Diablo Cottonweed	Herbaceous	- 3.2 *A1	-	NA	Openings on slopes, ridges, shallow soils, sedimentary or volcanic rocks (Mar–Jun)
<input type="radio"/> <i>Micropus californicus var. ?</i>	Slender Cottonweed	Herbaceous	- - -	-	NA	--

Asteraceae		Sorted Alphabetically by Family, Scientific Name					Asteraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Micropus californicus</i> var. <i>californicus</i>	Cottonweed	Herbaceous	- - -	-	NA	Clearings, often disturbed, dry or seasonally moist soils (Mar–Jun)	
<input type="radio"/> <i>Microseris acuminata</i>	Sierra Foothill Silverpuffs	Herbaceous	- - B	-	NA	Grassy, open rocky or clay soil (Apr–Jun)	
<input type="radio"/> <i>Microseris douglasii</i> subsp. ?	Douglas Silverpuffs	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Microseris douglasii</i> subsp. <i>douglasii</i>	Douglas Silverpuffs	Herbaceous	- - -	-	NA	Grassland, near vernal pools or serpentine (Mar–Jun)	
<input type="radio"/> <i>Microseris elegans</i>	Nodding Silverpuffs	Herbaceous	- - A2	-	NA	Generally inland clay grassland, often near vernal pools (Apr–Jun)	
<input type="radio"/> <i>Microseris</i> sp.	Silverpuffs	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Monolopia major</i>	Cupped Monolopia	Herbaceous	- - B	-	NA	Grassland, bare clay (Feb–Jul)	
<input type="radio"/> <i>Packera breweri</i> (JM93: <i>Senecio</i>)	Brewer's Ragwort	Herbaceous	- - B	-	NP	Common. Seasonally damp grassland, oak savanna, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Pentachaeta exilis</i> subsp. <i>exilis</i>	Meager Pentachaeta	Herbaceous	- - A1	-	NA	Grassland, foothill woodland (Mar–May)	
<input type="radio"/> <i>Pluchea odorata</i> var. <i>odorata</i> (JM93: no var.)	Saltmarsh-fleabane	Herbaceous	- - B	-	NP	Moist, often saline valley bottoms (Jun–Nov)	
<input type="radio"/> <i>Pseudognaphalium beneolens</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>beneolens</i>)	Scented White Everlasting	Herbaceous	- - C	-	NB	Dry, open slopes, disturbed sites (Jun–Oct)	
<input type="radio"/> <i>Pseudognaphalium biolettii</i> (JM93: <i>Gnaphalium bicolor</i>)	Two-color Everlasting	Herbaceous	- - A2	-	NP	Rocky slopes, roadsides, dunes, coastal scrub, chaparral, oak woodland (Apr–Jun)	
<input type="radio"/> <i>Pseudognaphalium californicum</i> (JM93: <i>Gnaphalium</i>)	California Everlasting	Herbaceous	- - -	-	NB	Sandy canyons, dry hills, coastal chaparral (Apr–Jul)	
<input type="radio"/> <i>Pseudognaphalium luteoalbum</i> (JM93: <i>Gnaphalium luteo-album</i>)	Weedy Cudweed	Herbaceous	- - -	-	ZA	Disturbed sites, fields, streambeds (Apr–Aug)	
<input type="radio"/> <i>Pseudognaphalium microcephalum</i> (JM93: <i>Gnaphalium canescens</i> subsp. <i>microcephalum</i>)	White Everlasting	Herbaceous	- - B	-	NB	Grassy hillsides, gravelly canyon bottoms, chaparral, coastal-sage scrub (Jun–Aug)	
<input type="radio"/> <i>Pseudognaphalium ramosissimum</i> (JM93: <i>Gnaphalium</i>)	Pink Everlasting	Herbaceous	- - -	-	NB	Dry, open slopes, woodland, sandy fields, dunes (Jul–Sep)	
<input type="radio"/> <i>Pseudognaphalium stramineum</i> (JM93: <i>Gnaphalium</i>)	Cotton-batting Plant	Herbaceous	- - -	-	NA	Many habitats, dunes, chaparral slopes, roadsides (Mar–Aug)	
<input type="radio"/> <i>Psilocarphus brevissimus</i> var. <i>brevissimus</i>	Dwarf Woollyheads	Herbaceous	- - B	-	NA	Drying edges of vernal pools, mud flats, drainages (May–Jun)	
<input type="radio"/> <i>Psilocarphus oregonus</i>	Oregon Woollyheads	Herbaceous	- - B	-	NA	Drying clay of vernal pools, drainages, moist rocky slopes (Mar–Jul)	
<input type="radio"/> <i>Psilocarphus tenellus</i> (JM93: var. <i>tenellus</i>)	Slender Woolly-marbles	Herbaceous	- - -	-	NA	Common. Dry, seasonally moist slopes, flats, burns, trails, rarely vernal pools (Mar–Jul)	
<input type="radio"/> <i>Rafinesquia californica</i>	California Chicory	Herbaceous	- - B	-	NA	Open sites in scrub, woodland; often common after fire (Apr–Jul)	
<input type="radio"/> <i>Rigiopappus leptocladus</i>	Bristle-head	Herbaceous	- - B	-	NA	Grassy sites, openings among shrubs or trees (Apr–Jul)	

Asteraceae		Sorted Alphabetically by Family, Scientific Name				Asteraceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Senecio aronicoides</i>	California Woolly Butterweed	Herbaceous	- - C	-	NP	Dry or drying sites in open woodland, upper foothill, montane forest (Apr–Jul)	
<input type="radio"/> <i>Senecio flaccidus</i> var. <i>douglasii</i>	Douglas' Threadleaf Ragwort	Woody	- - A2	-	NP	Generally disturbed dry, open, sandy or rocky sites (Jun–Oct)	
<input type="radio"/> <i>Senecio glomeratus</i> (JM93: <i>Erechtites glomerata</i>)	Cutleaf Burnweed	Herbaceous	- - -	M	ZA	Disturbed sites, mostly coastal (Apr–Sep)	
<input type="radio"/> <i>Senecio hydrophilus</i>	Water Ragwort	Herbaceous	- - A1	-	NP	Marshes, swampy places, standing water, alkaline sites (May–Sep)	
<input type="radio"/> <i>Senecio minimus</i> (JM93: <i>Erechtites minima</i>)	Coastal Burnweed	Herbaceous	- - -	M	ZA	Disturbed coastal sites (Jun–Sep)	
<input type="radio"/> <i>Senecio sylvaticus</i>	Woodland Ragwort	Herbaceous	- - -	-	ZA	Disturbed woodland, open, sandy sites (Mar–Sep)	
<input type="radio"/> <i>Senecio vulgaris</i>	Common Groundsel	Herbaceous	- - -	-	ZA	Common. Disturbed areas (Feb–Jul)	
<input type="radio"/> <i>Silybum marianum</i>	Milk Thistle	Herbaceous	- - -	L	ZAB	Roadsides, pastures, disturbed areas (Feb–Jun)	
<input type="radio"/> <i>Solidago elongata</i> (JM93: <i>S. canadensis</i> subsp. <i>elongata</i>)	West Coast Canada Goldenrod	Herbaceous	- - -	-	NP	Meadows, thickets (May–Oct)	
<input type="radio"/> <i>Solidago</i> sp.	Goldenrod	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Solidago velutina</i> subsp. <i>californica</i> (JM93: <i>S. californica</i>)	California Goldenrod	Herbaceous	- - -	-	NP	Woodland margins, grassland, disturbed soils (May–Nov)	
<input type="radio"/> <i>Soliva sessilis</i>	Common Soliva	Herbaceous	- - -	-	ZA	Disturbed areas, especially hard-packed paths (Apr–Jul)	
<input type="radio"/> <i>Sonchus asper</i> subsp. <i>asper</i>	Prickly Sow Thistle	Herbaceous	- - -	-	ZA	Common. Slightly moist disturbed sites, along streams (All year)	
<input type="radio"/> <i>Sonchus oleraceus</i>	Common Sow Thistle	Herbaceous	- - -	-	ZA	Abundant. Disturbed places (All year)	
<input type="radio"/> <i>Stebbinsoseris heterocarpa</i>	Grassland Silverpuffs	Herbaceous	- - B	-	NA	Open, sometimes disturbed sites, rocky to clay soils, generally inland (Apr–Jun)	
<input type="radio"/> <i>Stephanomeria elata</i>	Tall Wreath Plant	Herbaceous	- - A2	-	NA	Chaparral openings, grassy meadows, roadside embankments (Jul–Nov)	
<input type="radio"/> <i>Stephanomeria virgata</i> subsp. <i>pleurocarpa</i>	Twiggy Wreath Plant	Herbaceous	- - C	-	NA	Chaparral openings, grassland (Jun–Nov)	
<input type="radio"/> <i>Symphyotrichum chilense</i> (JM93: <i>Aster chilensis</i>)	Common California Wild Aster	Herbaceous	- - -	-	NP	Grassland, salt marshes, disturbed places (Jun–Oct)	
<input type="radio"/> <i>Symphyotrichum lentum</i> (JM93: <i>Aster lentus</i>)	Suisun Marsh Aster	Herbaceous	- 1B.2 *A2	-	NP	Marshes (May–Nov)	
<input type="radio"/> <i>Symphyotrichum subulatum</i> var. <i>parviflorum</i> (JM93: <i>Aster subulatus</i> var. <i>ligulatus</i>)	Annual Saltmarsh Aster	Herbaceous	- - -	-	NA	Marshes, disturbed places (Jul–Oct)	
<input type="radio"/> <i>Taraxacum officinale</i>	Common Dandelion	Herbaceous	- - -	-	ZBP	Abundant. Esp disturbed areas (All year)	
<input type="radio"/> <i>Tragopogon dubius</i>	Yellow Salsify	Herbaceous	- - -	-	ZAB	Uncommon. Disturbed places (May–Sep)	
<input type="radio"/> <i>Tragopogon porrifolius</i>	Purple Salsify	Herbaceous	- - -	-	ZBP	Common. Disturbed places (Mar–Nov)	
<input type="radio"/> <i>Uropappus lindleyi</i>	Silverpuffs	Herbaceous	- - -	-	NA	Common. Open grassland, woodland, chaparral, deserts, gen in loose soils (Mar–May)	
<input type="radio"/> <i>Urospermum picroides</i>	False Hawkbit	Herbaceous	- - -	-	ZAP	Uncommon. Disturbed places (Apr–Jul)	

Asteraceae		Sorted Alphabetically by Family, Scientific Name					Asteraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Wyethia angustifolia</i>	Narrow-leaved Mule's Ears	Herbaceous	- - -	-	NP	Grassland (Apr–Aug)	
<input type="radio"/> <i>Wyethia glabra</i>	Smooth Mule's Ears	Herbaceous	- - C	-	NP	Gen shady sites (Mar–Jun)	
<input type="radio"/> <i>Wyethia helenioides</i>	Gray Mule's Ears	Herbaceous	- - -	-	NP	Open grassland, woodland, scrub (Mar–May(Aug))	
<input type="radio"/> <i>Xanthium spinosum</i>	Spiny Cocklebur	Herbaceous	- - -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
<input type="radio"/> <i>Xanthium strumarium</i>	Cocklebur	Herbaceous	- - -	-	NA	Disturbed, seasonally wet, often alkaline sites, in grassland, marshes, watercourses (Jul–Oct)	
Azollaceae							Azollaceae
<input type="radio"/> <i>Azolla filiculoides</i>	Mosquito Fern	Fern-like	- - -	-	NP	Common. Ponds, slow streams -	
<input type="radio"/> <i>Azolla microphylla</i> (JM93: <i>A. mexicana</i>)	Mexican Mosquito Fern	Fern-like	- - -	-	NP	Ponds, slow streams -	
Berberidaceae							Berberidaceae
<input type="radio"/> <i>Berberis aquifolium</i> var. <i>dictyota</i>	Hollyleaf Oregon Grape	Woody	- - A2	-	NP	Slopes, canyons, conifer forest, oak woodland, chaparral (Mar–May)	
<input type="radio"/> <i>Berberis nervosa</i>	Longleaf Oregon Grape	Woody	- - A1	-	NP	Conifer forest (Mar–Jun)	
<input type="radio"/> <i>Berberis pinnata</i> subsp. <i>pinnata</i>	Shinyleaf Oregon-grape	Woody	- - C	-	NP	Rocky slopes, conifer forest, oak woodland, chaparral (Feb–May)	
<input type="radio"/> <i>Berberis</i> sp.	Oregon Grape	Woody	- - -	-	NP	--	
Betulaceae							Betulaceae
<input type="radio"/> <i>Alnus rhombifolia</i>	White Alder	Woody	- - -	-	NP	Along permanent streams (Apr–Jun)	
<input type="radio"/> <i>Alnus rubra</i>	Red Alder	Woody	- - A2	-	NP	Wet places, especially after logging (Feb–Mar)	
<input type="radio"/> <i>Alnus</i> sp.	Alder	Woody	- - -	-	NP	--	
<input type="radio"/> <i>Corylus cornuta</i> subsp. <i>californica</i> (JM93: var. <i>californica</i>)	California Hazelnut	Woody	- - -	-	NP	Common. Many habitats, esp moist, shady places (Jan–Mar)	
Blechnaceae							Blechnaceae
<input type="radio"/> <i>Woodwardia fimbriata</i>	Giant Chain Fern	Fern-like	- - -	-	NP	Near streams, springs, seeps -	
Boraginaceae							Boraginaceae
<input type="radio"/> <i>Amsinckia grandiflora</i>	Large-flowered Fiddleneck	Herbaceous	FE/CE 1B.1 *A1	-	NA	Grassy slopes (Mar–May)	
<input type="radio"/> <i>Amsinckia intermedia</i> (JM93: <i>A. menziesii</i> var. <i>intermedia</i>)	Common Fiddleneck	Herbaceous	- - -	-	NA	Abundant. Open, generally disturbed places (Mar–Jun)	
<input type="radio"/> <i>Amsinckia lunaris</i>	Bent-flowered Fiddleneck	Herbaceous	- 1B.2 *A2	-	NA	Gravelly slopes, grassland, openings in woodland (Mar–Jun)	
<input type="radio"/> <i>Amsinckia lycopsoides</i>	Bugloss-flowered Fiddleneck	Herbaceous	- - B	-	NA	Common. Open, grassland, foothill woodland, generally disturbed areas (Mar–Jun)	

Boraginaceae	Sorted Alphabetically by Family, Scientific Name					Boraginaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Amsinckia menziesii</i> (JM93: var. <i>menziesii</i>)	Small-flowered Fiddleneck	Herbaceous	- - -	-	NA	Shade-tolerant, open, disturbed areas at forest/woodland edges (May–Jul)
<input type="radio"/> <i>Amsinckia</i> sp.	Fiddleneck	Herbaceous	- - -	-	NA	--
<input type="radio"/> <i>Amsinckia tessellata</i> var. ?	Devil's Lettuce	Herbaceous	- - -	-	NA	--
<input type="radio"/> <i>Amsinckia tessellata</i> var. <i>tessellata</i>	Desert Fiddleneck	Herbaceous	- - B	-	NA	Rocky or sandy soils (Feb–Jun)
<input type="radio"/> <i>Cryptantha flaccida</i>	Weak-stemmed Cryptantha	Herbaceous	- - -	-	NA	Common; semi-barren, gravelly, loose soils, rocky sites, washes, slopes, ridges (Apr–Jul)
<input type="radio"/> <i>Cryptantha intermedia</i> var. <i>intermedia</i> (JM93: no var.)	Intermediate Cryptantha	Herbaceous	- - A1	-	NA	Sandy to gravelly flats, slopes, generally granitic or serpentine-based, grassland, chaparral, foothill woodland, occasionally open conifer forest (May–Jul)
<input type="radio"/> <i>Cryptantha micromeres</i>	Minute-flowered Cryptantha	Herbaceous	- - A1	-	NA	Open sites, disturbed, coarse soils, chaparral, woodland, burns (Mar–Jul)
<input type="radio"/> <i>Cryptantha muricata</i> var. <i>muricata</i> (JM93: no var.)	Prickly Cryptantha	Herbaceous	- - A2	-	NA	Coastal scrub, chaparral, foothill woodland, open conifer forest (Mar–Jul)
<input type="radio"/> <i>Cryptantha rattanii</i> (JM93: <i>C. decipiens</i>)	Deceptive Cryptantha	Herbaceous	- - A2	-	NA	Rocky, gravelly slopes, grassland, coastal scrub, chaparral, foothill woodland (Apr–Jul)
<input type="radio"/> <i>Cryptantha</i> sp.	Cryptantha	Herbaceous	- - -	-	NA	--
<input type="radio"/> <i>Cryptantha torreyana</i> var. <i>pumila</i> (JM93: no var.)	Dwarf Cryptantha	Herbaceous	- - A2	-	NA	Generally chaparral, foothill woodland (Apr–Jun)
<input type="radio"/> <i>Cynoglossum grande</i>	Grand Hound's Tongue	Herbaceous	- - -	-	NP	Chaparral, woodland (Feb–May)
<input type="radio"/> <i>Emmenanthe penduliflora</i> var. <i>penduliflora</i>	Whispering Bells	Herbaceous	- - B	-	NP	Chaparral to creosote-bush scrub, rocky, sandy, decomposed granite, serpentine soils (Apr–Jul)
<input type="radio"/> <i>Eriodictyon californicum</i>	California Yerba Santa	Woody	- - -	-	NP	Slopes, fields, roadsides, woodland, chaparral (Apr–Jul)
<input type="radio"/> <i>Eucrypta chrysanthemifolia</i> var. <i>chrysanthemifolia</i>	Common Eucrypta	Herbaceous	- - B	-	NA	Roadsides, burns, coastal bluffs, ravines (Mar–Jun)
<input type="radio"/> <i>Heliotropium curassavicum</i> var. <i>oculatum</i> (JM93: no var.)	Seaside Heliotrope	Herbaceous	- - -	-	NP	Moist to dry, saline to alkaline soils, gen near water (Feb–Oct)
<input type="radio"/> <i>Hydrophyllum occidentale</i>	Western Waterleaf	Herbaceous	- - A2	-	NP	Moist, shaded slopes, woodland, meadows, streambanks, chaparral (May–Jul)
<input type="radio"/> <i>Myosotis latifolia</i>	Broadleaved Forget-me-not	Herbaceous	- - -	L	ZP	Moist, disturbed, shady places (Feb–Jul)
<input type="radio"/> <i>Nemophila heterophylla</i>	Variable-leaf Nemophila	Herbaceous	- - -	-	NA	Common. Forest, chaparral, roadsides, streambanks (Feb–Jun)
<input type="radio"/> <i>Nemophila menziesii</i> var. <i>menziesii</i>	Baby Blue-eyes	Herbaceous	- - -	-	NA	Meadows, grassland, chaparral, woodland, slopes (Feb–May)
<input type="radio"/> <i>Nemophila parviflora</i> var. <i>parviflora</i>	Small-flower Nemophila	Herbaceous	- - C	-	NA	Woodland, forest, roadsides, slopes (Mar–Jul)

Boraginaceae	Sorted Alphabetically by Family, Scientific Name					Boraginaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Nemophila pedunculata</i>	Meadow Nemophila	Herbaceous	- - B	-	NA	Common. Ocean bluffs, grassland, slopes, meadows, sandbars, fields, woodland, streambanks (Feb–Jul)
<input type="radio"/> <i>Pectocarya penicillata</i>	Northern Pectocarya	Herbaceous	- - A1	-	NA	Disturbed sites, roadsides (Feb–May)
<input type="radio"/> <i>Pectocarya pusilla</i>	Little Pectocarya	Herbaceous	- - B	-	NA	Dry, semi-barren sites in grassland, chaparral, woodland, roadsides (Mar–Jun)
<input type="radio"/> <i>Phacelia breweri</i>	Brewer Phacelia	Herbaceous	- - A2	-	NP	Rocky soils, slopes, chaparral, oak woodland (Mar–Jun)
<input type="radio"/> <i>Phacelia californica</i>	California Phacelia	Herbaceous	- - -	-	NP	Bluffs, open slopes, road cuts, chaparral, woodland (Mar–Sep)
<input type="radio"/> <i>Phacelia ciliata</i>	Field Phacelia	Herbaceous	- - B	-	NA	Clay or gravelly slopes in grassland, fields (Feb–Jun)
<input type="radio"/> <i>Phacelia distans</i>	Common Phacelia	Herbaceous	- - -	-	NA	Common. Clay to rocky soils, slopes (Mar–May)
<input type="radio"/> <i>Phacelia divaricata</i>	Divaricate Phacelia	Herbaceous	- - A2	-	NA	Open areas, chaparral, woodland, grassland (Apr–Jun)
<input type="radio"/> <i>Phacelia heterophylla subsp. virgata</i>	Varileaf Phacelia	Herbaceous	- - -	-	NBP	Slopes, flats, roadsides (May–Jul)
<input type="radio"/> <i>Phacelia imbricata subsp. imbricata</i>	Rock Phacelia	Herbaceous	- - -	-	NP	Slopes, roadsides, flats, canyons, chaparral, woodland (Apr–Jul)
<input type="radio"/> <i>Phacelia malvifolia</i>	Stinging Phacelia	Herbaceous	- - A2	-	NA	Sandy or gravelly soils, scrub, conifer forest (Apr–Jul)
<input type="radio"/> <i>Phacelia nemoralis subsp. nemoralis</i>	Bristly Phacelia	Herbaceous	- - B	-	NBP	Moist slopes, streambanks, mixed-evergreen forest (Apr–Jul)
<input type="radio"/> <i>Phacelia ramosissima (JM93: var. ramosissima, latifolia)</i>	Branched Phacelia	Herbaceous	- - A2	-	NP	Diverse habitats incl salt marshes, canyons (Apr–Oct)
<input type="radio"/> <i>Phacelia rattanii</i>	Rattan Phacelia	Herbaceous	- - A1	-	NA	Shaded crevices, steep slopes (May–Jul)
<input type="radio"/> <i>Phacelia tanacetifolia</i>	Tansy-leaf Phacelia	Herbaceous	- - B	-	NA	Sandy to gravelly slopes, open areas (Mar–May)
<input type="radio"/> <i>Pholistoma auritum var. auritum</i>	Fiesta Flower	Herbaceous	- - B	-	NA	Ocean bluffs, talus slopes, woodland, streambanks, canyons (Mar–Jun)
<input type="radio"/> <i>Pholistoma membranaceum</i>	White Fiesta Flower	Herbaceous	- - -	-	NA	Beaches, bluffs, ravines, wooded slopes, desert washes (Feb–May)
<input type="radio"/> <i>Plagiobothrys acanthocarpus</i>	Adobe Popcornflower	Herbaceous	- - B	-	NA	Vernal pools, moist clay soil (Mar–May)
<input type="radio"/> <i>Plagiobothrys bracteatus</i>	Bracted Popcornflower	Herbaceous	- - -	-	NA	Common. Vernal pools, wet places in grassland, coastal-sage scrub, chaparral (Apr–Jun)
<input type="radio"/> <i>Plagiobothrys canescens var. canescens (JM93: no var.)</i>	Valley Popcornflower	Herbaceous	- - C	-	NA	Common. Grassland, woodland, coastal scrub, desert scrub, roadsides (Mar–Jun)
<input type="radio"/> <i>Plagiobothrys fulvus var. campestris (JM93: no var.)</i>	Field Popcornflower	Herbaceous	- - B	-	NA	Grassland, open woodland, loamy, sandy, or gravelly soil (Mar–May)
<input type="radio"/> <i>Plagiobothrys humistratus</i>	Low Popcornflower	Herbaceous	- - B	-	NA	Vernal pools, wet places, grassland (Mar–May)

Boraginaceae		Sorted Alphabetically by Family, Scientific Name					Boraginaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Plagiobothrys infectivus</i>	Dye Popcornflower	Herbaceous	- - A1	-	NA	Open grassland, friable clay to sandy soils (Mar–May)	
<input type="radio"/> <i>Plagiobothrys nothofulvus</i>	Rusty Popcornflower	Herbaceous	- - -	-	NA	Common; open woodland, grassland (Mar–May)	
<input type="radio"/> <i>Plagiobothrys sp.</i>	Popcornflower	Herbaceous	- - -	-	-	-	
<input type="radio"/> <i>Plagiobothrys stipitatus var. micranthus</i>	Valley Popcornflower	Herbaceous	- - -	-	NA	Common. Vernal pools, wet sites in grassland, conifer forest, sagebrush scrub (Apr–Jul)	
<input type="radio"/> <i>Plagiobothrys stipitatus var. stipitatus</i>	Showy Great Popcornflower	Herbaceous	- - C	-	NA	Vernal pools, wet sites in grassland (Mar–Jun)	
<input type="radio"/> <i>Plagiobothrys trachycarpus</i>	Rough-nutlet Popcornflower	Herbaceous	- - B	-	NA	Shallow vernal pools, wet places in grassland, scrub, chaparral, woodland (Mar–May)	
Brassicaceae							Brassicaceae
<input type="radio"/> <i>Athysanus pusillus</i>	Dwarf Sandweed	Herbaceous	- - -	-	NA	Grassy, open slopes, rocky outcrops, chaparral, flats, floodplains, cliffs, ledges (Feb–Jun)	
<input type="radio"/> <i>Barbarea orthoceras</i>	Erect-pod Winter Cress	Herbaceous	- - -	-	NP	Meadows, streambanks, moist woodland, grassland (Mar–Jul)	
<input type="radio"/> <i>Boechera breweri subsp. breweri (JM93: Arabis breweri var. breweri)</i>	Brewer's Rockcress	Herbaceous	- - A1	-	NP	Rocky outcrops, ledges, talus Mar–Jul	
<input type="radio"/> <i>Brassica nigra</i>	Black Mustard	Herbaceous	- - -	M	ZA	Common. Disturbed areas, fields (Apr–Sep)	
<input type="radio"/> <i>Brassica rapa</i>	Turnip	Herbaceous	- - -	L	ZA	Disturbed areas (Jan–May)	
<input type="radio"/> <i>Cakile maritima</i>	Horned Sea Rocket	Herbaceous	- - -	L	ZA	Beach dunes (May–Nov)	
<input type="radio"/> <i>Capsella bursa-pastoris</i>	Shepherd's Purse	Herbaceous	- - -	-	ZA	Disturbed areas (Jan–Oct)	
<input type="radio"/> <i>Cardamine californica</i>	Milk Maids	Herbaceous	- - -	-	NP	Gen shaded sites, canyons, woodland. One of first spring flowers (Jan–May)	
<input type="radio"/> <i>Cardamine oligosperma</i>	Western Bitter-cress	Herbaceous	- - -	-	NA	Wet meadows, shady banks, damp areas (Mar–Jul)	
<input type="radio"/> <i>Caulanthus flavescens (JM93: Guillenia)</i>	Yellow-flowered Jewelflower	Herbaceous	- - A2	-	NA	Dry, exposed slopes, generally on serpentine, open hillsides (Mar–May)	
<input type="radio"/> <i>Caulanthus lasiophyllus (JM93: Guillenia lasiophylla)</i>	California Mustard	Herbaceous	- - -	-	NA	Common. Desert flats, sandy banks, gravelly or rocky areas, talus slopes, shrubland, grassy fields, disturbed site Mar–Jun	
<input type="radio"/> <i>Descurainia sophia</i>	Feather-leaf Tansy Mustard	Herbaceous	- - -	L	ZA	Common. Disturbed areas, fields, canyon bottoms, desert (May–Aug)	
<input type="radio"/> <i>Draba cuneifolia (JM93: var. integrifolia)</i>	Wedge-leaf Whitlow Grass	Herbaceous	- - A1	-	NA	Open or disturbed places (May)	
<input type="radio"/> <i>Draba verna</i>	Spring Whitlow Grass	Herbaceous	- - B	-	NA	Open or disturbed areas (Feb–May)	
<input type="radio"/> <i>Erysimum capitatum var. capitatum (JM93: var. capitatum, var. angustatum)</i>	Western Wallflower	Herbaceous	- - C	-	NB	Common. Open areas, woodland, sandy areas, chaparral (Mar–Sep)	
<input type="radio"/> <i>Hirschfeldia incana</i>	Shortpod Mustard	Herbaceous	- - -	M	ZBP	Disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Idahoa scapigera</i>	Flat-pod	Herbaceous	- - A1	-	NA	Moist ledges, slopes, meadows, foothills (Mar–May)	

Brassicaceae		Sorted Alphabetically by Family, Scientific Name					Brassicaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
○ <i>Lepidium dictyotum</i> (JM93: var. <i>dictyotum</i>)	Alkali Peppergrass	Herbaceous	- - B	-	NA	Saline soils, dry stream beds, roadsides, sandy flats, fields, meadows, dried pools (Mar–Jun)	
○ <i>Lepidium didymum</i> (JM93: <i>Coronopus didymus</i>)	Lesser Swine Cress	Herbaceous	- - -	-	ZA	Common. Disturbed areas, fields, pastures (Mar–Jul)	
○ <i>Lepidium draba</i> (JM93: <i>Cardaria</i>)	Heart-podded Hoary Cress	Herbaceous	- - -	M	ZP	Disturbed areas, saline soils, pastures, fields (Apr–Aug)	
○ <i>Lepidium latifolium</i>	Perennial Peppergrass	Herbaceous	- - -	H	ZP	Pastures, disturbed areas, fields, grassland, saline meadows, streambanks, sagebrush scrub, edge of marshes (Jun–Sep)	
○ <i>Lepidium latipes</i> (JM93: var. <i>latipes</i>)	Dwarf Peppergrass	Herbaceous	- - B	-	NA	Alkaline soils, vernal pool margins, salt marsh edges, pastures (Mar–Jun)	
○ <i>Lepidium nitidum</i> (JM93: var. <i>nitidum</i> , <i>oreganum</i>)	Threadleaf Peppergrass	Herbaceous	- - -	-	NA	Alkaline soils, pastures, dry vernal pools, fields, beaches (Feb–Mar)	
○ <i>Lepidium oxycarpum</i>	Sharp-pod Peppergrass	Herbaceous	- - B	-	NA	Fields, vernal pool margins, alkaline flats (Mar–May)	
○ <i>Lepidium strictum</i>	Prostrate Peppergrass	Herbaceous	- - -	-	ZA	Uncommon. Disturbed areas, woodland, slopes (Apr–Jun)	
○ <i>Lobularia maritima</i>	Sweet Alyssum	Herbaceous	- - -	L	ZP	Disturbed areas, fields (Mar–Oct)	
○ <i>Lunaria annua</i>	Money Plant	Herbaceous	- - -	-	ZA	Disturbed areas (Apr–Jun)	
○ <i>Nasturtium officinale</i> (JM93: <i>Rorippa nasturtium-aquaticum</i>)	Water Cress	Herbaceous	- - -	-	NP	Streams, springs, marshes, lake margins, swamps (Mar–Nov)	
○ <i>Raphanus raphanistrum</i>	Jointed Charlock	Herbaceous	- - -	-	ZA	Disturbed areas, fields (Apr–Jul)	
○ <i>Raphanus sativus</i>	Radish	Herbaceous	- - -	L	ZA	Disturbed areas, fields (May–Jul)	
○ <i>Rorippa curvisiliqua</i>	Yellow Water Cress	Herbaceous	- - A2	-	NA	Uncommon. Streambanks, marshy ground, seepage areas, mud flats, meadows (May–Oct)	
○ <i>Rorippa palustris</i> subsp. <i>palustris</i> (JM93: var. <i>occidentalis</i>)	Marsh Water Cress	Herbaceous	- - A2	-	NA	Generally wet places (Mar–Sep)	
○ <i>Sinapis arvensis</i>	Charlock	Herbaceous	- - -	L	ZA	Disturbed areas (Mar–Sep)	
○ <i>Sisymbrium irio</i>	London Rocket	Herbaceous	- - -	M	ZA	Disturbed areas, fields, pastures (Jan–Apr)	
○ <i>Sisymbrium officinale</i>	Hedge Mustard	Herbaceous	- - -	-	ZA	Disturbed areas, fields, pastures (Apr–Sep)	
○ <i>Sisymbrium</i> sp.	Rocket / Mustard	Herbaceous	- - -	-	ZA	--	
○ <i>Streptanthus glandulosus</i> subsp. <i>glandulosus</i> (JM93: incl. <i>S. albidus</i> subsp. <i>peramoenus</i>)	Bristly Jewel Flower	Herbaceous	- - B	-	NA	Serpentine, bare slopes, chaparral & woodland openings (Apr–Jul)	
○ <i>Thysanocarpus curvipes</i>	Hairy Fringepod	Herbaceous	- - -	-	NA	Common. Slopes, washes, moist meadows, woodland, streambanks (Feb–Jun)	
○ <i>Thysanocarpus laciniatus</i> var. <i>laciniatus</i> (JM93: no var.)	Narrowleaf Fringepod	Herbaceous	- - B	-	NA	Oak woodland, rocky ridges, slopes, chaparral, washes (Mar–May)	

Brassicaceae		Sorted Alphabetically by Family, Scientific Name				Brassicaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Thysanocarpus radicans</i>	Ribbed Fringepod	Herbaceous	- - A2	-	NA	Moist slopes, pastures, open meadows, fields (Mar–Apr)	
<input type="radio"/> <i>Tropidocarpum gracile</i>	Lacepod	Herbaceous	- - B	-	NP	Grassy banks, open fields, roadsides, pastures (Mar–May)	
<input type="radio"/> <i>Turritis glabra</i> (JM93: <i>Arabis</i> , var. <i>glabra</i>)	Tower Mustard	Herbaceous	- - C	-	NP	Open fields, meadows, slopes (Apr–Jul)	
Callitrichaceae						Callitrichaceae	
<input type="radio"/> <i>Callitriche marginata</i>	California Water-starwort	Herbaceous	- - -	-	NA	Becoming stranded (often in vernal pools) (Mar–Jun)	
Campanulaceae						Campanulaceae	
<input type="radio"/> <i>Campanula exigua</i>	Chaparral / Rock Harebell	Herbaceous	- 1B.2 *A2	-	NA	Talus slopes, generally serpentine soil (May–Jun)	
<input type="radio"/> <i>Downingia ornatissima</i> var. <i>eximia</i>	Horned Downingia	Herbaceous	- - A1x	-	NA	Vernal pools, roadside ditches (Apr–May)	
<input type="radio"/> <i>Downingia pulchella</i>	Flatface Downingia	Herbaceous	- - B	-	NA	Vernal pools, roadside ditches (Apr–Jun)	
<input type="radio"/> <i>Downingia</i> sp.	Downingia	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Githopsis diffusa</i> subsp. <i>robusta</i>	Southern Bluecup	Herbaceous	- - A2	-	NA	Shaded or disturbed areas, burns (Apr–Jun)	
<input type="radio"/> <i>Githopsis specularioides</i>	Looking-glass Bluecup	Herbaceous	- - B	-	NA	Chaparral, oak woodland (Apr–May)	
<input type="radio"/> <i>Legenere limosa</i>	Legenere	Herbaceous	- 1B.1 *A1	-	NA	Wet areas, vernal pools, ponds (May–Jun)	
<input type="radio"/> <i>Triodanis biflora</i>	Small Venus-looking-glass	Herbaceous	- - A2	-	NA	Disturbed areas (Apr–Jun)	
Caprifoliaceae						Caprifoliaceae	
<input type="radio"/> <i>Lonicera hispidula</i> (JM93: var. <i>vacillans</i>)	Hairy Vine Honeysuckle	Woody	- - -	-	NP	Canyons, streamsides, woodland (May–Jun)	
<input type="radio"/> <i>Lonicera interrupta</i>	Chapparal Vine Honeysuckle	Woody	- - -	-	NP	Dry slopes, ridges, floodplains, oak woodland, chaparral (Apr–May)	
<input type="radio"/> <i>Lonicera involucrata</i> var. <i>ledebourii</i>	Black Twinberry Honeysuckle	Woody	- - B	-	NP	Moist places (May–Jul)	
<input type="radio"/> <i>Lonicera japonica</i>	Japanese Honeysuckle	Woody	- - -	-	ZP	Disturbed places (May–Jul)	
<input type="radio"/> <i>Lonicera subspicata</i> var. <i>denudata</i>	Southern Honeysuckle	Woody	- - B	-	NP	Chaparral slopes (Jun–Jul)	
<input type="radio"/> <i>Symphoricarpos albus</i> var. <i>laevigatus</i>	Snowberry	Woody	- - -	-	NP	Shady woodland, streambanks, N. slopes (May–Jul)	
<input type="radio"/> <i>Symphoricarpos mollis</i>	Creeping Snowberry	Woody	- - -	-	NP	Ridges, slopes, open places in woodland (Apr–May)	
Caryophyllaceae						Caryophyllaceae	
<input type="radio"/> <i>Cerastium fontanum</i> subsp. <i>vulgare</i>	Large Mouse-ear Chickweed	Herbaceous	- - -	-	ZA	Disturbed areas, grassy slopes, damp woodland, marshy ground (Spring)	
<input type="radio"/> <i>Cerastium glomeratum</i>	Sticky Mouse-ear Chickweed	Herbaceous	- - -	-	ZA	Dry hillsides, grassland, chaparral, disturbed areas (Spring)	
<input type="radio"/> <i>Herniaria hirsuta</i> var. <i>cinerea</i> (JM93: subsp.)	Ash-color Rupturewort	Herbaceous	- - -	-	ZA	Disturbed areas, alkaline hills, clay flats (Spring–fall)	

Caryophyllaceae		Sorted Alphabetically by Family, Scientific Name				Caryophyllaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Minuartia californica</i>	California Sandwort	Herbaceous	- - A2	-	NA	Gravelly, sandy slopes, grassy ridges, chaparral, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Minuartia douglasii</i>	Douglas' Stitchwort	Herbaceous	- - B	-	NA	Dry, rocky, slopes, flats in chaparral, oak and pine woodland, often serpentine (Spring–early summer)	
<input type="radio"/> <i>Minuartia pusilla</i>	Annual Sandwort	Herbaceous	- - A2	-	NA	Plains, open pine forest, chaparral slopes (Spring–summer)	
<input type="radio"/> <i>Moehringia macrophylla</i>	Big-leaf Sandwort	Herbaceous	- - A2	-	NP	Moist, shaded slopes, rocky ridges, summits, pine, oak forests, serpentine (Spring–summer)	
<input type="radio"/> <i>Polycarpon tetraphyllum</i>	Four-leaved Allseed	Herbaceous	- - -	-	ZA	Disturbed shaded areas, roadsides (Spring–fall)	
<input type="radio"/> <i>Sagina apetala</i>	Dwarf Pearlwort	Herbaceous	- - -	-	NA	Sandy disturbed areas, river bars, streamsides (Spring–early summer)	
<input type="radio"/> <i>Sagina decumbens subsp. occidentalis</i>	Western Pearlwort	Herbaceous	- - C	-	NA	Dry streams, chaparral, grassy areas, rock outcrops, vernal pools (Spring–early summer)	
<input type="radio"/> <i>Silene gallica</i>	Small-flower Catchfly	Herbaceous	- - -	-	ZA	Fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Silene laciniata subsp. californica (JM93: S. californica)</i>	California Pink	Herbaceous	- - B	-	NP	Chaparral, oak woodland, conifer forest, serpentine or not (Spring–summer)	
<input type="radio"/> <i>Spergula arvensis (JM93: subsp. arvensis)</i>	Stickwort	Herbaceous	- - -	-	ZA	Open slopes, pine woodland, sand dunes, fields, disturbed areas (Spring–early summer)	
<input type="radio"/> <i>Spergularia bocconi</i>	Boccone's Sand-spurry	Herbaceous	- - -	-	ZA	Salt marshes, alkaline areas, sandy soils (Spring)	
<input type="radio"/> <i>Spergularia macrotheca var. leucantha</i>	Short-style Sand-spurry	Herbaceous	- - A2	-	NP	Alkaline soils, floodplains, vernal pools, meadows, marshy ground (Apr–Jun)	
<input type="radio"/> <i>Spergularia macrotheca var. macrotheca</i>	Sticky Sand-spurry	Herbaceous	- - A2	-	NP	Salt flats, marshes, dunes, rocky outcrops, sandy or rocky coastal bluffs, gravelly ridges, alkaline fields (Spring–fall)	
<input type="radio"/> <i>Spergularia marina</i>	Saltmarsh Sand-spurrey	Herbaceous	- - -	-	NA	Mud flats, alkaline fields, sandy river bottoms, sandy coasts, salt marshes (Mar–Sep)	
<input type="radio"/> <i>Spergularia media var. media (JM93: no var.)</i>	Greater Sea-spurrey	Herbaceous	- - -	-	ZAP	Salt flats, salt marshes, sandy beaches (Summer–fall)	
<input type="radio"/> <i>Spergularia rubra</i>	Red Sand-spurry	Herbaceous	- - -	-	ZAP	Forest, meadows, mud flats, disturbed	
<input type="radio"/> <i>Spergularia sp.</i>	Sand-spurry	Herbaceous	- - -	-	-		
<input type="radio"/> <i>Stellaria media</i>	Common Chickweed	Herbaceous	- - -	-	ZA	Oak woodland, meadows, disturbed areas (Feb–Sep)	
<input type="radio"/> <i>Stellaria nitens</i>	Shiny Chickweed	Herbaceous	- - C	-	NA	Sand dunes, streambanks, open woodland, beneath boulders, disturbed areas (Spring)	
<input type="radio"/> <i>Stellaria pallida</i>	Lesser Chickweed	Herbaceous	- - -	-	ZA	Oak woodland, streambanks, grassy hills, flats, disturbed areas (Spring)	
Celastraceae						Celastraceae	
<input type="radio"/> <i>Euonymus occidentalis var. occidentalis</i>	Western Burning Bush	Woody	- - A1	-	NP	Shaded streambanks, canyons (Apr–Jun)	

Ceratophyllaceae		Sorted Alphabetically by Family, Scientific Name				Ceratophyllaceae		
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)		
Ceratophyllaceae								Ceratophyllaceae
<input type="radio"/> <i>Ceratophyllum demersum</i>	Hornwort	Herbaceous	- - -	-	NA	Common. Ditches, lakes, ponds, pools, slow watercourses; water 0.1–4 m deep, fresh to ± brackish, medium to high nutrient levels. (Jun–Aug)		
Chenopodiaceae								Chenopodiaceae
<input type="radio"/> <i>Allenrolfea occidentalis</i>	Iodine Bush	Woody	- - A2	-	NP	Flats, hummocks, in alkaline soils (Jun–Aug)		
<input type="radio"/> <i>Arthrocnemum subterminale</i> (JM93: <i>Salicornia</i>)	Glasswort Pickleweed	Herbaceous	- - A2	-	NP	Salt marshes, alkaline flats (Apr–Sep)		
<input type="radio"/> <i>Atriplex cordulata</i>	Heartscale	Herbaceous	- 1B.2 *A1	-	NA	Saline or alkaline soils (Jun–Jul)		
<input type="radio"/> <i>Atriplex depressa</i>	Brittlescale	Herbaceous	- 1B.2 *A2	-	NA	Alkaline or clay soils (Jun–Oct)		
<input type="radio"/> <i>Atriplex joaquinana</i> (JM93: <i>A. joaquiniana</i>)	San Joaquin Spearscale	Herbaceous	- 1B.2 *A2	-	NA	Alkaline soils (Apr–Sep)		
<input type="radio"/> <i>Atriplex lentiformis</i> (JM93: <i>subsp. lentiformis</i>)	Big Saltbush	Woody	- - A2	-	NP	Alkaline or saline washes, dry lakes, scrub (Jul–Oct)		
<input type="radio"/> <i>Atriplex leucophylla</i>	Beach Saltbush	Herbaceous	- - A2	-	NP	Sandy soils, dunes (Apr–Oct)		
<input type="radio"/> <i>Atriplex prostrata</i> (JM93: <i>A. triangularis</i>)	Fat-hen	Herbaceous	- - -	-	NA	Wet places, marshes (Apr–Oct)		
<input type="radio"/> <i>Atriplex rosea</i>	Tumbling Orach	Herbaceous	- - -	-	ZA	Common. Open, disturbed places, fields (Jul–Oct)		
<input type="radio"/> <i>Atriplex semibaccata</i>	Australian Saltbush	Herbaceous	- - -	M	ZP	Disturbed areas, scrub, woodland (Apr–Dec)		
<input type="radio"/> <i>Atriplex serenana</i> var. <i>serenana</i>	Bract Scale	Herbaceous	- - A1	-	NA	Alkaline flats, coastal bluffs (Apr–Oct)		
<input type="radio"/> <i>Atriplex</i> sp.	Saltbush / Orach	Herbaceous	- - -	-	-			
<input type="radio"/> <i>Bassia hyssopifolia</i>	Hyssop-leaf Bassia	Herbaceous	- - -	L	ZA	Disturbed sites, fields, roadsides (May–Nov)		
<input type="radio"/> <i>Chenopodium album</i>	Lamb's Quarters	Herbaceous	- - -	-	ZA	Common. Disturbed areas, fields (Jun–Oct)		
<input type="radio"/> <i>Chenopodium californicum</i>	California Goosefoot	Herbaceous	- - -	-	NP	Generally open sites, sandy to clay soils (Mar–Sep)		
<input type="radio"/> <i>Chenopodium desiccatum</i>	Aridland Goosefoot	Herbaceous	- - -	-	NA	Uncommon. Open places, scrub, conifer forest (Jul–Sep)		
<input type="radio"/> <i>Chenopodium macrospermum</i> (JM93: var. <i>halophilum</i>)	Coast Goosefoot	Herbaceous	- - -	-	ZA	Wet places, marshes (Jul–Oct)		
<input type="radio"/> <i>Chenopodium murale</i>	Wall Goosefoot	Herbaceous	- - -	-	ZA	Common. Disturbed areas, fields (All year)		
<input type="radio"/> <i>Chenopodium rubrum</i> var. <i>rubrum</i> (JM93: no var.)	Red Goosefoot	Herbaceous	- - A1	-	NA	Open, saline places, drying mudflats Aug–Oct		
<input type="radio"/> <i>Chenopodium</i> sp.	Goosefoot	Herbaceous	- - -	-	-			
<input type="radio"/> <i>Dysphania ambrosioides</i> (JM93: <i>Chenopodium</i>)	Mexican Tea	Herbaceous	- - -	-	ZAP	Disturbed places (Jul–Sep)		
<input type="radio"/> <i>Dysphania multifida</i> (JM93: <i>Chenopodium</i>)	Cutleaf Goosefoot	Herbaceous	- - -	-	ZA	Disturbed areas (Jul–Oct)		
<input type="radio"/> <i>Salicornia depressa</i> (JM93: <i>S. europaea</i>)	Slender Glasswort	Herbaceous	- - -	-	NA	Salt marshes, alkaline flats (Jul–Sep)		

Chenopodiaceae		Sorted Alphabetically by Family, Scientific Name				Chenopodiaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Salicornia pacifica</i> (JM93: <i>S. virginica</i>)	Pacific Pickleweed	Herbaceous	- - -	-	NP	Salt marshes, alkaline flats (Jul–Nov)	
<input type="radio"/> <i>Salsola soda</i>	Fleshy Russian Thistle	Herbaceous	- - -	M	ZA	Uppermost intertidal zone, saline or muddy flats, open areas in salt marshes (Jul–Oct)	
<input type="radio"/> <i>Salsola tragus</i>	Russian Thistle	Herbaceous	- - -	L	ZAP	Common. Disturbed places (Jul–Oct)	
<input type="radio"/> <i>Suaeda</i> sp.	Seepweed / Seablite	Woody	- - -	-	NP	--	
Cistaceae						Cistaceae	
<input type="radio"/> <i>Helianthemum scoparium</i>	Peak Rush-rose	Woody	- - B	-	NP	Dry sandy or rocky soil of hills, slopes, ridges (Feb–Sep)	
Commelinaceae						Commelinaceae	
<input type="radio"/> <i>Tradescantia fluminensis</i>	Spiderwort	Herbaceous	- - -	-	ZP	Shaded woodland, streambanks (Apr–Jul)	
Convolvulaceae						Convolvulaceae	
<input type="radio"/> <i>Calystegia malacophylla</i> subsp. <i>pedicellata</i>	Woolly Morning-glory	Herbaceous	- - A2	-	NP	Dry slopes, chaparral (Apr–Jul)	
<input type="radio"/> <i>Calystegia occidentalis</i> subsp. <i>occidentalis</i>	Western Morning-glory	Herbaceous	- - -	-	NP	Dry slopes, chaparral, pine woodland (May–Jul)	
<input type="radio"/> <i>Calystegia purpurata</i> subsp. <i>purpurata</i>	Climbing Morning-glory	Herbaceous	- - -	-	NP	Chaparral, coastal scrub (May–Jun)	
<input type="radio"/> <i>Calystegia sepium</i> subsp. <i>limnophila</i>	Hedge Bindweed	Herbaceous	- - A2	-	NP	Marshes, riverbanks (May–Jul)	
<input type="radio"/> <i>Calystegia</i> sp.	Morning-glory	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Calystegia subacaulis</i> subsp. <i>subacaulis</i>	Shortstem Morning-glory	Herbaceous	- - -	-	NP	Dry, open scrub or woodland (Apr–Jun)	
<input type="radio"/> <i>Convolvulus arvensis</i>	Bindweed	Herbaceous	- - -	-	ZP	Roadsides, open areas in many pl communities (Mar–Oct)	
<input type="radio"/> <i>Cressa truxillensis</i>	Alkali Weed	Herbaceous	- - -	-	NP	Saline and alkaline substrates (May–Oct)	
<input type="radio"/> <i>Cuscuta californica</i> var. <i>californica</i>	California Dodder	Herbaceous	- - A2	-	NA	On herbs, shrubs, roadsides, chaparral, grassland, yellow-pine forest (May–Sep)	
<input type="radio"/> <i>Cuscuta pacifica</i> var. <i>pacifica</i> (JM93: <i>C. salina</i> var. <i>major</i>)	Goldenthread	Herbaceous	- - B	-	NA	Gen on <i>Salicornia</i> , <i>Jaumea</i> in coastal salt marshes, tidal flats (Jul–Oct)	
<input type="radio"/> <i>Cuscuta subinclusa</i>	Canyon Dodder	Herbaceous	- - B	-	NP	Generally on herbs, shrubs, in forests near streams, river canyon bottoms, salt marshes (Mar–Oct(Dec?))	
Cornaceae						Cornaceae	
<input type="radio"/> <i>Cornus glabrata</i>	Brown Dogwood	Woody	- - A1	-	NP	Generally moist places (May–Jun)	
<input type="radio"/> <i>Cornus sericea</i> subsp. ?	American Dogwood	Woody	- - -	-	NP	--	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>occidentalis</i>	Western American Dogwood	Woody	- - C	-	NP	Generally moist places (May–Jul)	
<input type="radio"/> <i>Cornus sericea</i> subsp. <i>sericea</i>	American Creek Dogwood	Woody	- - C	-	NP	Generally moist habitats (May–Jul)	

Crassulaceae	Sorted Alphabetically by Family, Scientific Name					Crassulaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
Crassulaceae						Crassulaceae
<input type="radio"/> <i>Crassula aquatica</i>	Water Pygmy-weed	Herbaceous	- - C	- NA	Salt marshes, vernal pools, margins of lakes, ponds (Mar–Jun(Aug))	
<input type="radio"/> <i>Crassula connata</i>	Pygmy-weed	Herbaceous	- - -	- NA	Open areas (Feb–May)	
<input type="radio"/> <i>Crassula tillaea</i>	European Pygmy Weed	Herbaceous	- - -	- ZA	Open, gravelly sites (Feb–May)	
<input type="radio"/> <i>Dudleya cymosa subsp. cymosa</i>	Spreading Dudleya	Herbaceous	- - A1	- NP	Rocky outcrops, talus slopes, less often shaded canyon slopes (May–Jul)	
<input type="radio"/> <i>Dudleya cymosa subsp. paniculata</i>	Common / Hot Rock Dudleya	Herbaceous	- - B	- NP	Uncommon. Rocky outcrops, canyons (May–Jun)	
<input type="radio"/> <i>Dudleya farinosa</i>	Powdery Dudleya	Herbaceous	- - A1	- NP	Common: Coastal soils and cliffs (Jun–Aug)	
<input type="radio"/> <i>Sedella pentandra</i> (JM93: <i>Parvisedum pentandrum</i>)	Annual Stonecrop	Herbaceous	- - A2	- NA	Compacted ground, slate, shale, sandstone or serpentine outcrops (Mar–Jun)	
<input type="radio"/> <i>Sedum spathulifolium</i>	Broadleaf Stonecrop	Herbaceous	- - B	- NP	Outcrops, often in shade (Apr–Aug)	
Cucurbitaceae						Cucurbitaceae
<input type="radio"/> <i>Marah fabacea</i> (JM93: <i>M. fabaceus</i>)	California Man-root	Herbaceous	- - -	- NP	Streamsides, washes, shrubby open areas (Feb–Apr)	
<input type="radio"/> <i>Marah oregana</i> (JM93: <i>M. oreganus</i>)	Coast Man-root	Herbaceous	- - C	- NP	Shrubby or open areas, forest edges (Mar–May)	
<input type="radio"/> <i>Marah sp.</i>	Manroot	Herbaceous	- - -	- NP	- -	
Cupressaceae						Cupressaceae
<input type="radio"/> <i>Hesperocyparis macrocarpa</i> (JM93: <i>Cupressus</i>)	Monterey Cypress	Woody	- - -	N NP	Closed-cone-pine/cypress forests -	
<input type="radio"/> <i>Juniperus californica</i>	California Juniper	Woody	- - B	- NP	Dry slopes, flats, pinyon/juniper woodland -	
<input type="radio"/> <i>Sequoia sempervirens</i>	Redwood	Woody	- - B	- NP	Redwood forest -	
Cyperaceae						Cyperaceae
<input type="radio"/> <i>Bolboschoenus maritimus subsp. paludosus</i> (JM93: <i>Scirpus maritimus</i>)	Marine Bulrush	Grass-like	- - C	- NP	Common in CA. Brackish to saline coastal, inland marshes, shores (Summer)	
<input type="radio"/> <i>Bolboschoenus robustus</i> (JM93: <i>Scirpus</i>)	Seacoast Bulrush	Grass-like	- - -	- NP	Brackish to saline coastal marshes (Summer)	
<input type="radio"/> <i>Carex aquatilis var. dives</i>	Sitka Sedge	Grass-like	- - A2	- NP	Wet meadows, shores, fresh water (Jun–Sep)	
<input type="radio"/> <i>Carex barbarae</i>	Santa Barbara Sedge	Grass-like	- - B	- NP	Seasonally wet places (May–Aug)	
<input type="radio"/> <i>Carex brevicaulis</i>	Short-stemmed Sedge	Grass-like	- - A1	- NP	Coastal dunes, headlands, rocky or sandy soil (Apr–May)	
<input type="radio"/> <i>Carex globosa</i>	Round-fruited Sedge	Grass-like	- - A1	- NP	Well-drained soil of wooded areas, edges (Apr–Jun)	
<input type="radio"/> <i>Carex leporina</i> (JM93: <i>C. ovalis</i>)	Tracy Sedge	Grass-like	- - A1	- NP	Seasonally wet soil (May–Aug)	

Cyperaceae	Sorted Alphabetically by Family, Scientific Name					Cyperaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Carex leptopoda</i> (JM93: <i>C. deweyana</i> subsp. <i>leptopoda</i>)	Slender-footed Sedge	Grass-like	- - A1	-	NP	Moist soil, wooded areas (May–Aug)
<input type="radio"/> <i>Carex nebrascensis</i>	Nebraska Sedge	Grass-like	- - B	-	NP	Meadows, swamps (Jun–Sep)
<input type="radio"/> <i>Carex nudata</i>	Torrent Sedge	Grass-like	- - A2	-	NP	Rocky or sandy streambeds below high-water mark (Apr–Jul)
<input type="radio"/> <i>Carex obnupta</i>	Slough Sedge	Grass-like	- - A2	-	NP	Moist openings, shores, redwood forest (Jun–Sep)
<input type="radio"/> <i>Carex pellita</i> (JM93: <i>C. lanuginosa</i>)	Woolly Sedge	Grass-like	- - A1	-	NP	Generally marshy places, creekbanks (May–Sep)
<input type="radio"/> <i>Carex praegracilis</i>	Freeway Sedge	Grass-like	- - C	-	NP	Common. Often alkaline, ± moist places (Apr–Aug)
<input type="radio"/> <i>Carex serratodens</i>	Bifid Sedge	Grass-like	- - B	-	NP	Moist places (Apr–Jul)
<input type="radio"/> <i>Carex</i> sp.	Sedge	Grass-like	- - -	-	-	-
<input type="radio"/> <i>Carex subbracteata</i>	Small-bracted Sedge	Grass-like	- - B	-	NP	At least seasonally moist soil, grassland to open fores (Apr–Jun)
<input type="radio"/> <i>Carex tumulicola</i>	Foothill Sedge	Grass-like	- - C	-	NP	Meadows, open woodland (Apr–Jul)
<input type="radio"/> <i>Cyperus acuminatus</i>	Tapertip Flatsedge	Herbaceous	- - -	-	NA	Edges of temporary pools, ponds, streams (Jun–Oct)
<input type="radio"/> <i>Cyperus eragrostis</i>	Tall Nutsedge	Grass-like	- - -	-	NP	Vernal pools, streambanks (May–Nov)
<input type="radio"/> <i>Cyperus erythrorhizos</i>	Redroot Nutsedge	Grass-like	- - B	-	NA	Ditches, riverbanks, shores (Jul–Oct)
<input type="radio"/> <i>Cyperus odoratus</i>	Coarse Nutsedge	Grass-like	- - A1	-	NP	Wet disturbed soils (Jul–Oct)
<input type="radio"/> <i>Cyperus</i> sp.	Nutsedge	Grass-like	- - -	-	NAP	-
<input type="radio"/> <i>Cyperus strigosus</i>	False Nutsedge	Grass-like	- - A1	-	NP	Moist soils, pond margins, roadsides (Jul–Oct)
<input type="radio"/> <i>Eleocharis acicularis</i> var. <i>acicularis</i>	Needle Spikerush	Grass-like	- - C	-	NP	Common. Fresh wet soil to deeply submersed (Late spring–summer)
<input type="radio"/> <i>Eleocharis macrostachya</i>	Common Spikerush	Grass-like	- - -	-	NP	Common. Fresh to brackish wetland (Spring–summer)
<input type="radio"/> <i>Eleocharis parishii</i>	Parish Spikerush	Grass-like	- - A2	-	NP	Locally common. Fresh streams, vernal pools, foothill and riparian woodland, pine fores (Late spring–fall)
<input type="radio"/> <i>Eleocharis parvula</i>	Small Spikerush	Grass-like	- 4.3 *A1	-	NP	Brackish wet soil, coastal (Late winter–fall)
<input type="radio"/> <i>Eleocharis</i> sp.	Spikerush	Grass-like	- - -	-	NP	-
<input type="radio"/> <i>Isolepis cernua</i> (JM93: <i>Scirpus cernuus</i>)	Low Bulrush	Grass-like	- - B	-	NA	Sandy, sometimes brackish sea shores, bluffs, sand dunes, creeks, marshes (Late spring–winter)
<input type="radio"/> <i>Schoenoplectus acutus</i> var. <i>occidentalis</i> (JM93: <i>Scirpus</i>)	Common Tule	Grass-like	- - -	-	NP	Common. Marshes, shores, fens, shallow lakes, often emergent (Summer)
<input type="radio"/> <i>Schoenoplectus americanus</i> (JM93: <i>Scirpus</i>)	Olney's Three-square Bulrush	Grass-like	- - -	-	NP	Mineral-rich or brackish marshes, shores, fens, springs (Summer)
<input type="radio"/> <i>Schoenoplectus californicus</i> (JM93: <i>Scirpus</i>)	Southern Bulrush	Grass-like	- - -	-	NP	Common. Brackish to fresh marshes, shores (Spring–summer)

Cyperaceae		Sorted Alphabetically by Family, Scientific Name				Cyperaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Scirpus microcarpus</i>	Smallfruit Bulrush	Grass-like	- - -	-	NP	Marshes, wet meadows, streambanks, pond margins, sometimes weedy (Jun-Jul)
<input type="radio"/> <i>Scirpus, Isolepis, or Bolboschoenus sp. (JM93: Scirpus sp.)</i>	Bulrush	Grass-like	- - -	-	NAP	
Datisceae						Datisceae
<input type="radio"/> <i>Datisca glomerata</i>	Durango Root	Herbaceous	- - A2	-	NP	Dry streambeds or washes (May-Jul)
Dennstaedtiaceae						Dennstaedtiaceae
<input type="radio"/> <i>Pteridium aquilinum var. pubescens</i>	Bracken Fern	Fern-like	- - -	-	NP	Pastures, woodland, meadows, hillsides, partial to full sun -
Dipsacaceae						Dipsacaceae
<input type="radio"/> <i>Dipsacus sativus</i>	Fuller's Teasel	Herbaceous	- - -	M	ZB	Disturbed areas, fields, vacant lots, pastures (May-Jul)
<input type="radio"/> <i>Scabiosa atropurpurea</i>	Pincushion Flower	Herbaceous	- - -	-	ZA	Disturbed areas (Mar-Nov)
Dryopteridaceae						Dryopteridaceae
<input type="radio"/> <i>Dryopteris arguta</i>	Coastal Wood Fern	Fern-like	- - -	-	NP	Locally common. Open, wooded slopes, caves -
<input type="radio"/> <i>Polystichum munitum</i>	Western Sword Fern	Fern-like	- - -	-	NP	Common. Wooded hillsides, shaded slopes, rarely cliffs, outcrops -
Elaeagnaceae						Elaeagnaceae
<input type="radio"/> <i>Elaeagnus angustifolia</i>	Russian Olive	Woody	- - -	M	ZP	Uncommon. Disturbed, sometimes moist places (May-Jun)
Elatinaceae						Elatinaceae
<input type="radio"/> <i>Elatine californica</i>	California Waterwort	Herbaceous	- - A1	-	NA	Pools, ponds, rice fields, streambanks (Mar-Aug)
Equisetaceae						Equisetaceae
<input type="radio"/> <i>Equisetum arvense</i>	Common Horsetail	Fern-like	- - -	-	NP	Streambanks, wet meadows, springs, other wet, shaded places -
<input type="radio"/> <i>Equisetum hyemale subsp. affine</i>	Common Scouring Rush	Fern-like	- - -	-	NP	Streams, moist, sandy, gravelly areas -
<input type="radio"/> <i>Equisetum laevigatum</i>	Smooth Scouring Rush	Fern-like	- - B	-	NP	Moist, sandy or gravelly areas -
<input type="radio"/> <i>Equisetum sp.</i>	Scouring Rush	Fern-like	- - -	-	NP	- -
<input type="radio"/> <i>Equisetum telmateia subsp. braunii</i>	Giant Horsetail	Fern-like	- - -	-	NP	Streambanks, roadside ditches, seepage areas -
Ericaceae						Ericaceae
<input type="radio"/> <i>Arbutus menziesii</i>	Pacific Madrone	Woody	- - C	-	NP	Conifer, oak forests (Mar-May)

Ericaceae		Sorted Alphabetically by Family, Scientific Name					Ericaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Arctostaphylos auriculata</i>	Mount Diablo Manzanita	Woody	- 1B.3 *A2	-	NP	Sandstone, upland chaparral near coast. Mount Diablo and vicinity (Feb–Mar)	
<input type="radio"/> <i>Arctostaphylos crustacea</i> subsp. <i>crustacea</i> (JM93: <i>A. tomentosa</i> subsp. <i>crustacea</i>)	Brittle-leaf Manzanita	Woody	- - C	-	NP	Chaparral, conifer forest (Feb–Apr)	
<input type="radio"/> <i>Arctostaphylos glandulosa</i> subsp. <i>glandulosa</i>	Eastwood Manzanita	Woody	- - B	-	NP	Chaparral, conifer forest (Jan–Apr)	
<input type="radio"/> <i>Arctostaphylos glauca</i>	Big-berry Manzanita	Woody	- - C	-	NP	Rocky slopes, chaparral, woodland (Dec–Mar)	
<input type="radio"/> <i>Arctostaphylos manzanita</i> subsp. <i>laevigata</i>	Contra Costa Manzanita	Woody	- 1B.2 *A2	-	NP	Chaparral, rocky outcrops (Feb–May)	
<input type="radio"/> <i>Arctostaphylos manzanita</i> subsp. <i>manzanita</i>	Common / Giant Manzanita	Woody	- - C	-	NP	Chaparral, conifer forest (Feb–May)	
<input type="radio"/> <i>Arctostaphylos pallida</i>	Pallid Manzanita	Woody	FT/CE 1B.1 *A1	-	NP	Siliceous shales, slopes, ridges, chaparral (Jan–Mar)	
<input type="radio"/> <i>Arctostaphylos</i> sp.	Manzanita	Woody	- - -	-	NP	- -	
<input type="radio"/> <i>Gaultheria shallon</i>	Salal	Woody	- - A1	-	NP	Moist forest margins (Apr–Jul)	
<input type="radio"/> <i>Vaccinium ovatum</i>	California Huckleberry	Woody	- - B	-	NP	Edges, clearings in conifer forest (Mar–May)	
Euphorbiaceae							Euphorbiaceae
<input type="radio"/> <i>Chamaesyce maculata</i>	Spotted Spurge	Herbaceous	- - -	-	ZA	Disturbed places (Apr–Oct)	
<input type="radio"/> <i>Chamaesyce ocellata</i> subsp. <i>ocellata</i>	Valley Spurge	Herbaceous	- - B	-	NA	Dry, sandy places (May–Sep)	
<input type="radio"/> <i>Chamaesyce serpyllifolia</i> subsp. <i>serpyllifolia</i>	Thyme-leafed Spurge	Herbaceous	- - B	-	NA	Common. Dry habitats (Aug–Oct)	
<input type="radio"/> <i>Croton californicus</i>	Croton	Herbaceous	- - A2	-	NP	Sandy soils, dunes, washes (Apr–Jul)	
<input type="radio"/> <i>Croton setigerus</i> (JM93: <i>Eremocarpus</i>)	Turkey-mullein	Herbaceous	- - -	-	NA	Dry, open, often disturbed areas (May–Oct)	
<input type="radio"/> <i>Euphorbia crenulata</i>	Chinese Caps	Herbaceous	- - -	-	NA	Common. Dry places (Mar–Aug)	
<input type="radio"/> <i>Euphorbia lathyris</i>	Caper Spurge	Herbaceous	- - -	-	ZAB	Disturbed areas (All year)	
<input type="radio"/> <i>Euphorbia oblongata</i>	European Spurge	Herbaceous	- - -	L	ZP	Disturbed areas (Jun–Aug)	
<input type="radio"/> <i>Euphorbia peplus</i>	Petty Spurge	Herbaceous	- - -	-	ZA	Common. Disturbed areas (Feb–Aug)	
<input type="radio"/> <i>Euphorbia spathulata</i>	Wart Spurge	Herbaceous	- - -	-	NA	Open, gen disturbed places (Mar–Jun)	
Fabaceae							Fabaceae
<input type="radio"/> <i>Acacia cyclops</i>	Western Coastal Wattle	Woody	- - -	-	ZP	Uncommon. Disturbed areas, coastal dunes (Dec–Mar)	
<input type="radio"/> <i>Acacia dealbata</i>	Silver Wattle	Woody	- - -	M	ZP	Locally common. Disturbed areas, often roadsides (Feb–Apr)	
<input type="radio"/> <i>Acacia longifolia</i>	Sydney Golden Wattle	Woody	- - -	-	ZP	Uncommon. Disturbed places, especially sandy coastal areas (Jan–Apr)	
<input type="radio"/> <i>Acacia melanoxylon</i>	Blackwood Acacia	Woody	- - -	L	ZP	Uncommon. Disturbed areas (Feb–Mar)	
<input type="radio"/> <i>Acacia</i> sp.	Acacia	Woody	- - -	-	ZP	- -	
<input type="radio"/> <i>Acmispon americanus</i> var. <i>americanus</i> (JM93: <i>Lotus purshianus</i> var. <i>purshianus</i>)	Spanish Clover	Herbaceous	- - -	-	NA		

Fabaceae		Sorted Alphabetically by Family, Scientific Name					Fabaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Acmispon brachycarpus</i> (JM93: <i>Lotus humistratus</i>)	Colchita	Herbaceous	- - -	-	NA	Abundant. Grassland, oak and pine woodland, desert flats and mtns, roadsides (Mar–Jun)	
<input type="radio"/> <i>Acmispon glaber</i> var. <i>glaber</i> (JM93: <i>Lotus scoparius</i> var. <i>scoparius</i>)	Deerweed	Herbaceous	- - -	-	NP	Chaparral, roadsides, coastal sands; common (Mar–Aug)	
<input type="radio"/> <i>Acmispon</i> or <i>Hosackia</i> sp. (JM93: <i>Lotus</i> sp.)	Lotus	Herbaceous	- - -	-	-	-	
<input type="radio"/> <i>Acmispon parviflorus</i> (JM93: <i>Lotus micranthus</i>)	Small-flower Lotus	Herbaceous	- - -	-	NA	Abundant. Coastal bluffs to oak/pine or fir woodland, open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Acmispon strigosus</i> (JM93: <i>Lotus</i>)	Hairy Lotus	Herbaceous	- - A2	-	NP	Coastal scrub, chaparral, foothills, deserts, roadsides, other disturbed area (Mar–Jun)	
<input type="radio"/> <i>Acmispon wrangelianus</i> (JM93: <i>Lotus</i>)	California Lotus	Herbaceous	- - -	-	NA	Abundant. Coastal bluffs, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Astragalus asymmetricus</i>	San Joaquin Milkvetch	Herbaceous	- - B	-	NP	Grassy areas, open woodland, disturbed sites (Apr–Jul)	
<input type="radio"/> <i>Astragalus didymocarpus</i> var. <i>didymocarpus</i>	Two-seed Milkvetch	Herbaceous	- - A1	-	NA	Grassy areas (Feb–May)	
<input type="radio"/> <i>Astragalus gambelianus</i>	Gambel Milkvetch	Herbaceous	- - -	-	NA	Open, grassy areas, scrub (Mar–Jul)	
<input type="radio"/> <i>Astragalus oxyphysus</i>	Stanislaus Milkvetch	Herbaceous	- - A1?	-	NP	Arid grassland, scrub (Mar–Jun)	
<input type="radio"/> <i>Astragalus</i> sp.	Milkvetch	Herbaceous	- - -	-	NAP	-	
<input type="radio"/> <i>Cercis occidentalis</i>	Western Redbud	Woody	- - A1	-	NP	Dry, shrubby slopes, canyons, streambanks, chaparral, foothill woodland, yellow-pine forest (Mar–May)	
<input type="radio"/> <i>Cytisus scoparius</i>	Scotch Broom	Woody	- - -	H	ZP	Common. Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Genista monspessulana</i>	French Broom	Woody	- - -	H	ZP	Common. Disturbed places. (Mar–Jun)	
<input type="radio"/> <i>Glycyrrhiza lepidota</i>	Wild Licorice	Herbaceous	- - C	-	NP	In colonies, moist, gen open, disturbed areas incl streambanks, roadsides, alkaline soils or not (May–Jul)	
<input type="radio"/> <i>Hoita macrostachya</i>	Leather Root	Herbaceous	- - B	-	NP	Streamsides, marshes, spring-moist places (May–Aug)	
<input type="radio"/> <i>Lathyrus jepsonii</i> var. <i>californicus</i>	Bluff Pea	Herbaceous	- - B	-	NP	Forest, open areas (Apr–Aug)	
<input type="radio"/> <i>Lathyrus jepsonii</i> var. <i>jepsonii</i>	Delta Tule Pea	Herbaceous	- 1B.2 *A2	-	NP	Coastal, estuarine marshes (Apr–Aug)	
<input type="radio"/> <i>Lathyrus latifolius</i>	Perennial Sweet Pea	Herbaceous	- - -	-	ZP	Disturbed areas, esp roadsides (Apr–May)	
<input type="radio"/> <i>Lathyrus</i> sp.	Pea	Herbaceous	- - -	-	-	-	
<input type="radio"/> <i>Lathyrus vestitus</i> var. <i>vestitus</i>	Pale Purple Pacific Pea	Herbaceous	- - -	-	NP	North: Conifer forest. South: chaparral & oak woodland (Feb–Jul)	
<input type="radio"/> <i>Lotus corniculatus</i>	Bird's-foot Trefoil	Herbaceous	- - -	-	ZP	Open, disturbed areas (Jun–Sep)	
<input type="radio"/> <i>Lupinus adsurgens</i>	Drew's Silky Lupine	Herbaceous	- - -	-	NP	Dry slopes, montane forest (May–Jul)	
<input type="radio"/> <i>Lupinus affinis</i>	Sky Lupine	Herbaceous	- - A1	-	NA	Uncommon. Open areas (Mar–May)	
<input type="radio"/> <i>Lupinus albifrons</i> var. ?	Blue Bush / Silver Lupine	Woody	- - -	-	NP	-	

Fabaceae	Sorted Alphabetically by Family, Scientific Name					Fabaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>albifrons</i>	Bush Lupine	Woody	- - -	- NP	Common. Chaparral, foothill woodland (Mar–Jun)	
<input type="radio"/> <i>Lupinus albifrons</i> var. <i>collinus</i>	Bay Area Silver Lupine	Woody	- - -	- NP	Cliffs, forest openings (Mar–Jun)	
<input type="radio"/> <i>Lupinus arboreus</i>	Yellow Bush Lupine	Woody	- - A2	N NP	Coastal bluffs, dunes, or more inland (Apr–Jul)	
<input type="radio"/> <i>Lupinus bicolor</i>	Miniature Lupine	Herbaceous	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus concinnus</i>	Bajada Lupine	Herbaceous	- - A1	- NA	Common. Open or disturbed areas, burns (Mar–May)	
<input type="radio"/> <i>Lupinus formosus</i> var. <i>formosus</i>	Summer Lupine	Herbaceous	- - -	- NP	Dry clay soils, grassland, open areas under pines, gen in valleys (Apr–Sep)	
<input type="radio"/> <i>Lupinus latifolius</i> var. <i>latifolius</i>	Broadleaf Lupine	Herbaceous	- - -	- NP	Moist areas, open woodland (Apr–Jul)	
<input type="radio"/> <i>Lupinus luteolus</i>	Butter Lupine	Herbaceous	- - A1	- NA	Clearings, open or disturbed areas (May–Aug)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>densiflorus</i>	Gully Lupine	Herbaceous	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Apr–Jun)	
<input type="radio"/> <i>Lupinus microcarpus</i> var. <i>microcarpus</i>	Chick Lupine	Herbaceous	- - -	- NA	Abundant. Open or disturbed areas, occ seeded on roadbanks (Mar–Jun)	
<input type="radio"/> <i>Lupinus nanus</i>	Sky Lupine	Herbaceous	- - -	- NA	Abundant. Open or disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Lupinus pachylobus</i>	Big Pod Lupine	Herbaceous	- - B	- NA	Uncommon. Open or disturbed areas (Mar–May)	
<input type="radio"/> <i>Lupinus</i> sp.	Lupine	Herbaceous	- - -	- NAP	- -	
<input type="radio"/> <i>Lupinus succulentus</i>	Arroyo Lupine	Herbaceous	- - -	- NA	Abundant. Open or disturbed areas, often seeded on roadbanks (Feb–May)	
<input type="radio"/> <i>Lupinus variicolor</i>	Varied Lupine	Herbaceous	- - A1	- NP	Coastal terraces, beaches (Apr–Jul)	
<input type="radio"/> <i>Medicago arabica</i>	Spotted Bur Clover	Herbaceous	- - -	- ZA	Disturbed and agricultural areas, fields, woodland (Mar–Jun)	
<input type="radio"/> <i>Medicago lupulina</i>	Black Medick	Herbaceous	- - -	- ZA	Disturbed and agricultural areas, forest, mtns (Apr–Jul)	
<input type="radio"/> <i>Medicago polymorpha</i>	California Burclover	Herbaceous	- - -	L ZA	Common. Chaparral, oak woodland, streambanks, roadsides, disturbed areas (Mar–Jul)	
<input type="radio"/> <i>Medicago sativa</i>	Alfalfa	Herbaceous	- - -	- ZP	Disturbed, agricultural areas (Apr–Oct)	
<input type="radio"/> <i>Melilotus albus</i> (JM93: <i>M. alba</i>)	White Sweetclover	Herbaceous	- - -	- ZAB	Locally abundant. Pastures, open disturbed sites (May–Sep)	
<input type="radio"/> <i>Melilotus indicus</i> (JM93: <i>M. indica</i>)	Sourclover	Herbaceous	- - -	- ZAB	Open, disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Melilotus officinalis</i>	Yellow Sweetclover	Herbaceous	- - -	- ZAB	Open fields, disturbed sites, cultivated (May–Aug)	
<input type="radio"/> <i>Pediomelum californicum</i>	Indian Breadroot	Herbaceous	- - A1	- NP	Open chaparral, woodland (Apr–Jul)	
<input type="radio"/> <i>Pickeringia montana</i> var. <i>montana</i>	Chaparral Pea	Woody	- - B	- NP	Chaparral, open woodland (May–Aug)	
<input type="radio"/> <i>Robinia pseudoacacia</i>	Black Locust	Woody	- - -	L ZP	Locally common near abandoned houses, roadsides, canyon slopes, streambanks (May–Jun)	
<input type="radio"/> <i>Rupertia physodes</i>	California Tea	Herbaceous	- - -	- NP	Woodland (May–Sep)	
<input type="radio"/> <i>Spartium junceum</i>	Spanish Broom	Woody	- - -	H ZP	Disturbed areas (Apr–Jun)	

Fabaceae		Sorted Alphabetically by Family, Scientific Name					Fabaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Trifolium albopurpureum</i> (JM93: var. <i>albopurpureum</i>)	Rancheria Clover	Herbaceous	- - C	-	NA	Abundant. Dunes, grassland, wet meadows, slopes, disturbed areas, etc (Mar–Jun)	
<input type="radio"/> <i>Trifolium angustifolium</i>	Narrow-leaved Clover	Herbaceous	- - -	-	ZA	Disturbed areas (Late spring)	
<input type="radio"/> <i>Trifolium barbigerum</i> (JM93: var. <i>barbigerum</i>)	Bearded Clover	Herbaceous	- - B	-	NP	Wet meadows, open, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Trifolium bifidum</i> var. ?	Pinole Clover	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>bifidum</i>	Pinole Clover	Herbaceous	- - B	-	NA	Open, grassy areas, oak chaparral, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium bifidum</i> var. <i>decipiens</i>	Deceiving Clover	Herbaceous	- - -	-	NA	Open, grassy areas, forest (Apr–Jun)	
<input type="radio"/> <i>Trifolium campestre</i>	Hop Clover	Herbaceous	- - -	-	ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium ciliolatum</i>	Foothill Clover	Herbaceous	- - -	-	NA	Locally common. Grassland, chaparral, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>amplectens</i>	Fused-bract Pale Sack Clover	Herbaceous	- - -	-	NA	Grassland, coastal woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>depauperatum</i>	Dwarf Sack Clover	Herbaceous	- - B	-	NA	Wet meadows, grassland, roadsides, open spring-moist, heavy soils (Mar–May)	
<input type="radio"/> <i>Trifolium depauperatum</i> var. <i>truncatum</i>	Truncate Sack Clover	Herbaceous	- - -	-	NA	Grassy flats, disturbed slopes, openings in woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium dichotomum</i> (JM93: <i>T. albopurpureum</i> var. <i>dichotomum</i>)	Branched Indian Clover	Herbaceous	- - B	-	NA	Coastal dunes, open slopes, meadows, oak woodland, disturbed areas (Apr–Jun)	
<input type="radio"/> <i>Trifolium dubium</i>	Little Hop Clover	Herbaceous	- - -	-	ZA	Agricultural, disturbed areas, lawns (Spring)	
<input type="radio"/> <i>Trifolium fragiferum</i>	Strawberry Clover	Herbaceous	- - -	-	ZP	Roadsides, gen in saline soil (Late spring)	
<input type="radio"/> <i>Trifolium fucatum</i>	Bull Clover	Herbaceous	- - C	-	NA	Locally abundant. Moist, open grassland, marshes, roadsides, occasionally saline or serpentine soils (Apr–Jun)	
<input type="radio"/> <i>Trifolium glomeratum</i>	Clustered Clover	Herbaceous	- - -	-	ZA	Uncommon. Disturbed areas (Mar–May)	
<input type="radio"/> <i>Trifolium gracilentum</i> (JM93: var. <i>gracilentum</i>)	Pinpoint Clover	Herbaceous	- - -	-	NA	Open, disturbed places, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trifolium grayi</i> (JM93: <i>T. barbigerum</i> var. <i>andrewsii</i>)	Gray's Clover	Herbaceous	- - A1?	-	NA	Wet meadows, foothill slopes, pine woodland (Apr–Jun)	
<input type="radio"/> <i>Trifolium hirtum</i>	Rose Clover	Herbaceous	- - -	M	ZA	Disturbed areas, roadsides (Apr–May)	
<input type="radio"/> <i>Trifolium incarnatum</i>	Crimson Clover	Herbaceous	- - -	-	ZA	Uncommon. Disturbed areas (May–Aug)	
<input type="radio"/> <i>Trifolium macraei</i>	Macrae's Clover	Herbaceous	- - A2	-	NA	Disturbed areas, dunes (Mar–May)	
<input type="radio"/> <i>Trifolium microcephalum</i>	Small-head Clover	Herbaceous	- - -	-	NA	Streambanks, moist, disturbed areas, roadsides, serpentine, conifer forest (Apr–Aug)	
<input type="radio"/> <i>Trifolium microdon</i>	Thimble Clover	Herbaceous	- - -	-	NA	Common locally. Open, moist or dry, gen disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Trifolium obtusiflorum</i>	Clammy Clover	Herbaceous	- - B	-	NA	Moist disturbed areas, gravel bars, marshes (Apr–Jul)	
<input type="radio"/> <i>Trifolium oliganthum</i>	Few-flowered Clover	Herbaceous	- - C	-	NA	Woody or shrubby slopes, roadsides (Mar–Jun)	

Fabaceae		Sorted Alphabetically by Family, Scientific Name					Fabaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Trifolium olivaceum</i> (JM93: <i>T. albopurpureum</i> var. <i>olivaceum</i>)	Olive Clover	Herbaceous	- - B	-	NA	Grassy slopes, valley meadows, disturbed areas (Apr–May)	
<input type="radio"/> <i>Trifolium pratense</i>	Red Clover	Herbaceous	- - -	-	ZP	Disturbed areas (Apr–Oct)	
<input type="radio"/> <i>Trifolium repens</i>	White Clover	Herbaceous	- - -	-	ZP	Disturbed areas (Feb–Dec)	
<input type="radio"/> <i>Trifolium sp.</i>	Clover	Herbaceous	- - -	-	--		
<input type="radio"/> <i>Trifolium subterraneum</i>	Subterranean Clover	Herbaceous	- - -	-	ZA	Meadows, roadsides, disturbed areas (Mar–Apr)	
<input type="radio"/> <i>Trifolium variegatum</i> var. ? (JM93: no var.)	White-tip Clover	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Trifolium willdenovii</i>	Tomcat Clover	Herbaceous	- - -	-	NA	Abundant. Disturbed, gen spring-moist, heavy soils, occas serpentine (Mar–Jun)	
<input type="radio"/> <i>Trifolium wormskioldii</i>	Cow Clover	Herbaceous	- - A2	-	NA	Beaches to mtn meadows, ridges, generally open moist or marshy places (May–Oct)	
<input type="radio"/> <i>Vicia americana</i> subsp. <i>americana</i> (JM93: var. <i>americana</i>)	American Vetch	Herbaceous	- - -	-	NP	Generally open, moist forest, along streams, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia benghalensis</i>	One-sided Purple Vetch	Herbaceous	- - -	-	ZP	Grassland, roadsides, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia disperma</i>	Two-seeded Vetch	Herbaceous	- - -	-	ZA	Roadsides, urban weed (Mar–May)	
<input type="radio"/> <i>Vicia gigantea</i>	Giant Vetch	Herbaceous	- - C	-	NP	Coastal shrub, coastal forest, chaparral (Mar–Aug)	
<input type="radio"/> <i>Vicia hassei</i>	Slender Vetch	Herbaceous	- - A2	-	NA	Coastal scrub, chaparral, oak woodland (Mar–May)	
<input type="radio"/> <i>Vicia ludoviciana</i> subsp. <i>ludoviciana</i> (JM93: var. <i>ludoviciana</i>)	Deerpea Vetch	Herbaceous	- - -	-	NA	Slopes, canyons, streambanks, in grassland, coastal-sage scrub, chaparral, oak woodland, riparian woodland (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. ?	Narrowleaf / Spring Vetch	Herbaceous	- - -	-	ZA	--	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>nigra</i>	Narrow-leaved Vetch	Herbaceous	- - -	-	ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sativa</i> subsp. <i>sativa</i>	Spring Vetch	Herbaceous	- - -	-	ZA	Roadsides, disturbed areas, grassland, open areas in oak and riparian woodlands (Mar–Jun)	
<input type="radio"/> <i>Vicia sp.</i>	Vetch	Herbaceous	- - -	-	--		
<input type="radio"/> <i>Vicia villosa</i> subsp. ?	Vetch	Herbaceous	- - -	-	ZAB	--	
<input type="radio"/> <i>Vicia villosa</i> subsp. <i>varia</i>	Sparsely Hairy Vetch	Herbaceous	- - -	-	ZAB	Grassland, roadside, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Vicia villosa</i> subsp. <i>villosa</i>	Hairy Vetch	Herbaceous	- - -	-	ZAB	Grassland, roadside, disturbed areas (May–Jul)	
Fagaceae						Fagaceae	
<input type="radio"/> <i>Chrysolepis chrysophylla</i> var. <i>minor</i>	Golden Chinquapin	Woody	- - A2	-	NP	Conifer forest, closed-cone-pine forest, chaparral (Jun–Sep)	
<input type="radio"/> <i>Quercus agrifolia</i> var. <i>agrifolia</i>	Coast Live Oak	Woody	- - A2	-	NP	Valleys, slopes, mixed-evergreen forest, woodland (Mar–Apr)	
<input type="radio"/> <i>Quercus berberidifolia</i>	Scrub Oak	Woody	- - C	-	NP	Dry slopes, chaparral (Feb–Apr)	

Fagaceae		Sorted Alphabetically by Family, Scientific Name					Fagaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Quercus chrysolepis</i>	Canyon Live Oak	Woody	- - -	-	NP	Canyons, shaded slopes, chaparral, mixed-evergreen forest, woodland (Apr–May)	
<input type="radio"/> <i>Quercus douglasii</i>	Blue Oak	Woody	- - -	-	NP	Dry slopes, interior foothills, woodland (Apr–May)	
<input type="radio"/> <i>Quercus durata</i> var. <i>durata</i>	Leather Oak	Woody	- - A2	-	NP	Chaparral, generally serpentine (Apr–May)	
<input type="radio"/> <i>Quercus garryana</i> var. <i>garryana</i>	Oregon / Garry Oak	Woody	- - B	-	NP	Slopes, mixed-evergreen or conifer forest (Apr–Jun)	
<input type="radio"/> <i>Quercus kelloggii</i>	California Black Oak	Woody	- - -	-	NP	Slopes, valleys, woodland, conifer forest (Apr–May)	
<input type="radio"/> <i>Quercus lobata</i>	Valley Oak	Woody	- - B	-	NP	Slopes, valleys, savanna (Mar–Apr)	
<input type="radio"/> <i>Quercus palmeri</i>	Palmer Oak	Woody	- - A2	-	NP	Uncommon. Rocky slopes, flats (Apr–May)	
<input type="radio"/> <i>Quercus</i> sp.	Oak	Woody	- - -	-	NP	- -	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>frutescens</i>	Dwarf Interior Live Oak	Woody	- - C	-	NP	Valleys, chaparral (Mar–May)	
<input type="radio"/> <i>Quercus wislizeni</i> var. <i>wislizeni</i>	Interior Live Oak	Woody	- - -	-	NP	Interior canyons, slopes, pine/oak woodland (Mar–May)	
Frankeniaceae							Frankeniaceae
<input type="radio"/> <i>Frankenia salina</i>	Alkali Heath	Herbaceous	- - -	-	NP	Salt marshes, alkali flats (Apr–Sep)	
Garryaceae							Garryaceae
<input type="radio"/> <i>Garrya elliptica</i>	Coast Silk Tassel	Woody	- - B	-	NP	Seacliffs, sand dunes, chaparral, foothill-pine woodland (Jan–Mar)	
<input type="radio"/> <i>Garrya fremontii</i>	Green-leaf Silk Tassel	Woody	- - A1	-	NP	Chaparral, foothill woodland, montane forest (Jan–Apr)	
Gentianaceae							Gentianaceae
<input type="radio"/> <i>Cicendia quadrangularis</i>	Timwort	Herbaceous	- - B	-	NA	Open places (Mar–May)	
<input type="radio"/> <i>Zeltnera davyi</i> (JM93: <i>Centaurium</i>)	Davy's Centaury	Herbaceous	- - B	-	NA	Moist coastal bluffs, dunes, open forest (May–Aug)	
<input type="radio"/> <i>Zeltnera muehlenbergii</i> (JM93: <i>Centaurium</i>)	Monterey Centaury	Herbaceous	- - -	-	NA	Moist coastal bluffs, forest openings (Jun–Aug)	
<input type="radio"/> <i>Zeltnera</i> sp. (JM93: <i>Centaurium</i>)	Centaury	Herbaceous	- - -	-	NA	- -	
Geraniaceae							Geraniaceae
<input type="radio"/> <i>Erodium botrys</i>	Long-beaked Filaree	Herbaceous	- - -	-	ZA	Dry, open or disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Erodium brachycarpum</i>	Southern European Filaree	Herbaceous	- - -	-	ZA	Dry, open or disturbed sites (Mar–Jul)	
<input type="radio"/> <i>Erodium cicutarium</i>	Redstem Filaree	Herbaceous	- - -	L	ZA	Open, disturbed sites, grassland, scrub (Feb–Sep)	
<input type="radio"/> <i>Erodium moschatum</i>	Greenstem Filaree	Herbaceous	- - -	-	ZA	Open, disturbed sites (Feb–Sep)	
<input type="radio"/> <i>Geranium carolinianum</i>	Carolina Geranium	Herbaceous	- - -	-	NA	Open to shaded sites, grassland, scrub, forest (Feb–Aug)	
<input type="radio"/> <i>Geranium dissectum</i>	Cut-leaved Geranium	Herbaceous	- - -	M	ZA	Open, disturbed sites (Mar–Jul)	

Geraniaceae		Sorted Alphabetically by Family, Scientific Name				Geraniaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Geranium molle</i>	Hairy Dove's Foot Geranium	Herbaceous	- - -	-	ZA	Open to shaded sites, disturbed ground (Feb–Aug)	
<input type="radio"/> <i>Geranium robertianum</i>	Herb Robert	Herbaceous	- - -	-	ZAB	Open to shaded sites (Apr–Sep)	
Grossulariaceae						Grossulariaceae	
<input type="radio"/> <i>Ribes amarum</i>	Bitter Gooseberry	Woody	- - A2	-	NP	Chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes californicum</i> var. <i>californicum</i>	Hillside Gooseberry	Woody	- - -	-	NP	Forest openings, woodland (Feb–Mar)	
<input type="radio"/> <i>Ribes divaricatum</i> var. <i>pubiflorum</i>	Straggle Gooseberry	Woody	- - B	-	NP	Uncommon. Coastal bluffs, forest edges (Mar–May)	
<input type="radio"/> <i>Ribes malvaceum</i> var. <i>malvaceum</i>	Chaparral Currant	Woody	- - B	-	NP	Chaparral, oak woodland (Oct–Apr)	
<input type="radio"/> <i>Ribes menziesii</i> var. <i>menziesii</i> (JM93: no var.)	Canyon Gooseberry	Woody	- - -	-	NP	Common. Forest openings, chaparral (Feb–Apr)	
<input type="radio"/> <i>Ribes sanguineum</i> var. <i>glutinosum</i>	Red-flowering Currant	Woody	- - C	-	NP	Many habitats (Feb–Apr)	
<input type="radio"/> <i>Ribes</i> sp.	Gooseberry / Currant	Woody	- - -	-	NP	--	
<input type="radio"/> <i>Ribes speciosum</i>	Fuchsia-flowered Gooseberry	Woody	- - A1	-	NP	Coastal-sage scrub, chaparral (Jan–May)	
Haloragaceae						Haloragaceae	
<input type="radio"/> <i>Myriophyllum sibiricum</i>	Siberian Water-milfoil	Herbaceous	- - -	-	NP	Ponds, streams, lakes (Jun–Sep)	
<input type="radio"/> <i>Myriophyllum</i> sp.	Water-milfoil	Herbaceous	- - -	-	--		
Hydrocharitaceae						Hydrocharitaceae	
<input type="radio"/> <i>Najas marina</i>	Holly-leaved Water-nymph	Herbaceous	- - -	-	NA	Ponds, lakes, marshes, rivers (Jul–Aug)	
Hypericaceae						Hypericaceae	
<input type="radio"/> <i>Hypericum perforatum</i>	Klamathweed	Herbaceous	- - -	M	ZP	Open, disturbed areas in many plant communities (May–Aug)	
Iridaceae						Iridaceae	
<input type="radio"/> <i>Iris douglasiana</i>	Douglas Iris	Herbaceous	- - A2	-	NP	Common. Grassy places, esp near coast (May–Jul)	
<input type="radio"/> <i>Iris longipetala</i>	Coast Iris	Herbaceous	- 4.2 *A1	-	NP	Moist, coastal prairie or open coastal forest (Mar–Jun)	
<input type="radio"/> <i>Iris pseudacorus</i>	Yellow Flag	Herbaceous	- - -	L	ZP	Common. Pond margins, estuaries (Apr–Jun)	
<input type="radio"/> <i>Iris</i> sp.	Iris	Herbaceous	- - -	-	--		
<input type="radio"/> <i>Sisyrinchium bellum</i>	Western Blue-eyed-grass	Herbaceous	- - -	-	NP	Common. Open, gen moist, grassy areas, woodland (Mar–May)	
Isoetaceae						Isoetaceae	
<input type="radio"/> <i>Isoetes howellii</i>	Howell Quillwort	Fern-like	- - A2	-	NP	Vernal pools, lake margins (Late spring-summer)	
Juglandaceae						Juglandaceae	
<input type="radio"/> <i>Juglans hindsii</i> (JM93: <i>californica</i> var. <i>hindsii</i>)	N. California Black Walnut	Woody	- 1B.1 *A2	-	NP	Along streams, disturbed slopes (Apr–May)	

Juncaceae		Sorted Alphabetically by Family, Scientific Name				Juncaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
Juncaceae						Juncaceae
<input type="radio"/> <i>Juncus balticus</i> subsp. <i>ater</i> (JM93: no subsp.)	Baltic Rush	Grass-like	- - -	- NP	Moist to ± dry sites (Jul–Nov)	
<input type="radio"/> <i>Juncus bufonius</i> var. ?	Toad Rush	Grass-like	- - -	- NA	--	
<input type="radio"/> <i>Juncus bufonius</i> var. <i>bufonius</i>	Toad Rush	Grass-like	- - -	- NA	Damp sunny ground, gen disturbed (May–Sep)	
<input type="radio"/> <i>Juncus bufonius</i> var. <i>congestus</i>	Clustered Toad Rush	Grass-like	- - -	- NA	Drying pools, shores, disturbed ground, gen saline (Apr–Aug)	
<input type="radio"/> <i>Juncus dubius</i>	Mariposa Rush	Grass-like	- - A1	- NP	Wet places (Jul–Sep)	
<input type="radio"/> <i>Juncus effusus</i> subsp. <i>pacificus</i> (JM93: var.)	Pacific Rush	Grass-like	- - -	- NP	Seeps, shores, marshes, generally damp sunny ground (May–Oct)	
<input type="radio"/> <i>Juncus lescurii</i> (JM93: <i>J. lesueurii</i>)	San Francisco Rush	Grass-like	- - A2	- NP	Salt or freshwater marshes, shores of creeks and lakes (Aug–Sep)	
<input type="radio"/> <i>Juncus mexicanus</i>	Mexican Rush	Grass-like	- - B	- NP	Common. Coast to montane meadows (Aug–Nov)	
<input type="radio"/> <i>Juncus occidentalis</i>	Western Rush	Grass-like	- - -	- NP	Moist gen sunny areas (May–Sep)	
<input type="radio"/> <i>Juncus oxymiris</i>	Pointed Rush	Grass-like	- - A2	- NP	Uncommon. Swales, montane meadows, damp, sunny ground (Jul–Aug)	
<input type="radio"/> <i>Juncus patens</i>	Spreading Rush	Grass-like	- - -	- NP	Marshy places, creeks, seeps (Jun–Oct)	
<input type="radio"/> <i>Juncus phaeocephalus</i> var. ?	Brownhead Rush	Grass-like	- - -	- NP	--	
<input type="radio"/> <i>Juncus phaeocephalus</i> var. <i>paniculatus</i>	Panicled Brownhead Rush	Grass-like	- - C	- NP	Wet places, coastal, inland (Jun–Sep)	
<input type="radio"/> <i>Juncus phaeocephalus</i> var. <i>phaeocephalus</i>	Brownheaded Rush	Grass-like	- - B	- NP	Coastal meadows, dune hollows, marsh edges (Jun–Aug)	
<input type="radio"/> <i>Juncus</i> sp.	Rush	Grass-like	- - -	-	--	
<input type="radio"/> <i>Juncus tenuis</i>	Slender Rush	Grass-like	- - -	- NP	Uncommon. Damp places (Jun–Sep)	
<input type="radio"/> <i>Juncus xiphioides</i>	Iris-leaved Rush	Grass-like	- - -	- NP	Wet places (Jul–Oct)	
<input type="radio"/> <i>Luzula comosa</i> var. <i>comosa</i> (JM93: no var.)	Common Woodrush	Grass-like	- - -	- NP	Meadows, open woodland, conifer forest (Jun–Jul)	
Juncaginaceae						Juncaginaceae
<input type="radio"/> <i>Triglochin maritima</i>	Common Arrow-grass	Grass-like	- - C	- NP	Coastal salt marshes, interior saline, brackish, alkaline marshes (Apr–Aug)	
<input type="radio"/> <i>Triglochin scilloides</i> (JM93: <i>Lilaea</i>)	Flowering-quillwort	Grass-like	- - C	- NA	Vernal pools, streams, ponds, lake margins (Mar–Oct)	
<input type="radio"/> <i>Triglochin striata</i>	Three-ribbed Arrow-grass	Grass-like	- - A2	- NP	Uncommon. Brackish to freshwater coastal marshes, springs (May–Sep)	
Lamiaceae						Lamiaceae
<input type="radio"/> <i>Acanthomintha lanceolata</i>	Santa Clara Thornmint	Herbaceous	- 4.2 *A2	- NP	Arid, rocky slopes, openings in scrub or woodland, gen on serpentine (Mar–Jun)	
<input type="radio"/> <i>Clinopodium douglasii</i> (JM93: <i>Satureja</i>)	Yerba Buena	Herbaceous	- - -	- NP	Shady places, chaparral, woodland (Apr–Sep)	

Lamiaceae		Sorted Alphabetically by Family, Scientific Name				Lamiaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
<input type="radio"/> <i>Lamium amplexicaule</i>	Clasping Henbit	Herbaceous	- - -	- ZAB	Disturbed sites, cult or abandoned fields (Apr–Sep)	
<input type="radio"/> <i>Lamium purpureum</i>	Red Henbit	Herbaceous	- - -	- ZA	Uncommon. Disturbed sites, meadows (Apr–Sep)	
<input type="radio"/> <i>Lepechinia calycina</i>	Pitcher Sage	Woody	- - B	- NP	Common. Rocky slopes; chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Lycopus americanus</i>	Cutleaf Bugleweed	Herbaceous	- - A1	- NP	Moist areas, marshes, streambanks (Aug–Sep)	
<input type="radio"/> <i>Lycopus asper</i>	Bugleweed	Herbaceous	- - A1	- NP	Moist areas, marshes, streambanks (Jun–Oct)	
<input type="radio"/> <i>Lycopus sp.</i>	Water Horehound / Bugleweed	Herbaceous	- - -	- NP	- -	
<input type="radio"/> <i>Marrubium vulgare</i>	Common Horehound	Herbaceous	- - -	L ZP	Disturbed sites, gen overgrazed pastures (Mar–Nov)	
<input type="radio"/> <i>Melissa officinalis</i>	Lemon Balm	Herbaceous	- - -	- ZP	Moist sites, meadows, fields (Jun–Sep)	
<input type="radio"/> <i>Mentha arvensis</i>	Field Mint	Herbaceous	- - B	- NP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha piperita</i>	Peppermint	Herbaceous	- - -	- ZP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha pulegium</i>	Pennyroyal	Herbaceous	- - -	M ZP	Moist places, fields (Jul–Oct)	
<input type="radio"/> <i>Mentha sp.</i>	Mint	Herbaceous	- - -	-	- -	
<input type="radio"/> <i>Mentha spicata</i> (JM93: var. <i>spicata</i>)	Spearmint	Herbaceous	- - -	- ZP	Moist places, marshes, lakeshores, fields (Jul–Oct)	
<input type="radio"/> <i>Monardella douglasii</i> (JM93: subsp. <i>douglasii</i>)	Douglas Coyotemint	Herbaceous	- - A2	- NA	Grassland, oak woodland, chaparral, serpentine outcrops (May–Nov)	
<input type="radio"/> <i>Monardella villosa subsp. villosa</i> (JM93: <i>M. villosa subsp. villosa, globosa, M. antonina subsp. antonina</i>)	Coyote-mint	Herbaceous	- - -	- NP	Dry rocky slopes, oak woods, chaparral (May–Aug)	
<input type="radio"/> <i>Pogogyne serpylloides</i>	Thymeleaf Beardstyle	Herbaceous	- - -	- NA	Grassy, brushy areas (Mar–Jun)	
<input type="radio"/> <i>Prunella vulgaris var. ?</i>	European Selfheal	Herbaceous	- - -	- ZP	- -	
<input type="radio"/> <i>Prunella vulgaris var. lanceolata</i>	Narrow-leaved Selfheal	Herbaceous	- - A1	- NP	Moist areas, gen conifer forest, woodland (May–Sep)	
<input type="radio"/> <i>Prunella vulgaris var. vulgaris</i>	European Selfheal	Herbaceous	- - -	- ZP	Moist, disturbed sites (Jun–Sep)	
<input type="radio"/> <i>Salvia columbariae</i>	Chia	Herbaceous	- - B	- NA	Dry, disturbed sites, chaparral, coastal-sage scrub (Mar–Jun)	
<input type="radio"/> <i>Salvia mellifera</i>	Black Sage	Woody	- - -	- NP	Coastal-sage scrub, lower chaparral (Mar–Jun)	
<input type="radio"/> <i>Salvia spathacea</i>	California Hummingbird Sage	Herbaceous	- - A1	- NP	Common, oak woodland, chaparral, coastal-sage scrub, slopes (Mar–May)	
<input type="radio"/> <i>Scutellaria californica</i>	California Skullcap	Herbaceous	- - A2	- NP	Open sites, scrub, woodland (Jun–Jul)	
<input type="radio"/> <i>Scutellaria siphocampyloides</i>	Grayleaf Skullcap	Herbaceous	- - A1	- NP	Open sites, seeps, dry stream beds, scrub, woodland (May–Jul)	
<input type="radio"/> <i>Scutellaria tuberosa</i>	Blue / Tuberous Skullcap	Herbaceous	- - C	- NP	Dry sites, chaparral, oak woodland (Mar–Jul)	
<input type="radio"/> <i>Stachys ajugoides</i> (JM93: var. <i>ajugoides</i>)	Bugle Hedge-nettle	Herbaceous	- - A2	- NA	Moist, open places, gen remain wet into summer (Jun–Sep)	
<input type="radio"/> <i>Stachys albens</i>	White-stem Hedge-nettle	Herbaceous	- - B	- NP	Swamps, seeps (May–Oct)	

Lamiaceae		Sorted Alphabetically by Family, Scientific Name				Lamiaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Stachys bullata</i>	California Hedge-nettle	Herbaceous	- - A1?	-	NP	Dry slopes near coast (Mar–Sep)
<input type="radio"/> <i>Stachys pycnantha</i>	Short-spiked Hedge-nettle	Herbaceous	- - C	-	NP	Streambanks, springs, pine/oak forest (Jun–Oct)
<input type="radio"/> <i>Stachys rigida</i> var. <i>quercetorum</i> (JM93: <i>S. ajugoides</i> var. <i>rigida</i>)	Common Rigid Hedge-nettle	Herbaceous	- - -	-	NP	Moist to ± dry places (Mar–Oct)
<input type="radio"/> <i>Trichostema lanceolatum</i>	Vinegar Weed	Herbaceous	- - -	-	NA	Dry, open, gen disturbed habitats (Jun–Nov)
Lauraceae						Lauraceae
<input type="radio"/> <i>Umbellularia californica</i>	California Bay	Woody	- - -	-	NP	Common. Canyons, valleys, chaparral (Nov–May)
Lemnaceae						Lemnaceae
<input type="radio"/> <i>Lemna gibba</i>	Swollen Duckweed	Herbaceous	- - -	-	NP	Common. Fresh or brackish water -
<input type="radio"/> <i>Lemna minor</i>	Common Lesser Duckweed	Herbaceous	- - -	-	NP	Common. Freshwater (Aug)
<input type="radio"/> <i>Lemna</i> sp.	Duckweed	Herbaceous	- - -	-	NP	- -
Liliaceae						Liliaceae
<input type="radio"/> <i>Asparagus officinalis</i> subsp. <i>officinalis</i>	Garden Asparagus	Herbaceous	- - -	-	ZP	Disturbed places, roadsides, fields (Mar–Sep)
<input type="radio"/> <i>Calochortus albus</i>	White Globe Lily	Herbaceous	- - -	-	NP	Common. Shady to open woodland, scrub (Apr–Jun)
<input type="radio"/> <i>Calochortus argillosus</i>	Clay Mariposa Lily	Herbaceous	- - -	-	NP	Hard clay from volcanic or metamorphic rocks (Apr–Jun)
<input type="radio"/> <i>Calochortus luteus</i>	Yellow Mariposa Lily	Herbaceous	- - C	-	NP	Heavy soils in grassland, woodland, mixed-evergreen forest (Apr–Jun)
<input type="radio"/> <i>Calochortus pulchellus</i>	Mount Diablo Fairy-lantern	Herbaceous	- 1B.2 *A2	-	NP	Wooded slopes, rarely chaparral, gen n aspect (Apr–Jun)
<input type="radio"/> <i>Calochortus</i> sp.	Mariposa Lily	Herbaceous	- - -	-	NP	- -
<input type="radio"/> <i>Calochortus splendens</i>	Splendid Lilac Mariposa Lily	Herbaceous	- - A1	-	NP	Dry, often granitic soils, grassland, chaparral, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calochortus superbus</i>	Superb Mariposa Lily	Herbaceous	- - -	-	NP	Common. Open grassland, woodland, dry meadows, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Calochortus umbellatus</i>	Oakland Star-tulip	Herbaceous	- 4.2 *A2	-	NP	Open chaparral or woodland, gen on serpentine (Mar–May)
<input type="radio"/> <i>Calochortus venustus</i>	White Butterfly Mariposa Lily	Herbaceous	- - B	-	NP	Sandy (often granitic) soil in grassland, woodland, yellow-pine forest (May–Jul)
<input type="radio"/> <i>Clintonia andrewsiana</i>	Red-bead Lily	Herbaceous	- - A1	-	NP	Shaded, damp redwood forest (May–Jul)
<input type="radio"/> <i>Fritillaria affinis</i> (JM93: var. <i>affinis</i>)	Checker Lily	Herbaceous	- - -	-	NP	Common. Oak or pine scrub, grassland (Mar–Jun)
<input type="radio"/> <i>Fritillaria agrestis</i>	Stinkbells	Herbaceous	- 4.2 *A2	-	NP	Clay (gen serpentine) banks, depressions (Mar–Jun)
<input type="radio"/> <i>Fritillaria liliacea</i>	Fragrant Fritillary	Herbaceous	- 1B.2 *A1	-	NP	Heavy soil, open hills, fields near coast (Feb–Apr)
<input type="radio"/> <i>Lilium pardalinum</i> subsp. <i>pardalinum</i>	Leopard Lily	Herbaceous	- - A2	-	NP	Moist places (drier along coast) (May–Aug)

Liliaceae		Sorted Alphabetically by Family, Scientific Name					Liliaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Prosartes hookeri</i> (JM93: <i>Disporum</i>)	Hooker Fairy Bells	Herbaceous	- - C	-	NP	Montane conifer, mixed-evergreen forest, exposed roadside (Mar–Jun)	
<input type="radio"/> <i>Prosartes smithii</i> (JM93: <i>Disporum</i>)	Large-flower Fairy Bells	Herbaceous	- - A1?	-	NP	Moist, shady coastal, montane forest (Mar–Jun)	
Linaceae							Linaceae
<input type="radio"/> <i>Hesperolinon breweri</i>	Brewer's Western Flax	Herbaceous	- 1B.2 *A2	-	NA	Chaparral or grassland, occ on serpentine (May–Jun)	
<input type="radio"/> <i>Hesperolinon californicum</i>	California Flax	Herbaceous	- - A2	-	NA	Rocky areas, chaparral, grassland, occasionally on serpentine (Apr–Jul)	
<input type="radio"/> <i>Hesperolinon sp.</i>	Flax	Herbaceous	- - -	-	NA		
<input type="radio"/> <i>Linum bienne</i>	Narrow-leaf Flax	Herbaceous	- - -	-	ZP	Garden escape; grassland, woodland, disturbed places, esp coastal (Mar–Jun)	
<input type="radio"/> <i>Linum sp.</i>	Flax	Herbaceous	- - -	-	- -		
Loasaceae							Loasaceae
<input type="radio"/> <i>Mentzelia affinis</i>	Hydra Blazing Star	Herbaceous	- - A2	-	NA	Rocky or gray-white soils in grassland, woodland, creosote-bush scrub (Apr–May)	
<input type="radio"/> <i>Mentzelia dispersa</i>	Nevada Blazing Star	Herbaceous	- - A2	-	NA	Loamy to sandy or rocky slopes, roadsides (May–Aug)	
<input type="radio"/> <i>Mentzelia lindleyi</i>	Lindley Blazing Star	Herbaceous	- - A2	-	NA	Rocky, open slopes, coastal-sage scrub, oak/pine woodland (May–Jun)	
Lythraceae							Lythraceae
<input type="radio"/> <i>Lythrum californicum</i>	California Loosestrife	Herbaceous	- - B	-	NP	Marshes, pond and stream margins (Apr–Sep)	
<input type="radio"/> <i>Lythrum hyssopifolia</i>	Grass-poly	Herbaceous	- - -	-	ZAP	Marshes, drying pond margins, disturbed ground (Apr–Oct)	
<input type="radio"/> <i>Lythrum sp.</i>	Loosestrife	Herbaceous	- - -	-	- -		
<input type="radio"/> <i>Lythrum tribracteatum</i>	Three-bract Loosestrife	Herbaceous	- - -	-	ZA	Wet areas, drying ponds (May–Jun)	
Malvaceae							Malvaceae
<input type="radio"/> <i>Eremalche parryi</i> subsp. <i>parryi</i>	Parry's Mallow	Herbaceous	- - A1x	-	NA	Grassland, scrub, foothill woodland (Mar–May)	
<input type="radio"/> <i>Malacothamnus fasciculatus</i> var. <i>nuttallii</i> (JM93: no var.)	Chaparral Mallow	Woody	- - -	-	NP	Open chaparral (May–Jul)	
<input type="radio"/> <i>Malacothamnus fremontii</i>	Fremont Bush-mallow	Woody	- - B	-	NP	Open chaparral to pine woodland (May–Jul)	
<input type="radio"/> <i>Malva nicaeensis</i>	Bull Mallow	Herbaceous	- - -	-	ZA	Disturbed places (Mar–Jun)	
<input type="radio"/> <i>Malva parviflora</i>	Cheeseweed	Herbaceous	- - -	-	ZA	Common. Disturbed places (Mar–May)	
<input type="radio"/> <i>Malva pseudolavatera</i> (JM93: <i>Lavatera cretica</i>)	Cretan Mallow	Herbaceous	- - -	-	ZAB	Uncommon. Disturbed places on coastal bluffs, dunes, occ inland (Apr–Jun)	
<input type="radio"/> <i>Malva sp.</i>	Mallow	Herbaceous	- - -	-	- -		

Malvaceae		Sorted Alphabetically by Family, Scientific Name					Malvaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Malvella leprosa</i>	Alkal-mallow	Herbaceous	- - -	-	NP	Valleys, gen saline (Apr–Nov)	
<input type="radio"/> <i>Sidalcea diploscypha</i>	Fringed Checkerbloom	Herbaceous	- - A2	-	NA	Grassland, open woodland, valleys, occasionally near vernal pools, generally serpentine (Apr–May)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. ?	Common Checkerbloom	Herbaceous	- - -	-	NP	--	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>laciniata</i>	Geranium-leaved Checkerbloom	Herbaceous	- - C	-	NP	Grassland, open woodland (Mar–Jun)	
<input type="radio"/> <i>Sidalcea malviflora</i> subsp. <i>malviflora</i>	Common Checkerbloom	Herbaceous	- - B	-	NP	Coastal prairie, scrub, open forest (Mar–Jul)	
Melanthiaceae							Melanthiaceae
<input type="radio"/> <i>Toxicoscordion fremontii</i> (JM93: <i>Zigadenus</i>)	Common Star Lily	Herbaceous	- - -	-	NP	Grassy or wooded slopes, outcrops (Feb–Jun)	
<input type="radio"/> <i>Trillium chloropetalum</i>	Giant Trillium	Herbaceous	- - -	-	NP	Edges of redwood forest, chaparral, gen moist slopes, canyon banks in alluvial soils (Apr–May)	
<input type="radio"/> <i>Trillium ovatum</i> subsp. <i>ovatum</i>	Western White Trillium	Herbaceous	- - A2	-	NP	Redwood, mixed-evergreen forest on moist wooded slopes (Feb–Apr)	
Molluginaceae							Molluginaceae
<input type="radio"/> <i>Glinus lotoides</i>	Damascisa	Herbaceous	- - -	-	ZA	Uncommon. Moist or seasonally dry margins of wetlands (Jun–Nov)	
Montiaceae							Montiaceae
<input type="radio"/> <i>Calandrinia breweri</i>	Brewer Red Maids	Herbaceous	- 4.2 *A2	-	NA	Sandy to loamy soil, disturbed sites, burns (Feb–May)	
<input type="radio"/> <i>Calandrinia ciliata</i>	Red Maids	Herbaceous	- - -	-	NA	Common. Sandy to loamy soil, grassy areas, cult fields (Feb–May)	
<input type="radio"/> <i>Claytonia exigua</i> subsp. <i>exigua</i>	Common Pale Claytonia	Herbaceous	- - C	-	NA	Dry or moist, disturbed bare clay to sandy soils, often serpentine (Apr–Jul)	
<input type="radio"/> <i>Claytonia gypsophiloides</i>	Santa Lucia Claytonia	Herbaceous	- - A2	-	NA	Moist, bare, often stony sites, in sun or shade, often serpentine (Mar–May)	
<input type="radio"/> <i>Claytonia parviflora</i> subsp. ?	Miner's Lettuce	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Claytonia parviflora</i> subsp. <i>utahensis</i> (JM93: subsp. <i>parviflora</i>)	Small-flower Claytonia	Herbaceous	- - -	-	NA	Rock crevices, alluvial fans, boulder fields (Apr–Jul)	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>mexicana</i>	Angle-leaf Miner's Lettuce	Herbaceous	- - -	-	NA	Shrubland, woodland, rock crevices, rockslides (Feb–Apr)	
<input type="radio"/> <i>Claytonia perfoliata</i> subsp. <i>perfoliata</i>	Common Miner's Lettuce	Herbaceous	- - -	-	NA	Vernally moist, often shady or disturbed sites (Jan–May)	
<input type="radio"/> <i>Lewisia rediviva</i> var. <i>rediviva</i> (JM93: no var.)	Bitter Root	Herbaceous	- - A2	-	NP	Rocky, sandy ground, open conifer woodland, scrub (Mar–Jun)	
<input type="radio"/> <i>Montia fontana</i>	Water Chickweed	Herbaceous	- - C	-	NA	Common. Ponds, streams, vernal pools, seeps (Jul–Aug)	

Myricaceae	Sorted Alphabetically by Family, Scientific Name					Myricaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
Myricaceae						Myricaceae
<input type="radio"/> <i>Morella californica</i> (JM93: <i>Myrica</i>)	California Bayberry	Woody	- - A2	- NP	Coastal dunes and scrub, closed-cone pine, redwood forest (Mar–Apr)	
Myrsinaceae						Myrsinaceae
<input type="radio"/> <i>Anagallis arvensis</i>	Scarlet Pimpernel	Herbaceous	- - -	- ZA	Common. Disturbed places, ocean beaches (Mar–May)	
<input type="radio"/> <i>Glaux maritima</i>	Sea Milkwort	Herbaceous	- - A1	- NP	Coastal salt marshes, saline meadows (May–Jul)	
<input type="radio"/> <i>Trientalis latifolia</i>	Starflower	Herbaceous	- - C	- NP	Shaded places, esp woodland (Apr–Jul)	
Myrtaceae						Myrtaceae
<input type="radio"/> <i>Eucalyptus camaldulensis</i>	River Red Gum	Woody	- - -	L ZP	Common. Disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Eucalyptus globulus</i>	Blue Gum	Woody	- - -	M ZP	Common. Disturbed areas (Oct–Jan)	
<input type="radio"/> <i>Eucalyptus sp.</i>	Gum	Woody	- - -	- ZP	- -	
<input type="radio"/> <i>Eucalyptus tereticornis</i>	Forest Red Gum	Woody	- - -	- ZP	Uncommon. Disturbed coastal areas (May–Aug)	
<input type="radio"/> <i>Eucalyptus viminalis</i>	Manna Gum	Woody	- - -	- ZP	Uncommon. Disturbed areas (Jul–Sep)	
Nyctaginaceae						Nyctaginaceae
<input type="radio"/> <i>Mirabilis laevis</i> var. <i>crassifolia</i> (JM93: <i>M. californica</i>)	Wishbone Bush	Herbaceous	- - A1	- NP	Grassy areas, chaparral, dunes, dry rocky areas, washes Dec–Jun)	
Nymphaeaceae						Nymphaeaceae
<input type="radio"/> <i>Nymphaea mexicana</i>	Yellow Or Banana Waterlily	Herbaceous	- - -	- ZP	Lakes, ponds, slow streams (Apr–Jul)	
<input type="radio"/> <i>Nymphaea odorata</i>	Fragrant Or White Waterlily	Herbaceous	- - -	- ZP	Quiet waters, ponds, edges of lakes (Apr–Aug)	
Oleaceae						Oleaceae
<input type="radio"/> <i>Forestiera pubescens</i>	Desert Olive	Woody	- - A2	- NP	Streambanks, canyons, washes (Mar–Apr)	
<input type="radio"/> <i>Fraxinus dipetala</i>	California Ash	Woody	- - A2	- NP	Canyons, slopes, chaparral, oak/pine woodland (Apr–Jun)	
<input type="radio"/> <i>Fraxinus latifolia</i>	Oregon Ash	Woody	- - B	- NP	Canyons, streambanks, woodland (Mar–May)	
<input type="radio"/> <i>Fraxinus sp.</i>	Ash	Woody	- - -	- NP	- -	
<input type="radio"/> <i>Olea europaea</i>	Olive	Woody	- - -	L ZP	Generally waif, persisting from cultivation (Feb–Jun)	
Onagraceae						Onagraceae
<input type="radio"/> <i>Camissonia contorta</i>	Contorted Suncup	Herbaceous	- - B	- NA	Sandy soil, slopes, flats, often disturbed, grassland, chaparral, pinyon/juniper woodland (May–Ju)	

Onagraceae		Sorted Alphabetically by Family, Scientific Name				Onagraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Camissoniopsis intermedia</i> (JM93: <i>Camissonia</i>)	Intermediate Suncup	Herbaceous	- - A2	-	NA	Shrubby slopes, especially burns (Mar–May)
<input type="radio"/> <i>Camissoniopsis micrantha</i> (JM93: <i>Camissonia</i>)	Small-flower Suncup	Herbaceous	- - B	-	NA	Coastal dunes, beaches, sandy fields, washes (Mar–May)
<input type="radio"/> <i>Clarkia affinis</i>	Small Clarkia	Herbaceous	- - C	-	NA	Openings in woodland, chaparral (May–Jun)
<input type="radio"/> <i>Clarkia biloba</i> subsp. <i>biloba</i>	Mule-ear Clarkia	Herbaceous	- - A2	-	NA	Foothill woodland, serpentine or not (May–Aug)
<input type="radio"/> <i>Clarkia concinna</i> subsp. ?	Red Ribbons	Herbaceous	- - -	-	NA	- -
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>automixa</i>	Santa Clara Red Ribbons	Herbaceous	- 4.3 *A1	-	NA	Woodland (Apr–Jun)
<input type="radio"/> <i>Clarkia concinna</i> subsp. <i>concinna</i>	Red Ribbons	Herbaceous	- - C	-	NA	Mixed-evergreen forest, woodland, coastal scrub (Apr–Jul)
<input type="radio"/> <i>Clarkia franciscana</i>	Presidio Clarkia	Herbaceous	FE/CE 1B.1 *A1	-	NA	Serpentine soil (May–Jun)
<input type="radio"/> <i>Clarkia gracilis</i> subsp. <i>gracilis</i>	Herald-of-summer	Herbaceous	- - B	-	NA	Common. Openings in woodland, forest (Apr–Jul)
<input type="radio"/> <i>Clarkia modesta</i>	Waltham Creek Fairy Fan	Herbaceous	- - A2	-	NA	Shady places in woodland (Apr–May)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. ?	Winecup Clarkia	Herbaceous	- - -	-	NA	- -
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>purpurea</i>	Purple Winecup Clarkia	Herbaceous	- - A2	-	NA	Uncommon. Grassland (Apr–Jun)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>quadrivulnera</i>	Four-spot	Herbaceous	- - -	-	NA	Common. Open, grassy or shrubby places (Apr–Aug)
<input type="radio"/> <i>Clarkia purpurea</i> subsp. <i>viminea</i>	Large Clarkia	Herbaceous	- - A2	-	NA	Open, grassy or shrubby places (May–Jul)
<input type="radio"/> <i>Clarkia rubicunda</i>	Ruby Chalice Clarkia	Herbaceous	- - -	-	NA	Openings in woodland, forest, chaparral near coast (May–Aug)
<input type="radio"/> <i>Clarkia</i> sp.	Clarkia	Herbaceous	- - -	-	NA	- -
<input type="radio"/> <i>Clarkia unguiculata</i>	Elegant Clarkia	Herbaceous	- - -	-	NA	Common. Woodland (Apr–Sep)
<input type="radio"/> <i>Epilobium brachycarpum</i>	Panicled / Weedy Willowherb	Herbaceous	- - -	-	NA	Common. Dry open or disturbed woodland, grassland, roadsides (Jun–Sep)
<input type="radio"/> <i>Epilobium campestre</i> (JM93: <i>E. pygmaeum</i>)	Smooth Willowherb	Herbaceous	- - B	-	NA	Vernal pools, clay mud flats (May–Sep)
<input type="radio"/> <i>Epilobium canum</i> subsp. <i>canum</i>	California Fuchsia	Herbaceous	- - -	-	NP	Dry slopes, ridges (Jun–Dec)
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>ciliatum</i>	Common Willowherb	Herbaceous	- - -	-	NP	Common. Disturbed places, moist meadows, streambanks, roadsides (Jun–Oct)
<input type="radio"/> <i>Epilobium ciliatum</i> subsp. <i>watsonii</i>	Watson Willowherb	Herbaceous	- - B	-	NP	Moist coastal bluffs, streamsides, ± disturbed sites (May–Oct)
<input type="radio"/> <i>Epilobium densiflorum</i>	Dense-flower Willowherb	Herbaceous	- - C	-	NA	Streambanks, outwashes, seasonal moist flats (May–Oct)
<input type="radio"/> <i>Epilobium minutum</i>	Chaparral Willowherb	Herbaceous	- - -	-	NA	Dry, open, disturbed areas, vernal pools, often after fire (Apr–Sep)
<input type="radio"/> <i>Epilobium</i> sp.	Willowherb	Herbaceous	- - -	-	NA	- -
<input type="radio"/> <i>Epilobium torreyi</i>	Narrowleaf Willowherb	Herbaceous	- - A2	-	NA	Seasonally moist streambanks, seeps, roadside ditches (May–Aug)

Onagraceae		Sorted Alphabetically by Family, Scientific Name					Onagraceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Eremothera boothii</i> subsp. <i>decorticans</i> (JM93: <i>Camissonia</i>)	Shredding Evening-primrose	Herbaceous	- - A1	-	NA	Open, generally steep and rocky, especially shale slopes (Mar–Jun)	
<input type="radio"/> <i>Ludwigia hexapetala</i>	Uruguayan Primrose-willow	Herbaceous	- - A1?	H	NP	Lake margins, wetlands (May–Dec)	
<input type="radio"/> <i>Ludwigia peploides</i> subsp. <i>peploides</i>	Yellow Water Primrose	Herbaceous	- - B	-	NP	Lakeshores, streambanks, seasonal wetlands (May–Oct)	
<input type="radio"/> <i>Ludwigia</i> sp.	Water Primrose	Herbaceous	- - -	-	NP	- -	
<input type="radio"/> <i>Oenothera deltooides</i> subsp. <i>howellii</i>	Antioch Dunes Evening-primrose	Herbaceous	FE/CE 1B.1 *A2	-	NP	Sandy bluffs, dunes (Mar–Sep)	
<input type="radio"/> <i>Oenothera elata</i> subsp. <i>hirsutissima</i>	Hairy Evening-primrose	Herbaceous	- - -	-	NBP	Moist places, generally inland (Jun–Sep)	
<input type="radio"/> <i>Oenothera elata</i> subsp. <i>hookeri</i>	Hooker Evening-primrose	Herbaceous	- - B	-	NB	Moist, coastal, slightly inland, sandy bluffs (Jun–Sep)	
<input type="radio"/> <i>Taraxia ovata</i> (JM93: <i>Camissonia</i>)	Golden Eggs Suncup	Herbaceous	- - C	-	NP	Grassy fields, gen clay soil (Mar–Jun)	
<input type="radio"/> <i>Tetrapteron graciliflorum</i> (JM93: <i>Camissonia graciliflora</i>)	Hill Sun Cup	Herbaceous	- - A2	-	NA	Open or shrubby slopes, generally clay soils, grassland, oak and Joshua-tree woodland (Mar–Apr)	
Orchidaceae							Orchidaceae
<input type="radio"/> <i>Corallorhiza maculata</i> var. <i>maculata</i> (JM93: <i>no var.</i>)	Spotted Coralroot	Herbaceous	- - A2	-	NP	Shaded mixed-evergreen or conifer forest, in decomposing lf litter (May–Aug)	
<input type="radio"/> <i>Corallorhiza striata</i>	Striped Coralroot	Herbaceous	- - A1	-	NP	Open to shaded mixed-evergreen or conifer forest, in decomposing leaf litter (Feb–Jul)	
<input type="radio"/> <i>Epipactis gigantea</i>	Stream / Brook Orchid	Herbaceous	- - A1	-	NP	Seeps, wet meadows, streambank (Mar–Oct)	
<input type="radio"/> <i>Epipactis helleborine</i>	Broad-leaved Helleborine Orchid	Herbaceous	- - -	-	ZP	Generally dry slopes, roadcuts, mixed-conifer forest (Apr–Dec)	
<input type="radio"/> <i>Piperia elegans</i> subsp. <i>elegans</i> (JM93: <i>no subsp.</i>)	Elegant Rein Orchid	Herbaceous	- - B	-	NP	Generally dry, open sites, scrub, conifer forest (May–Sep)	
<input type="radio"/> <i>Piperia elongata</i>	Wood Rein-orchid	Herbaceous	- - A2	-	NP	Generally dry sites, scrub, chaparral, mixed-evergreen or conifer forest (May–Jul)	
<input type="radio"/> <i>Piperia</i> sp.	Rein Orchid	Herbaceous	- - -	-	NP	- -	
<input type="radio"/> <i>Piperia unalascensis</i>	Alaska Rein Orchid	Herbaceous	- - A1	-	NP	Generally dry sites, scrub, woodland, forest (May–Aug)	
<input type="radio"/> <i>Spiranthes porrifolia</i>	Western Ladies Tresses	Herbaceous	- - A1	-	NP	Wet meadows, freshwater marshes, seeps, grassland, oak woodland (Jun–Sep)	
Orobanchaceae							Orobanchaceae
<input type="radio"/> <i>Bellardia trixago</i>	Mediterranean Linseed	Herbaceous	- - -	L	ZA	Disturbed grassland. (Apr–Jun)	
<input type="radio"/> <i>Castilleja affinis</i> subsp. <i>affinis</i>	Common Indian Paintbrush	Herbaceous	- - -	-	NP	Chaparral, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Castilleja attenuata</i>	Valley Tassels	Herbaceous	- - -	-	NA	Grassland (Mar–May)	

Orobanchaceae		Sorted Alphabetically by Family, Scientific Name				Orobanchaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Castilleja densiflora</i> subsp. <i>densiflora</i>	Common Owl's-clover	Herbaceous	- - -	-	NA	Grassland (Mar–May)	
<input type="radio"/> <i>Castilleja exserta</i> subsp. <i>exserta</i>	Purple Owl's-clover	Herbaceous	- - -	-	NA	Open fields, grassland (Mar–May)	
<input type="radio"/> <i>Castilleja foliolosa</i>	Woolly Paintbrush	Herbaceous	- - -	-	NP	Dry, open, rocky slopes, edges of chaparral (Mar–Jun)	
<input type="radio"/> <i>Castilleja rubicundula</i> subsp. <i>lithospermoides</i>	Yellow Cream Sacs	Herbaceous	- - B	-	NA	Open grassland (Apr–Jun)	
<input type="radio"/> <i>Castilleja subinclusa</i> subsp. <i>franciscana</i>	Franciscan Paintbrush	Herbaceous	- - A2	-	NP	Coastal scrub (Mar–Jul)	
<input type="radio"/> <i>Castilleja wightii</i>	Seaside / Coastal Paintbrush	Herbaceous	- - C	-	NP	Coastal scrub (Mar–Aug)	
<input type="radio"/> <i>Chloropyron molle</i> subsp. <i>molle</i> (JM93: <i>Cordylanthus mollis</i> subsp. <i>mollis</i>)	Soft Salty Bird's-beak	Herbaceous	FE/CR 1B.2 *A2	-	NA	Coastal salt marshes (Jul–Nov)	
<input type="radio"/> <i>Cordylanthus pilosus</i> subsp. <i>pilosus</i>	Hairy Bird's Beak	Herbaceous	- - B	-	NA	Open foothill woodland, chaparral, on serpentine (Jul–Sep)	
<input type="radio"/> <i>Cordylanthus rigidus</i> subsp. <i>rigidus</i>	Stiff Bird's Beak	Herbaceous	- - A1	-	NA	Open foothill woodland, chaparral margins, conifer forest (Jul–Sep)	
<input type="radio"/> <i>Kopsiopsis strobilacea</i> (JM93: <i>Boschniakia</i>)	Cone Flower / Ground Cone	Herbaceous	- - A1	-	NP	Open woodland, chaparral, generally on <i>Arctostaphylos</i> (Apr–Jun)	
<input type="radio"/> <i>Orobanche bulbosa</i>	Chaparral Broom-rape	Herbaceous	- - A2	-	NP	Openings in chaparral, generally on <i>Adenostoma fasciculatum</i> (Apr–Jun)	
<input type="radio"/> <i>Orobanche californica</i> subsp. <i>jepsonii</i>	Jepson Broomrape	Herbaceous	- - A1	-	NP	Uncommon. Gen dry flats, slopes. Root parasite on sunflower family (Jul–Sep)	
<input type="radio"/> <i>Orobanche fasciculata</i>	Clustered Broomrape	Herbaceous	- - B	-	NP	Dry, gen bare places. Root parasite on various shrubs (Apr–Jul)	
<input type="radio"/> <i>Orobanche uniflora</i>	Naked Broom-rape	Herbaceous	- - C	-	NP	Generally moist places, on herbs, especially <i>Sedum</i> , <i>Saxifragaceae</i> , <i>Asteraceae</i> (Apr–Jul)	
<input type="radio"/> <i>Orobanche vallicola</i>	Sharp-lobe Broom-rape	Herbaceous	- - A2	-	NP	Riparian woodland, on <i>Sambucus nigra</i> (Jul–Sep)	
<input type="radio"/> <i>Pedicularis densiflora</i>	Warrior's Plume	Herbaceous	- - -	-	NP	Dry chaparral, oak/pine forest (Mar–May)	
<input type="radio"/> <i>Triphysaria eriantha</i> subsp. <i>eriantha</i>	Yellow Johnny-tuck	Herbaceous	- - -	-	NA	Grassland, foothills (Mar–May)	
<input type="radio"/> <i>Triphysaria pusilla</i>	Dwarf Owl's Clover	Herbaceous	- - -	-	NA	Grassland (Apr–Jun)	
<input type="radio"/> <i>Triphysaria versicolor</i> subsp. <i>faucibarbata</i>	Smooth Owl's Clover	Herbaceous	- - A2	-	NA	Grassland (Apr–Jun)	
Oxalidaceae						Oxalidaceae	
<input type="radio"/> <i>Oxalis corniculata</i>	Creeping Wood Sorrel	Herbaceous	- - -	-	ZP	Disturbed areas (Most of Year)	
<input type="radio"/> <i>Oxalis pes-caprae</i>	Bermuda Buttercup	Herbaceous	- - -	M	ZP	Disturbed areas, roadsides, grassland, dunes (Jan–May)	
<input type="radio"/> <i>Oxalis pilosa</i> (JM93: <i>O. albicans</i> subsp. <i>pilosa</i>)	Hairy Wood Sorrel	Herbaceous	- - A2	-	NP	Coastal grassland, scrub, chaparral (Feb–Sep)	
<input type="radio"/> <i>Oxalis</i> sp.	Wood Sorrel	Herbaceous	- - -	-	- -		
Papaveraceae						Papaveraceae	
<input type="radio"/> <i>Dendromecon rigida</i>	Bush Poppy	Woody	- - A2	-	NP	Dry slopes, washes, esp recent burns (Apr–Jun)	

Papaveraceae		Sorted Alphabetically by Family, Scientific Name				Papaveraceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Dicentra formosa</i>	Pacific Bleeding Heart	Herbaceous	- - A2	-	NP	Damp, shaded areas (Mar–Jul)	
<input type="radio"/> <i>Ehrendorferia chrysantha</i> (JM93: <i>Dicentra</i>)	Golden Eardrops	Herbaceous	- - A2	-	NP	Dry slopes, burns, disturbed areas (Apr–Sep)	
<input type="radio"/> <i>Eschscholzia californica</i>	California Poppy	Herbaceous	- - -	-	NP	Grassy, open areas (Feb–Sep)	
<input type="radio"/> <i>Fumaria officinalis</i>	Fumitory	Herbaceous	- - -	-	ZA	Disturbed places (Apr–Jul)	
<input type="radio"/> <i>Meconella californica</i>	California Meconella	Herbaceous	- - A1	-	NA	Open, rocky areas (Mar–May)	
<input type="radio"/> <i>Papaver californicum</i>	Fire Poppy	Herbaceous	- - A2	-	NA	Burns, disturbed areas, open woodland (Apr–May)	
<input type="radio"/> <i>Papaver heterophyllum</i> (JM93: <i>Stylomecon heterophylla</i>)	Wind Poppy	Herbaceous	- - -	-	NA	Grassy areas, openings in chaparral (Apr–May)	
<input type="radio"/> <i>Platystemon californicus</i>	Cream Cups	Herbaceous	- - -	-	NA	Open grassland, sandy soil, burns (Mar–May)	
Phrymaceae						Phrymaceae	
<input type="radio"/> <i>Mimulus aurantiacus</i> var. <i>aurantiacus</i> (JM93: no var.)	Bush Monkeyflower	Woody	- - -	-	NP	Disturbed areas, coastal cliffs, canyon sides (Mar–Jun)	
<input type="radio"/> <i>Mimulus cardinalis</i>	Scarlet Monkeyflower	Herbaceous	- - C	-	NP	Moist to wet places along streams, seepage areas (May–Sep)	
<input type="radio"/> <i>Mimulus douglasii</i>	Mickey Mouse Monkeyflower	Herbaceous	- - A2	-	NA	Bare clay, serpentine or granitic soils; generally along upper banks of small creeks (Feb–Apr)	
<input type="radio"/> <i>Mimulus guttatus</i>	Golden Monkeyflower	Herbaceous	- - -	-	NP	Common. Wet places, gen terrestrial, occ emergent or floating in mats (Mar–Aug)	
<input type="radio"/> <i>Mimulus latidens</i>	Broadtooth Monkeyflower	Herbaceous	- - A2	-	NA	Vernally wet depressions (Apr–Jun)	
<input type="radio"/> <i>Mimulus pilosus</i>	Downy Monkeyflower	Herbaceous	- - A2	-	NA	Moist, sandy areas, especially by small streams, disturbed areas (Apr–Aug)	
Pinaceae						Pinaceae	
<input type="radio"/> <i>Pinus attenuata</i>	Knobcone Pine	Woody	- - A2	-	NP	Closed-cone-pine forest, chaparral -	
<input type="radio"/> <i>Pinus coulteri</i>	Coulter Pine	Woody	- - A2	-	NP	Chaparral, lower mixed-conifer, mixed-hardwood forests -	
<input type="radio"/> <i>Pinus radiata</i>	Monterey Pine	Woody	- - -	-	NP	Closed-cone-pine forest, oak woodland -	
<input type="radio"/> <i>Pinus sabiniana</i>	Gray Pine	Woody	- - -	-	NP	Foothill woodland, n oak woodland, chaparral, infertile soils in mixed-conifer and hardwood forests -	
Plantaginaceae						Plantaginaceae	
<input type="radio"/> <i>Antirrhinum multiflorum</i>	Mayflower Snapdragon	Herbaceous	- - A1	-	NA	Rocky or disturbed areas, burns (Apr–Aug)	
<input type="radio"/> <i>Antirrhinum vexillocalyculatum</i> subsp. <i>vexillocalyculatum</i> (JM93: <i>A. vexillocalyculatum</i> subsp. <i>vexillo-calyculatum</i>)	Wiry Snapdragon	Herbaceous	- - B	-	NA	Disturbed areas (Jun–Aug)	
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. ?	White Chinese Houses	Herbaceous	- - -	-	NA	- -	

Plantaginaceae	Sorted Alphabetically by Family, Scientific Name					Plantaginaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. <i>bartsiiifolia</i>	White Chinese Houses	Herbaceous	- - A1	-	NA	Open, sandy places (Mar–Jun)
<input type="radio"/> <i>Collinsia bartsiiifolia</i> var. <i>stricta</i> (JM93: var. <i> davidsonii</i>)	Small White Chinese Houses	Herbaceous	- - A1	-	NA	Chaparral, open oak woodland, dry mixed woodland (May–Jul)
<input type="radio"/> <i>Collinsia heterophylla</i> var. <i>heterophylla</i> (JM93: no var.)	Chinese-houses	Herbaceous	- - -	-	NA	Shady places in chaparral, open mixed woodland, oak woodland (Mar–Jun)
<input type="radio"/> <i>Collinsia parviflora</i>	Blue-eyed Mary	Herbaceous	- - A2	-	NA	Common. Moist, ± shady places, montane (Mar–Jul)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. ?	Few-flowered Collinsia	Herbaceous	- - -	-	NA	--
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>collina</i>	Few-flowered Collinsia	Herbaceous	- - C	-	NA	Disturbed grassy fields, roadbanks, open chaparral, open oak and dry mixed woodland (Mar–Apr)
<input type="radio"/> <i>Collinsia sparsiflora</i> var. <i>sparsiflora</i>	Few-flowered Collinsia	Herbaceous	- - C	-	NA	Grassy, sometimes disturbed or rocky places, chaparral, oak woodland, dry mixed woodland (Mar–May)
<input type="radio"/> <i>Cymbalaria muralis</i>	Kenilworth Ivy	Herbaceous	- - -	-	ZP	Rock walls, shady, disturbed areas (May–Sep)
<input type="radio"/> <i>Digitalis purpurea</i>	Foxglove	Herbaceous	- - -	L	ZB	Acid soil in open woodland, disturbed areas (May–Jul)
<input type="radio"/> <i>Keckiella corymbosa</i>	Redwood / Red Rock Penstemon	Woody	- - A1	-	NP	Rocky slopes in conifer or hardwood forests, (chaparral) (Jun–Oct)
<input type="radio"/> <i>Kickxia elatine</i>	Sharp Point Fluvellin	Herbaceous	- - -	-	ZA	Disturbed, open places (Apr–Oct)
<input type="radio"/> <i>Nuttallanthus texanus</i> (JM93: <i>Linaria canadensis</i>)	Blue Toadflax	Herbaceous	- - B	-	NA	Sand or gravel (Mar–May)
<input type="radio"/> <i>Penstemon centranthifolius</i>	Scarlet Bugler	Herbaceous	- - A2	-	NP	Dry, open chaparral or oak woodland (Apr–Jul)
<input type="radio"/> <i>Penstemon heterophyllus</i> var. <i>heterophyllus</i>	Foothill Penstemon	Herbaceous	- - B	-	NP	Grassland, chaparral, forest openings (Apr–Jul)
<input type="radio"/> <i>Plantago coronopus</i>	Cut-leaf Plantain	Herbaceous	- - -	-	NA	Coastal bluffs, salt marshes, trampled places, chaparral, grassy flats (Apr–Jul)
<input type="radio"/> <i>Plantago erecta</i>	California Dwarf Plantain	Herbaceous	- - -	-	NA	Sandy, clay, serpentine soil; grassy slopes, flats, open woodland (Mar–May)
<input type="radio"/> <i>Plantago lanceolata</i>	English Plantain	Herbaceous	- - -	L	ZA	Common. Disturbed areas (Apr–Aug)
<input type="radio"/> <i>Plantago major</i>	Common Plantain	Herbaceous	- - -	-	ZAP	Disturbed areas (Apr–Sep)
<input type="radio"/> <i>Plantago subnuda</i>	Coast Plantain	Herbaceous	- - A1	-	NP	Wet meadows, ditches, coastal bluffs, marshes (May–Sep)
<input type="radio"/> <i>Tonella tenella</i>	Small-flowered Tonella	Herbaceous	- - A2	-	NA	Moist, shaded places in chaparral, oak and mixed woodland (Mar–Jun)
<input type="radio"/> <i>Veronica americana</i>	American Brooklime	Herbaceous	- - C	-	NP	Common. Moist to wet soil, springs, slow streams, meadows, lakeshores (May–Aug)
<input type="radio"/> <i>Veronica anagallis-aquatica</i>	Water Speedwell	Herbaceous	- - -	-	ZP	Wet meadows, streambanks, slow streams (May–Sep)
<input type="radio"/> <i>Veronica arvensis</i>	Common Speedwell	Herbaceous	- - -	-	ZA	Meadows (Apr–Jul)

Plantaginaceae		Sorted Alphabetically by Family, Scientific Name				Plantaginaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Veronica catenata</i>	Chain Speedwell	Herbaceous	- - -	-	ZP	Wet meadows, slow streams (Jul-Sep)	
<input type="radio"/> <i>Veronica peregrina subsp. xalapensis</i>	Purslane Speedwell	Herbaceous	- - -	-	NA	Moist places (Apr-Aug)	
<input type="radio"/> <i>Veronica persica</i>	Persian Speedwell	Herbaceous	- - -	-	ZA	Wet, disturbed areas, fields (Feb-May)	
<input type="radio"/> <i>Veronica sp.</i>	Speedwell	Herbaceous	- - -	-	--		
Platanaceae						Platanaceae	
<input type="radio"/> <i>Platanus racemosa</i>	Western Sycamore	Woody	- - B	-	NP	Common. Streamsides, canyons, arroyos (Feb-Apr)	
Plumbaginaceae						Plumbaginaceae	
<input type="radio"/> <i>Limonium californicum</i>	Western Marsh-rosemary	Herbaceous	- - B	-	NP	Common; coastal dunes, salt marshes (Jul-Dec)	
Poaceae						Poaceae	
<input type="radio"/> <i>Agrostis avenacea</i>	Pacific Bent Grass	Grass-like	- - -	L	ZP	Open, often disturbed place (Jun-Jul)	
<input type="radio"/> <i>Agrostis capillaris</i>	Colonial Bent Grass	Grass-like	- - -	-	ZP	Roadsides, open, disturbed places (Jul-Sep)	
<input type="radio"/> <i>Agrostis exarata</i>	Spike Bent Grass	Grass-like	- - -	-	NP	Common. Moist or disturbed areas, open woodland, conifer forest (Jun-Aug)	
<input type="radio"/> <i>Agrostis hallii</i>	Hall's Bent Grass	Grass-like	- - A2	-	NP	Open oak woodland, conifer forest (May-Jul)	
<input type="radio"/> <i>Agrostis microphylla</i>	Small-leaf Bent Grass	Grass-like	- - -	-	NA	Thin, rocky soils, cliffs, vernal pools, occasionally on serpentine (May-Jul)	
<input type="radio"/> <i>Agrostis pallens</i>	Dune Bent Grass	Grass-like	- - -	-	NP	Common. Open meadows, woodland, forest, subalpine (Jun-Aug)	
<input type="radio"/> <i>Agrostis sp.</i>	Bent Grass	Grass-like	- - -	-	--		
<input type="radio"/> <i>Agrostis stolonifera</i>	Creeping Bent Grass	Grass-like	- - -	L	ZP	Ditches, lake margins, marshes (Jun-Sep)	
<input type="radio"/> <i>Aira caryophyllea</i>	Silver Hair Grass	Grass-like	- - -	-	ZA	Sandy soils, open or disturbed sites (Apr-Jun)	
<input type="radio"/> <i>Alopecurus saccatus</i>	Pacific Foxtail	Grass-like	- - B	-	NA	Vernal pools, moist, open meadows (Mar-May)	
<input type="radio"/> <i>Arundo donax</i>	Giant Reed	Grass-like	- - -	H	ZP	Moist places, seeps, ditchbanks (Mar-Sep)	
<input type="radio"/> <i>Avena barbata</i>	Slender Wild Oat	Grass-like	- - -	M	ZA	Disturbed sites (Mar-Jun)	
<input type="radio"/> <i>Avena fatua</i>	Wild Oat	Grass-like	- - -	M	ZA	Disturbed sites (Apr-Jun)	
<input type="radio"/> <i>Brachypodium distachyon</i>	Purple False-brome	Grass-like	- - -	M	ZA	Disturbed areas, dry slopes (Apr-Jul)	
<input type="radio"/> <i>Briza maxima</i>	Rattlesnake Grass	Grass-like	- - -	L	ZA	Shaded sites, roadsides, pastures, weedy on coastal dunes (Apr-Jul)	
<input type="radio"/> <i>Briza minor</i>	Little Quaking Grass	Grass-like	- - -	-	ZA	Shaded or moist, open sites (Apr-Jul)	
<input type="radio"/> <i>Bromus arenarius</i>	Australian Chess	Grass-like	- - -	-	ZA	Open, disturbed places (Apr-Jul)	
<input type="radio"/> <i>Bromus carinatus var. carinatus</i>	California Brome	Grass-like	- - -	-	NP	Coastal prairies, openings in chaparral, plains, open oak and pine woodland (Apr-Aug)	
<input type="radio"/> <i>Bromus catharticus var. catharticus (JM93: no var.)</i>	Rescue Grass	Grass-like	- - -	-	ZAP	Open, disturbed places (Apr-Nov)	

Poaceae	Sorted Alphabetically by Family, Scientific Name						Poaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Bromus diandrus</i>	Ripgut Grass	Grass-like	- - -	M	ZA	Open, gen disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Bromus hordeaceus</i>	Soft Chess	Grass-like	- - -	L	ZA	Fields, disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Bromus japonicus</i>	Japanese Brome	Grass-like	- - -	L	ZA	Open, disturbed areas (May–Jul)	
<input type="radio"/> <i>Bromus laevipes</i>	Woodland Brome	Grass-like	- - -	-	NP	Shrubland, conifer forest, shaded streambanks, roadsides (May–Jul)	
<input type="radio"/> <i>Bromus madritensis subsp. madritensis</i>	Foxtail Chess	Grass-like	- - -	-	ZA	Disturbed areas, roadsides (Apr–Jan)	
<input type="radio"/> <i>Bromus madritensis subsp. rubens</i>	Red Brome	Grass-like	- - -	H	ZA	Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Bromus sterilis</i>	Poverty Brome	Grass-like	- - -	-	ZA	Open, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Bromus tectorum</i>	Cheat Grass	Grass-like	- - -	H	ZA	Open, disturbed areas (May–Aug)	
<input type="radio"/> <i>Calamagrostis rubescens</i>	Pine Reed Grass	Grass-like	- - B	-	NP	Wooded slopes, montane forest, chaparral, meadows (Jun–Sep)	
<input type="radio"/> <i>Cortaderia jubata</i>	Hairy Pampas Grass	Grass-like	- - -	H	ZP	Disturbed sites, many habitats, esp coastal (Sep–Feb)	
<input type="radio"/> <i>Cortaderia selloana</i>	Smooth Pampas Grass	Grass-like	- - -	H	ZP	Disturbed sites (Sep–Mar)	
<input type="radio"/> <i>Crypsis schoenoides</i>	Swamp Prickle Grass	Grass-like	- - -	-	ZA	Wet places (Jun–Oct)	
<input type="radio"/> <i>Crypsis vaginiflora</i>	Prickle Grass	Grass-like	- - -	-	ZA	Wet soils, lake margins, vernal pools (Jun–Sep)	
<input type="radio"/> <i>Cynodon dactylon</i>	Bermuda Grass	Grass-like	- - -	M	ZP	Disturbed sites (Jun–Aug)	
<input type="radio"/> <i>Cynosurus echinatus</i>	Bristly Dogtail Grass	Grass-like	- - -	M	ZA	Open, disturbed sites (May–Jul)	
<input type="radio"/> <i>Dactylis glomerata</i>	Orchard Grass	Grass-like	- - -	L	ZP	Disturbed, often moist sites (May–Aug)	
<input type="radio"/> <i>Danthonia californica (JM93: var. californica)</i>	California Oat Grass	Grass-like	- - C	-	NP	Gen moist meadows, open woodland (Apr–Aug)	
<input type="radio"/> <i>Deschampsia cespitosa subsp. cespitosa</i>	Tufted Hair Grass	Grass-like	- - C	-	NP	Meadows, streambanks, coastal marshes, forest (Jul–Aug)	
<input type="radio"/> <i>Deschampsia cespitosa subsp. holciformis</i>	Pacific Hair Grass	Grass-like	- - A2	-	NP	Coastal marshes, meadows (May–Jul)	
<input type="radio"/> <i>Deschampsia danthonioides</i>	Annual Hair Grass	Grass-like	- - C	-	NA	Moist to drying, open sites, meadows, streambanks, vernal pools, occ alkali soil (Mar–Aug)	
<input type="radio"/> <i>Deschampsia elongata</i>	Slender Hair Grass	Grass-like	- - C	-	NP	Wet sites, meadows, lakeshores, shaded slopes (May–Sep)	
<input type="radio"/> <i>Digitaria ischaemum</i>	Smooth Crab Grass	Grass-like	- - -	-	ZA	Disturbed areas (Sep–Nov)	
<input type="radio"/> <i>Digitaria sanguinalis</i>	Hairy Crab Grass	Grass-like	- - -	-	ZA	Disturbed areas (Jun–Sep)	
<input type="radio"/> <i>Distichlis spicata</i>	Salt Grass	Grass-like	- - -	-	NP	Salt marshes, coastal dunes, moist, alkaline areas (Apr–Sep)	
<input type="radio"/> <i>Echinochloa crus-galli</i>	Barnyard Grass	Grass-like	- - -	-	ZA	Gen wet, disturbed sites, fields, roadsides (Jun–Oct)	
<input type="radio"/> <i>Ehrharta erecta</i>	Panic Veldt Grass	Grass-like	- - -	M	ZP	Disturbed, often shady, moist sites (Mar–Jun)	
<input type="radio"/> <i>Elymus caput-medusae (JM93: Taeniatherum)</i>	Medusa Head	Grass-like	- - -	H	ZA	Disturbed areas (Apr–Jul)	
<input type="radio"/> <i>Elymus condensatus (JM93: Leymus)</i>	Giant Wild-rye	Grass-like	- - -	-	NP	Dry slopes, open woodland (Jun–Aug)	

Poaceae	Sorted Alphabetically by Family, Scientific Name						Poaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Elymus glaucus</i> subsp. <i>glaucus</i>	Western Wild-rye	Grass-like	- - -	-	NP	Open areas, chaparral, woodland, forest (Jun–Aug)	
<input type="radio"/> <i>Elymus multisetus</i>	Big Squirreltail	Grass-like	- - C	-	NP	Open, sandy to rocky areas (May–Jul)	
<input type="radio"/> <i>Elymus stebbinsii</i>	Parish's Wheat Grass	Grass-like	- - A2	-	NP	Dry slopes, chaparral, conifer forest (Jun–Jul)	
<input type="radio"/> <i>Elymus trachycaulus</i> subsp. <i>trachycaulus</i>	Slender Wheat Grass	Grass-like	- - B	-	NP	Dry to moist, open areas, forest, woodland (Jun–Aug)	
<input type="radio"/> <i>Elymus triticoides</i> (JM93: <i>Leymus</i>)	Beardless Wild Rye	Grass-like	- - -	-	NP	Dry to moist, often saline, meadows (Jun–Jul)	
<input type="radio"/> <i>Eragrostis curvula</i> (JM93: var. <i>curvula</i>)	Weeping Love Grass	Grass-like	- - -	-	NP	Roadsides (Aug–Oct)	
<input type="radio"/> <i>Festuca arundinacea</i>	Tall Fescue	Grass-like	- - -	M	ZP	Disturbed places (May–Jun)	
<input type="radio"/> <i>Festuca bromoides</i> (JM93: <i>Vulpia</i>)	Brome Fescue	Grass-like	- - -	-	ZA	Uncommon. Dry, disturbed places, coastal-sage scrub, chaparral (May–Jun)	
<input type="radio"/> <i>Festuca californica</i>	California Fescue	Grass-like	- - C	-	NP	Dry, open forest, moist streambanks, chaparral (May–Jun)	
<input type="radio"/> <i>Festuca elmeri</i>	Elmer Fescue	Grass-like	- - A2	-	NP	Moist, wooded slopes, under trees in rich soil (May–Jul)	
<input type="radio"/> <i>Festuca idahoensis</i>	Idaho Fescue	Grass-like	- - C	-	NP	Dry, open or shady places (Jul–Sep)	
<input type="radio"/> <i>Festuca microstachys</i> (JM93: <i>Vulpia</i> , var. <i>ciliata</i> , <i>confusa</i> , <i>microstachys</i> , <i>pauciflora</i>)	Hairy Fescue	Grass-like	- - -	-	NA	Disturbed, open, gen sandy soils (Apr–Jun)	
<input type="radio"/> <i>Festuca myuros</i> (JM93: <i>Vulpia</i> , var. <i>hirsuta</i> , <i>myuros</i>)	Rattail Sixweeks Grass	Grass-like	- - -	-	ZA	Common. Generally open places, sandy soils (Feb–May)	
<input type="radio"/> <i>Festuca octoflora</i> (JM93: <i>Vulpia</i> , var. <i>hirtella</i> , <i>octoflora</i>)	Slender Six-weeks Fescue	Grass-like	- - A2	-	NA	Sandy to rocky soils, open sites (Mar–Jun)	
<input type="radio"/> <i>Festuca perennis</i> (JM93: <i>Lolium perenne</i> , <i>L. multiflorum</i>)	Rye Grass	Grass-like	- - -	-	ZP	Dry to moist disturbed sites, abandoned fields (May–Sep)	
<input type="radio"/> <i>Festuca rubra</i>	Red Fescue	Grass-like	- - B	-	NP	Sand dunes, grassland, subalpine forest (May–Jul)	
<input type="radio"/> <i>Festuca</i> sp. (annual) (JM93: <i>Vulpia</i> sp.)	Annual Fescue	Grass-like	- - -	-	NA	--	
<input type="radio"/> <i>Festuca</i> sp. (perennial)	Perennial Fescue	Grass-like	- - -	-	--	--	
<input type="radio"/> <i>Festuca temulenta</i> (JM93: <i>Lolium temulentum</i>)	Long-glumed Rye Grass	Grass-like	- - -	-	ZA	Open, disturbed sites (Apr–Jun)	
<input type="radio"/> <i>Gastridium phleoides</i> (JM93: <i>G. ventricosum</i>)	Nit Grass	Grass-like	- - -	-	ZA	Open, generally dry, disturbed sites (Apr–Nov)	
<input type="radio"/> <i>Glyceria occidentalis</i>	Western Manna Grass	Grass-like	- - A2	-	NP	Freshwater marshes, ponds and steams (Jun–Aug)	
<input type="radio"/> <i>Holcus lanatus</i>	Common Velvet Grass	Grass-like	- - -	M	ZP	Moist sites, roadbanks, cult fields, meadows (Jun–Aug)	
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. ?	California / Meadow Barley	Grass-like	- - -	-	NP		
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>brachyantherum</i>	Northern Barley	Grass-like	- - -	-	NP	Meadows, pastures, streambanks (May–Aug)	
<input type="radio"/> <i>Hordeum brachyantherum</i> subsp. <i>californicum</i>	California Barley	Grass-like	- - B	-	NP	Meadows, pastures, streambanks (Mar–Jul)	

Poaceae	Sorted Alphabetically by Family, Scientific Name					Poaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
○ <i>Hordeum depressum</i>	Low Barley	Grass-like	- - B	- NA	Moist sites, vernal pools, generally alkaline soils (Apr–May)	
○ <i>Hordeum jubatum</i> subsp. <i>jubatum</i> (JM93: no subsp.)	Foxtail Barley	Grass-like	- - B	- NAP	Roadsides, disturbed areas, meadows, marshes (May–Jul)	
○ <i>Hordeum marinum</i> subsp. <i>gussoneanum</i>	Mediterranean Barley	Grass-like	- - -	- ZA	Dry to moist, disturbed sites (Apr–Jun)	
○ <i>Hordeum murinum</i> subsp. <i>glaucum</i>	Smooth Barley	Grass-like	- - -	- ZA	Moist, generally disturbed sites (Mar–Jul)	
○ <i>Hordeum murinum</i> subsp. <i>leporinum</i>	Hare Barley	Grass-like	- - -	- ZA	Moist, gen disturbed sites. Common (Feb–May)	
○ <i>Hordeum murinum</i> subsp. <i>murinum</i>	Wall Barley	Grass-like	- - -	- ZA	Moist, gen disturbed sites (Feb–May)	
○ <i>Koeleria gerardii</i> (JM93: var. <i>hindsii</i>)	Annual June Grass	Grass-like	- - -	- ZA	Open, disturbed sites (Apr–Jul)	
○ <i>Koeleria macrantha</i>	June Grass	Grass-like	- - C	- NP	Dry, open sites, clay to rocky soils, shrubland, woodland, conifer forest (May–Jul)	
○ <i>Lamarckia aurea</i>	Goldentop	Grass-like	- - -	- ZA	Open ground, moist seeps, rocky hillsides, sandy soil (Feb–May)	
○ <i>Leersia oryzoides</i>	Rice Cut Grass	Grass-like	- - A2	- NP	Marshes, streams, ponds (Aug–Oct)	
○ <i>Leptochloa fusca</i> subsp. <i>fascicularis</i> (JM93: <i>L. fascicularis</i>)	Bearded Sprangletop	Grass-like	- - A2	- NA	Marshes, wetlands, often wet disturbed areas, often associated with agriculture (Jun–Oct)	
○ <i>Melica bulbosa</i>	Oniongrass	Grass-like	- - A1?	- NP	Dry rocky slopes, conifer forest (Jul–Aug)	
○ <i>Melica californica</i>	California Melic	Grass-like	- - C	- NP	Open or rocky hillsides, oak woodland, conifer forest (Apr–May)	
○ <i>Melica imperfecta</i>	Little California Melica	Grass-like	- - -	- NP	Dry rocky hillsides, chaparral, woodland (Apr–May)	
○ <i>Melica torreyana</i>	Torrey's Melic	Grass-like	- - -	- NP	Chaparral, conifer forest (Mar–Jun)	
○ <i>Panicum capillare</i>	Witch Grass	Grass-like	- - B	- NA	Open places, fields, roadsides (Feb–Dec)	
○ <i>Panicum sp.</i>	Panic Grass	Grass-like	- - -	-	- -	
○ <i>Parapholis incurva</i>	Sickle Grass	Grass-like	- - -	- ZA	Disturbed, well drained soils of salt marshes, generally above highest tide level (Apr–Jun)	
○ <i>Paspalum dilatatum</i>	Dallis Grass	Grass-like	- - -	- ZP	Disturbed areas (May–Nov)	
○ <i>Paspalum distichum</i>	Knot Grass	Grass-like	- - -	- ZP	Edges of lakes, ponds, rice fields, wet roadside ditches (Jun–Oct)	
○ <i>Paspalum sp.</i>	Dallis / Knot Grass	Grass-like	- - -	- ZP	- -	
○ <i>Pennisetum clandestinum</i>	Kikuyu Grass	Grass-like	- - -	L ZP	Disturbed areas (All year)	
○ <i>Phalaris aquatica</i>	Harding Grass	Grass-like	- - -	M ZP	Disturbed areas, roadsides (Apr–Aug)	
○ <i>Phalaris arundinacea</i>	Reed Canary Grass	Grass-like	- - A2	- NP	Moist habitats, woodland (May–Sep)	
○ <i>Phalaris canariensis</i>	Canary Grass	Grass-like	- - -	- ZA	Disturbed areas (Apr–Jul)	
○ <i>Phalaris minor</i>	Little-seeded Canary Grass	Grass-like	- - -	- ZA	Disturbed areas, cultivated fields (Apr–Jul)	
○ <i>Phalaris paradoxa</i>	Hood Canary Grass	Grass-like	- - -	- ZA	Disturbed areas, cultivated field (May–Aug)	
○ <i>Phragmites australis</i>	Common Reed	Grass-like	- - B	N NP	Pond & lake margins, sloughs, marshes (Jul–Nov)	

Poaceae		Sorted Alphabetically by Family, Scientific Name					Poaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Pleuropogon californicus</i> var. <i>californicus</i> (JM93: no var.)	California Semaphore Grass	Grass-like	- - B	-	NA	Wet places, vernal pools, wet grassland (Mar–Jun)	
<input type="radio"/> <i>Poa annua</i>	Annual Blue Grass	Grass-like	- - -	-	ZA	Abundant. Disturbed moist ground (Feb–Sep)	
<input type="radio"/> <i>Poa bulbosa</i> subsp. <i>vivipara</i> (JM93: no subsp.)	Bulbous Blue Grass	Grass-like	- - -	-	ZP	Disturbed places (Mar–Jul)	
<input type="radio"/> <i>Poa howellii</i>	Howell Blue Grass	Grass-like	- - B	-	NA	Rocky banks, shaded slopes of woodland, chaparral, disturbed places (Apr–Jun)	
<input type="radio"/> <i>Poa pratensis</i> subsp. <i>pratensis</i>	Kentucky Blue Grass	Grass-like	- - -	-	ZP	Common. Disturbed habitats, including saline or alkaline soils (Mar–Jul)	
<input type="radio"/> <i>Poa secunda</i> subsp. <i>secunda</i>	One-sided Blue Grass	Grass-like	- - -	-	NP	Common. Dry slopes to saline/alkaline meadows to alpine (Mar–Aug)	
<input type="radio"/> <i>Polypogon interruptus</i>	Ditch Beard Grass	Grass-like	- - -	-	ZP	Common. Streambanks (May–Aug)	
<input type="radio"/> <i>Polypogon monspeliensis</i>	Rabbitfoot Grass	Grass-like	- - -	L	ZA	Moist places, along streams (Apr–Aug)	
<input type="radio"/> <i>Polypogon viridis</i> (JM93: <i>Agrostis</i>)	Water Beard Grass	Grass-like	- - -	-	ZP	Common. Disturbed areas, wet areas, ponds, streambanks (May–Jun)	
<input type="radio"/> <i>Puccinellia nutkaensis</i>	Alaska Alkali Grass	Grass-like	- - A1	-	NP	Marshes, wet sites (Jun)	
<input type="radio"/> <i>Puccinellia nuttalliana</i>	Nuttall Alkali Grass	Grass-like	- - A1	-	NP	Saline meadows (Jun–Sep)	
<input type="radio"/> <i>Puccinellia simplex</i>	California Alkali Grass	Grass-like	- - A2	-	NA	Saline flats, mineral springs (Mar–May)	
<input type="radio"/> <i>Scribneria bolanderi</i>	Scribner Grass	Grass-like	- - A2	-	NP	Dry, disturbed areas (Mar–Jun)	
<input type="radio"/> <i>Sorghum bicolor</i>	Sorghum	Grass-like	- - -	-	ZA	Disturbed areas, roadsides (All year)	
<input type="radio"/> <i>Sorghum halepense</i>	Johnson Grass	Grass-like	- - -	-	ZP	Disturbed areas, ditch banks, roadsides (May–Aug)	
<input type="radio"/> <i>Sorghum</i> sp.	Johnson Grass / Sorghum	Grass-like	- - -	-	ZAP	- -	
<input type="radio"/> <i>Spartina alterniflora</i>	Smooth Saltwater Cordgrass	Grass-like	- - -	H	ZP	Coastal salt marshes (Jul–Sep)	
<input type="radio"/> <i>Spartina densiflora</i>	Dense-flowered Cordgrass	Grass-like	- - -	H	ZP	Coastal salt marshes (May–Aug)	
<input type="radio"/> <i>Spartina foliosa</i>	California Cordgrass	Grass-like	- - B	-	NP	Salt marshes, mudflats, shores (Jun–Nov)	
<input type="radio"/> <i>Sporobolus airoides</i>	Alkali Sacaton	Grass-like	- - A2	-	NP	Seasonally moist, alkaline areas (Apr–Oct)	
<input type="radio"/> <i>Stipa cernua</i> (JM93: <i>Nassella</i>)	Nodding Needle Grass	Grass-like	- - C	-	NP	Grassland, chaparral, juniper woodland (Feb–Jul)	
<input type="radio"/> <i>Stipa lepida</i> (JM93: <i>Nassella</i>)	Foothill Needle Grass	Grass-like	- - C	-	NP	Dry slopes, chaparral, grassland, savanna, coastal scrub (Mar–Jun)	
<input type="radio"/> <i>Stipa miliacea</i> var. <i>miliacea</i> (JM93: <i>Piptatherum miliaceum</i>)	Smilo Grass	Grass-like	- - -	L	ZP	Salt marshes, streambanks, chaparral, open woodland, disturbed (Mar–Sep)	
<input type="radio"/> <i>Stipa pulchra</i> (JM93: <i>Nassella</i>)	Purple Needle Grass	Grass-like	- - C	-	NP	Oak woodland, chaparral, grassland (Mar–Jun)	
<input type="radio"/> <i>Trisetum canescens</i>	Tall False Oat	Grass-like	- - B	-	NP	Open to shaded sites, meadows, chaparral, conifer forest (May–Aug)	
Polemoniaceae							Polemoniaceae
<input type="radio"/> <i>Allophyllum divaricatum</i>	Straggling Gilia	Herbaceous	- - A2	-	NA	Sandy areas, chaparral, woodland (Apr–Jun)	
<input type="radio"/> <i>Allophyllum gilioides</i> subsp. <i>gilioides</i>	Purple False Gilia	Herbaceous	- - B	-	NA	Open, sandy, gen damp or grassy areas (Apr–Jun)	

Polemoniaceae		Sorted Alphabetically by Family, Scientific Name				Polemoniaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Collomia heterophylla</i>	Variable-leaf Collomia	Herbaceous	- - A2	-	NA	Sandy to gravelly, open areas (Apr–Jun)	
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>achilleifolia</i>	California Gilia	Herbaceous	- - B	-	NA	Open or shaded, gen grassy places, sandy or rocky soil (Mar–Jun)	
<input type="radio"/> <i>Gilia achilleifolia</i> subsp. <i>multicaulis</i>	Many-stem California Gilia	Herbaceous	- - -	-	NA	Open or shaded, gen grassy places, sandy or rocky soil (Feb–Jun)	
<input type="radio"/> <i>Gilia capitata</i> subsp. ?	Globe Gilia	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Gilia capitata</i> subsp. <i>staminea</i>	Pale Gilia	Herbaceous	- - B	-	NA	Sandhills, flats (Mar–May)	
<input type="radio"/> <i>Gilia clivorum</i>	Purplespot Gilia	Herbaceous	- - C	-	NA	Common. Open, grassy areas (Feb–Jun)	
<input type="radio"/> <i>Gilia</i> sp.	Gilia	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Gilia tricolor</i> subsp. ?	Bird's-eye Gilia	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Gilia tricolor</i> subsp. <i>diffusa</i>	Diffuse Bird's-eye Gilia	Herbaceous	- - B	-	NA	Open grassland, hills, valleys (Mar–Jul)	
<input type="radio"/> <i>Gilia tricolor</i> subsp. <i>tricolor</i>	Bird's-eye Gilia	Herbaceous	- - B	-	NA	Open, grassland, hills, valleys (Mar–May)	
<input type="radio"/> <i>Leptosiphon acicularis</i> (JM93: <i>Linanthus</i>)	Bristly Leptosiphon	Herbaceous	- 4.2 *A1	-	NA	Grassy areas, woodland, chaparral (Apr–May)	
<input type="radio"/> <i>Leptosiphon ambiguus</i> (JM93: <i>Linanthus</i>)	Serpentine Leptosiphon	Herbaceous	- 4.2 *A2	-	NA	Grassy areas gen serpentine (Apr–May)	
<input type="radio"/> <i>Leptosiphon androsaceus</i> (JM93: <i>Linanthus</i>)	Pinklobe Leptosiphon	Herbaceous	- - C	-	NA	Open or shaded areas in woodland, chaparral (Apr–Jun)	
<input type="radio"/> <i>Leptosiphon bicolor</i> (JM93: <i>Linanthus</i>)	Bicolor Leptosiphon	Herbaceous	- - -	-	NA	Common. Open, grassy areas, chaparral, woodland (Mar–Jun)	
<input type="radio"/> <i>Leptosiphon ciliatus</i> (JM93: <i>Linanthus</i>)	Whisker Brush	Herbaceous	- - B	-	NA	Common. Open or wooded areas (Mar–Jul)	
<input type="radio"/> <i>Leptosiphon parviflorus</i> (JM93: <i>Linanthus</i>)	Common Linanthus	Herbaceous	- - B	-	NA	Abundant. Open or wooded areas (Mar–Jun)	
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. ? (JM93: no subsp.)	Evening Snow	Herbaceous	- - A2	-	NA	--	
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. <i>dichotomus</i> (JM93: no subsp.)	Evening Snow	Herbaceous	- - A2	-	NA	Common. Drying openings, especially serpentine (Apr–Jun)	
<input type="radio"/> <i>Linanthus dichotomus</i> subsp. <i>meridianus</i> (JM93: no subsp.)	Day-blooming Evening Snow	Herbaceous	- - A2	-	NA	Drying openings, especially serpentine (Apr–Jun)	
<input type="radio"/> <i>Linanthus</i> or <i>Leptosiphon</i> sp. (JM93: <i>Linanthus</i> sp.)	Linanthus	Herbaceous	- - -	-	NA	--	
<input type="radio"/> <i>Microsteris gracilis</i> (JM93: <i>Phlox</i>)	Slender Annual Phlox	Herbaceous	- - -	-	NA	Dry to moist areas (Mar–Aug)	
<input type="radio"/> <i>Navarretia atractyloides</i>	Holly-leaf Navarretia	Herbaceous	- - A2	-	NA	Open, rocky or sandy areas (May–Jul)	
<input type="radio"/> <i>Navarretia intertexta</i> subsp. <i>intertexta</i>	Needle-leaved Navarretia	Herbaceous	- - A1	-	NA	Open, seasonally wet areas, meadows, vernal pools (May–Jul)	
<input type="radio"/> <i>Navarretia mellita</i>	Honeyscent Navarretia	Herbaceous	- - C	-	NA	Open, wet, sandy or gravelly areas, chaparral (May–Jul)	
<input type="radio"/> <i>Navarretia pubescens</i>	Downy Navarretia	Herbaceous	- - C	-	NA	Open, slopes, gravel, clay (May–Jul)	
<input type="radio"/> <i>Navarretia squarrosa</i>	Skunkweed	Herbaceous	- - -	-	NA	Common. Open, wet, gravelly flats, slopes (Jun–Aug)	

Polygalaceae		Sorted Alphabetically by Family, Scientific Name				Polygalaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
Polygalaceae						Polygalaceae
<input type="radio"/> <i>Polygala californica</i>	California Milkwort	Herbaceous	- - A1	- NP	Coastal prairie and forest, chaparral, occasionally on serpentine (Apr–Jul)	
Polygonaceae						Polygonaceae
<input type="radio"/> <i>Chorizanthe membranacea</i>	Pink Spineflower	Herbaceous	- - A2	- NA	Common. Sand, gravel or rocks (Apr–Jul)	
<input type="radio"/> <i>Chorizanthe polygonoides</i> var. <i>polygonoides</i>	Knotweed Spineflower	Herbaceous	- - A1	- NA	Sand or gravel (Apr–Jun)	
<input type="radio"/> <i>Eriogonum angulosum</i>	Angle-stem Wild Buckwheat	Herbaceous	- - A2	- NA	Common. Clay (All year)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. ?	California Buckwheat	Woody	- - -	- NP	--	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>foliolosum</i>	Leafy California Buckwheat	Woody	- - A2	- NP	Gravel (All year)	
<input type="radio"/> <i>Eriogonum fasciculatum</i> var. <i>polifolium</i>	Mojave Desert Calif. Buckwheat	Woody	- - A1	- NP	Sand, gravel or rocks (All year)	
<input type="radio"/> <i>Eriogonum gracile</i> var. <i>gracile</i>	Slender Woolly Wild Buckwheat	Herbaceous	- - B	- NA	Common. Sand (All year)	
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>caninum</i>	Tiburon Buckwheat	Herbaceous	- 1B.2 *A1	- NA	Serpentine (May–Oct)	
<input type="radio"/> <i>Eriogonum luteolum</i> var. <i>luteolum</i>	Golden-carpet Wild Buckwheat	Herbaceous	- - A2	- NA	Common. Serpentine (Jul–Nov)	
<input type="radio"/> <i>Eriogonum nudum</i> var. ?	Naked-stem Buckwheat	Herbaceous	- - -	- NP	--	
<input type="radio"/> <i>Eriogonum nudum</i> var. <i>auriculatum</i>	Ear-shaped Wild Buckwheat	Herbaceous	- - A1	- NP	Common. Sand or gravel (May–Oct)	
<input type="radio"/> <i>Eriogonum roseum</i>	Wand Wild Buckwheat	Herbaceous	- - B	- NA	Common. Sand or gravel (May–Nov)	
<input type="radio"/> <i>Eriogonum</i> sp.	Wild Buckwheat	Herbaceous	- - -	- NAP	--	
<input type="radio"/> <i>Eriogonum umbellatum</i> var. <i>smallianum</i> (JM93: var. <i>bahiiiforme</i>)	Small's Sulphur Flower	Herbaceous	- - *A2	- NP	Serpentine (Jul–Sep)	
<input type="radio"/> <i>Eriogonum wrightii</i> var. <i>trachygonum</i>	Rough-node Bastard-sage	Herbaceous	- - A2	- NP	Common. Gravel (Jul–Oct)	
<input type="radio"/> <i>Persicaria amphibia</i> (JM93: <i>Polygonum amphibia</i> var. <i>emersum</i> , <i>stipulaceum</i>)	Water Smartweed	Herbaceous	- - -	- NP	Shallow lakes, streams, shores (Jun–Nov)	
<input type="radio"/> <i>Persicaria hydropiperoides</i> (JM93: <i>Polygonum</i>)	False Waterpepper	Herbaceous	- - A2	- NP	Wet banks, shallow water, marshes, moist prairies (Jun–Oct)	
<input type="radio"/> <i>Persicaria lapathifolia</i> (JM93: <i>Polygonum lapathifolium</i>)	Willow Weed	Herbaceous	- - C	- NA	Moist places, disturbed areas, roadsides, fields (Jun–Oct)	
<input type="radio"/> <i>Persicaria maculosa</i> (JM93: <i>Polygonum persicaria</i>)	Lady's Thumb	Herbaceous	- - -	- ZA	Moist disturbed areas, field (Jun–Nov)	
<input type="radio"/> <i>Persicaria punctata</i> (JM93: <i>Polygonum punctatum</i>)	Water Smartweed	Herbaceous	- - C	- NAP	Shallow water, shores, marshes, floodplain forest (Jun–Nov)	
<input type="radio"/> <i>Polygonum aviculare</i> subsp. <i>depressum</i> (JM93: <i>P. arenastrum</i>)	Knotweed	Herbaceous	- - -	- ZA	Disturbed places (May–Nov)	
<input type="radio"/> <i>Polygonum</i> or <i>Persicaria</i> sp. (JM93: <i>Polygonum</i> sp.)	Knotweed / Smartweed	Herbaceous	- - -	-	--	
<input type="radio"/> <i>Pterostegia drymarioides</i>	Pink Creeper	Herbaceous	- - -	- NA	Common. Sand or gravel (Mar–Jul)	

Polygonaceae		Sorted Alphabetically by Family, Scientific Name				Polygonaceae	
<input checked="" type="checkbox"/> Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="checkbox"/> <i>Rumex acetosella</i>	Sheep Sorrel	Herbaceous	- - -	M	ZP	± Disturbed, often acidic places (Apr–Jul)	
<input type="checkbox"/> <i>Rumex conglomeratus</i>	Green Dock	Herbaceous	- - -	-	ZP	Common. Moist places (May–Aug)	
<input type="checkbox"/> <i>Rumex crassus</i> (JM93: <i>R. salicifolius</i> var. <i>crassus</i>)	Dense Willow Dock	Herbaceous	- - A2	-	NP	Coastal dunes, sandy shores, marshes (Feb–Jul)	
<input type="checkbox"/> <i>Rumex crispus</i>	Curly Dock	Herbaceous	- - -	L	ZP	Abundant. Disturbed places (All year)	
<input type="checkbox"/> <i>Rumex fueginus</i> (JM93: <i>R. maritimus</i>)	Golden Dock	Herbaceous	- - A2	-	NA	Riparian, disturbed places, shores, marshes, bogs, wet meadows (May–Aug)	
<input type="checkbox"/> <i>Rumex obtusifolius</i>	Bitter Dock	Herbaceous	- - -	-	ZP	Moist places (May–Sep)	
<input type="checkbox"/> <i>Rumex occidentalis</i>	Western Dock	Herbaceous	- - A1	-	NP	Uncommon. Wet habitats (May–Aug)	
<input type="checkbox"/> <i>Rumex pulcher</i>	Fiddle Dock	Herbaceous	- - -	-	ZP	Disturbed places, meadows, moist or dry habitats (May–Sep)	
<input type="checkbox"/> <i>Rumex salicifolius</i> (JM93: var. <i>salicifolius</i>)	Thin-leaf Willow Dock	Herbaceous	- - B	-	NP	Wet places, margins, rocky slopes (May–Jul)	
<input type="checkbox"/> <i>Rumex sp.</i>	Dock	Herbaceous	- - -	-	-		
<input type="checkbox"/> <i>Rumex transitorius</i> (JM93: <i>R. salicifolius</i> var. <i>transitorius</i>)	Ovate Willow Dock	Herbaceous	- - -	-	NP	Coastal dunes, wet margins of meadows (Apr–Jun)	
Polypodiaceae						Polypodiaceae	
<input type="checkbox"/> <i>Polypodium calirhiza</i>	Polypody Fern	Fern-like	- - -	-	NP	On plants, rocky cliffs or outcrops, roadcuts, often granitic or volcanic, rarely dunes -	
<input type="checkbox"/> <i>Polypodium scolieri</i>	Leather-leaf Fern	Fern-like	- - A1	-	NP	Coast, generally in heavy fog-drip or salt-spray zones, on plants (especially Douglas fir, Eucalyptus), dunes, rocky cliffs, bluffs, mtn ridges, granitic or volcanic rocks, mossy logs, or in soil -	
Pontederiaceae						Pontederiaceae	
<input type="checkbox"/> <i>Eichhornia crassipes</i>	Water Hyacinth	Herbaceous	- - -	H	ZP	Locally abundant. Ponds, sloughs, waterways (Jun–Oct)	
Portulacaceae						Portulacaceae	
<input type="checkbox"/> <i>Portulaca oleracea</i>	Purslane	Herbaceous	- - -	-	ZP	Disturbed soil (Late spring-early fall)	
Potamogetonaceae						Potamogetonaceae	
<input type="checkbox"/> <i>Potamogeton crispus</i>	Crisp-leaved Pondweed	Herbaceous	- - -	M	ZP	Uncommon. Shallow water, ponds, streams. (Jul–Sep)	
<input type="checkbox"/> <i>Potamogeton illinoensis</i>	Shining Pondweed	Herbaceous	- - -	-	NP	Lakes, ponds, streams (Jun–Aug)	
<input type="checkbox"/> <i>Potamogeton nodosus</i>	Long-leaved Pondweed	Herbaceous	- - -	-	NP	Shallow water, lakes, ponds, streams (May–Aug)	
<input type="checkbox"/> <i>Potamogeton</i> or <i>Stuckenia</i> sp. (JM93: <i>Potamogeton</i> sp.)	Pondweed	Herbaceous	- - -	-	NP	-	

Potamogetonaceae		Sorted Alphabetically by Family, Scientific Name				Potamogetonaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Stuckenia pectinata</i> (JM93: <i>Potamogeton pectinatus</i>)	Fennel-leaf Pondweed	Herbaceous	- - -	-	NP	Common. Ponds, lakes, marshes, streams (May–Jul)
Primulaceae						Primulaceae
<input type="radio"/> <i>Dodecatheon clevelandii</i> subsp. ?	Shooting Star	Herbaceous	- - -	-	NP	- -
<input type="radio"/> <i>Dodecatheon clevelandii</i> subsp. <i>patulum</i>	Padre Shooting Star	Herbaceous	- - C	-	NP	Moist places, often on serpentine or in ± alkaline sites (Mar–May)
<input type="radio"/> <i>Dodecatheon hendersonii</i>	Mosquitobills Shooting Star	Herbaceous	- - -	-	NP	Gen in shady sites (Mar–Jul)
Pteridaceae						Pteridaceae
<input type="radio"/> <i>Adiantum aleuticum</i>	Five-finger Fern	Fern-like	- - A2	-	NP	Shady, moist banks, streamsides, serpentine -
<input type="radio"/> <i>Adiantum jordanii</i>	California Maidenhair	Fern-like	- - -	-	NP	Shaded hillsides, moist woodland -
<input type="radio"/> <i>Aspidotis californica</i>	California Lace Fern	Fern-like	- - A2	-	NP	Rock outcrops, crevices -
<input type="radio"/> <i>Cheilanthes covillei</i>	Coville's Lip Fern	Fern-like	- - A2	-	NP	Crevices, bases of rocks, sun or shade -
<input type="radio"/> <i>Cheilanthes intertexta</i>	Coastal Lip Fern	Fern-like	- - A2	-	NP	Crevices, bases of rocks -
<input type="radio"/> <i>Pellaea andromedifolia</i>	Coffee Fern	Fern-like	- - -	-	NP	Generally rocky or dry areas -
<input type="radio"/> <i>Pellaea mucronata</i> var. <i>mucronata</i>	Bird's-foot Fern	Fern-like	- - C	-	NP	Rocky or dry areas -
<input type="radio"/> <i>Pentagramma triangularis</i> subsp. <i>triangularis</i>	Goldenback Fern	Fern-like	- - -	-	NP	Gen shaded, sometimes rocky or wooded areas -
Ranunculaceae						Ranunculaceae
<input type="radio"/> <i>Actaea rubra</i>	Baneberry	Herbaceous	- - B	-	NP	Deep soils, moist, open to shaded sites, mixed-evergreen or conifer forests (May–Sep)
<input type="radio"/> <i>Aquilegia eximia</i>	Serpentine Columbine	Herbaceous	- - A2	-	NP	Generally serpentine seeps, occasionally moist ravines, mixed-evergreen or conifer forests (May–Oct)
<input type="radio"/> <i>Aquilegia formosa</i>	Crimson Columbine	Herbaceous	- - C	-	NP	Streambanks, seeps, moist places, chaparral, oak woodland, mixed-evergreen or conifer forests (Apr–Sep)
<input type="radio"/> <i>Clematis lasiantha</i>	Chaparral Clematis	Woody	- - -	-	NP	Hillsides, chaparral, open woodland (Jan–Jun)
<input type="radio"/> <i>Clematis ligusticifolia</i>	Western Virgin's Bower	Woody	- - C	-	NP	Along streams, wet places (Jun–Sep)
<input type="radio"/> <i>Clematis</i> sp.	Virgin's Bower / Pipestems	Woody	- - -	-	NP	- -
<input type="radio"/> <i>Delphinium californicum</i> subsp. <i>californicum</i>	California Larkspur	Herbaceous	- - A2	-	NP	Generally slopes in dense chaparral, w side of coast ranges (Apr–Jun)
<input type="radio"/> <i>Delphinium decorum</i> subsp. <i>decorum</i>	Coast Larkspur	Herbaceous	- - C	-	NP	Open coastal grassland, chaparral (Mar–May)
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>hesperium</i>	Western Larkspur	Herbaceous	- - B	-	NP	Oak woodland, w slope coast ranges (Mar–Jun)
<input type="radio"/> <i>Delphinium hesperium</i> subsp. <i>pallescens</i>	Pale-flowered Western Larkspur	Herbaceous	- - C	-	NP	Oak woodland, e slope coast ranges (Mar–May)
<input type="radio"/> <i>Delphinium nudicaule</i>	Red Or Orange Larkspur	Herbaceous	- - C	-	NP	Moist talus, wooded, rocky slopes (Mar–Jun)
<input type="radio"/> <i>Delphinium patens</i> subsp. <i>patens</i>	Spreading Larkspur	Herbaceous	- - -	-	NP	Grassland, open woodland (Mar–Jun)

Ranunculaceae		Sorted Alphabetically by Family, Scientific Name				Ranunculaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Delphinium sp.</i>	Larkspur	Herbaceous	- - -	-	NP	-	
<input type="radio"/> <i>Delphinium variegatum subsp. variegatum</i>	Royal Larkspur	Herbaceous	- - C	-	NP	Grassland, open oak woodland (Mar–May)	
<input type="radio"/> <i>Enemion occidentale (JM93: Isopyrum)</i>	Western Rue-anemone	Herbaceous	- - A1	-	NP	Shaded slopes, chaparral, oak woodland, conifer forest (Mar–May)	
<input type="radio"/> <i>Ranunculus aquatilis var. diffusus (JM93: var. capillaceus)</i>	Thread-leaf Water Buttercup	Herbaceous	- - -	-	NP	Ponds, lakes, streams (Mar–Sep)	
<input type="radio"/> <i>Ranunculus californicus var. californicus (JM93: no var.)</i>	California Buttercup	Herbaceous	- - -	-	NP	Grassland, open woodland (Mar–Aug)	
<input type="radio"/> <i>Ranunculus canus var. canus (JM93: no var.)</i>	Sacramento Valley Buttercup	Herbaceous	- - B	-	NP	Grassland, open oak woodlan (Mar–Jul)	
<input type="radio"/> <i>Ranunculus hebecarpus</i>	Downy Buttercup	Herbaceous	- - -	-	NA	Grassland, open woodland (Mar–May)	
<input type="radio"/> <i>Ranunculus lobbii</i>	Lobb's Aquatic Buttercup	Herbaceous	- 4.2 *A2	-	NA	Ponds (Mar–May)	
<input type="radio"/> <i>Ranunculus muricatus</i>	Prickleseed Buttercup	Herbaceous	- - -	-	ZA	Stream-banks, drainages, low meadows (Apr–Jun)	
<input type="radio"/> <i>Ranunculus occidentalis var. occidentalis (JM93: no var.)</i>	Western Buttercup	Herbaceous	- - A2	-	NP	Grassy slopes in meadows or open woodland (Mar–Jul)	
<input type="radio"/> <i>Ranunculus orthorhynchus var. bloomeri</i>	Bloomer Buttercup	Herbaceous	- - A1	-	NP	Meadows, marshy areas (Mar–May)	
<input type="radio"/> <i>Ranunculus repens</i>	Creeping Buttercup	Herbaceous	- - -	L	ZP	Meadows, road banks, disturbed places, edges of marshes, streams (All year)	
<input type="radio"/> <i>Ranunculus sceleratus var. sceleratus</i>	Cursed Crowsfoot	Herbaceous	- - -	-	NA	Ponds, riverbanks (Apr–Jun,Oct)	
<input type="radio"/> <i>Ranunculus sp.</i>	Buttercup	Herbaceous	- - -	-	-	-	
<input type="radio"/> <i>Thalictrum fendleri var. polycarpum</i>	Foothill Meadow-rue	Herbaceous	- - -	-	NP	Moist, open to shaded places, woodland, forest (Mar–Jun)	
Rhamnaceae		Rhamnaceae					
<input type="radio"/> <i>Ceanothus cuneatus var. cuneatus</i>	Buckbrush	Woody	- - C	-	NP	Sandy to rocky flats, slopes, ridges (Feb–May)	
<input type="radio"/> <i>Ceanothus oliganthus var. sorediatus</i>	Jim Brush	Woody	- - -	-	NP	Slopes, ridges, chaparral, conifer forest (Jan–May)	
<input type="radio"/> <i>Ceanothus thyrsiflorus var. thyrsiflorus (JM93: no var.)</i>	Blue Blossom	Woody	- - -	-	NP	Bluffs, slopes, canyons, chaparral, coastal scrub, closed-cone-pine forest (Mar–Jun)	
<input type="radio"/> <i>Frangula californica subsp. californica (JM93: Rhamnus)</i>	California Coffee Berry	Woody	- - -	-	NP	Coastal-sage scrub, chaparral, forest, woodland (May–Jul)	
<input type="radio"/> <i>Frangula californica subsp. tomentella (JM93: Rhamnus tomentella subsp. tomentella)</i>	Hoary Coffee Berry	Woody	- - B	-	NP	Chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus crocea</i>	Spiny Redberry	Woody	- - C	-	NP	Coastal-sage scrub, chaparral, woodland (Jan–Apr)	
<input type="radio"/> <i>Rhamnus ilicifolia</i>	Hollyleaf Redberry	Woody	- - C	-	NP	Chaparral, montane forest (Mar–Jun)	
Rosaceae		Rosaceae					
<input type="radio"/> <i>Acaena pinnatifida var. californica</i>	California Acaena	Herbaceous	- - C	-	NP	Coastal grassland, open, rocky slopes (Mar–May)	

Rosaceae	Sorted Alphabetically by Family, Scientific Name					Rosaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Adenostoma fasciculatum</i> var. <i>fasciculatum</i> (JM93: no var.)	Chamise	Woody	- - -	-	NP	Dry slopes, ridges, chaparral (May–Jun)
<input type="radio"/> <i>Amelanchier utahensis</i>	Utah Service-berry	Woody	- - C	-	NP	Open, rocky slopes, canyons, banks of creeks, deserts, conifer forest (Apr–Jun)
<input type="radio"/> <i>Aphanes occidentalis</i>	Lady's Mantle	Herbaceous	- - -	-	NA	Seasonally moist grassland, chaparral, woodland (Mar–May)
<input type="radio"/> <i>Cercocarpus betuloides</i> var. <i>betuloides</i>	Birch-leaf Mountain-mahogany	Woody	- - C	-	NP	Dry, rocky slopes, chaparral (Mar–May)
<input type="radio"/> <i>Cotoneaster pannosus</i> (JM93: <i>C. pannosa</i>)	Silverleaf Cotoneaster	Woody	- - -	M	ZP	Disturbed places, mixed-evergreen forest (May–Jul)
<input type="radio"/> <i>Crataegus monogyna</i>	Hawthorn	Woody	- - -	L	ZP	Naturalized in scattered places (May–Aug)
<input type="radio"/> <i>Drymocallis glandulosa</i> var. <i>glandulosa</i> (JM93: <i>Potentilla glandulosa</i> ssp. <i>glandulosa</i>)	Sticky Cinquefoil	Herbaceous	- - -	-	NP	Gen ± shady or moist areas (May–Jul)
<input type="radio"/> <i>Duchesnea indica</i> var. <i>indica</i> (JM93: no var.)	Mock-strawberry	Herbaceous	- - -	-	ZP	Disturbed areas (May–Aug)
<input type="radio"/> <i>Fragaria vesca</i>	Wood Strawberry	Herbaceous	- - -	-	NP	Gen partial shade in forest (Jan–Jul)
<input type="radio"/> <i>Heteromeles arbutifolia</i>	Christmas Berry / Toyon	Woody	- - -	-	NP	Chaparral, oak woodland, mixed-evergreen forest ((May)Jun–Aug)
<input type="radio"/> <i>Holodiscus discolor</i> var. <i>discolor</i> (JM93: no var.)	Oceanspray	Woody	- - -	-	NP	Moist woodland edges, rocky slopes (May–Aug)
<input type="radio"/> <i>Horkelia californica</i> var. <i>californica</i> (JM93: subsp. <i>californica</i>)	Few-lobed Calif. Horkelia	Herbaceous	- - A1	-	NP	Grassy openings, edges of coastal scrub, especially n slopes (Mar–Sep)
<input type="radio"/> <i>Horkelia californica</i> var. <i>elata</i> (JM93: subsp. <i>dissita</i>)	Many-lobed Calif. Horkelia	Herbaceous	- - A2	-	NP	Shady meadow edges, seasonal streams, open chaparral (Jun–Sep)
<input type="radio"/> <i>Horkelia californica</i> var. <i>frondosa</i> (JM93: subsp. <i>frondosa</i>)	Unlobed California Horkelia	Herbaceous	- - C	-	NP	Coastal scrub, canyons, poison-oak thickets (May–Oct)
<input type="radio"/> <i>Horkelia</i> sp.	Horkelia	Herbaceous	- - -	-	NP	--
<input type="radio"/> <i>Oemleria cerasiformis</i>	Oso Berry	Woody	- - -	-	NP	Chaparral, canyons, streambanks, lowland wet to dry open woodland, coast to shaded conifer forest (Feb–Apr)
<input type="radio"/> <i>Physocarpus capitatus</i>	Pacific Ninebark	Woody	- - C	-	NP	Moist banks, n-facing slopes, mixed-conifer forest (May–Jul)
<input type="radio"/> <i>Potentilla anserina</i> subsp. <i>pacifica</i>	Pacific Cinquefoil	Herbaceous	- - A1	-	NP	Coastal wetlands, often brackish (Mar–Oct)
<input type="radio"/> <i>Prunus cerasifera</i>	Cherry Plum	Woody	- - -	L	ZP	Roadsides, streambanks, chaparral as waif (Feb–Mar)
<input type="radio"/> <i>Prunus dulcis</i>	Almond	Woody	- - -	-	ZP	Canyons, roadsides, grassland (as waif) (Feb–Mar)
<input type="radio"/> <i>Prunus emarginata</i>	Bitter Cherry	Woody	- - C	-	NP	Rocky slopes, canyons, chaparral, mixed-evergreen, conifer forest (Apr–Jun)
<input type="radio"/> <i>Prunus ilicifolia</i> subsp. <i>ilicifolia</i>	Holly-leaved Cherry	Woody	- - C	-	NP	Canyons, slopes, scrubland, woodland (Apr–May)
<input type="radio"/> <i>Prunus subcordata</i>	Sierra Plum	Woody	- - C	-	NP	Mixed-evergreen or conifer forest (Mar–May)

Rosaceae		Sorted Alphabetically by Family, Scientific Name					Rosaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Prunus virginiana</i> var. <i>demissa</i>	Western Choke Cherry	Woody	- - -	-	NP	Rocky slopes, canyons, scrubland, oak/pine woodland (May–Jun)	
<input type="radio"/> <i>Pyracantha angustifolia</i>	Woolly Firethorn	Woody	- - -	L	ZP	Disturbed areas, fencerows, abandoned fields, roadsides (Feb–Jun)	
<input type="radio"/> <i>Rosa californica</i>	California Rose	Woody	- - -	-	NP	Gen ± moist areas in sun, esp streambanks (Feb–Nov)	
<input type="radio"/> <i>Rosa gymnocarpa</i> var. <i>gymnocarpa</i> (JM93: no var.)	Wood Rose	Woody	- - -	-	NP	Common. Gen in shade of forest, scrub ((Feb)Apr–Jul)	
<input type="radio"/> <i>Rosa</i> sp.	Rose	Woody	- - -	-	NP	--	
<input type="radio"/> <i>Rosa spithamea</i>	Coast Ground Rose	Woody	- - C	-	NP	Open forest, chaparral, especially after fire (Apr–Aug)	
<input type="radio"/> <i>Rubus armeniicus</i> (JM93: <i>R. discolor</i>)	Himalayan Blackberry	Woody	- - -	H	ZP	Common. Disturbed areas, roadsides (Mar–Jun)	
<input type="radio"/> <i>Rubus parviflorus</i>	Thimbleberry	Woody	- - -	-	NP	Common; moist semi-shaded areas, esp edges of woodland (Mar–Aug)	
<input type="radio"/> <i>Rubus spectabilis</i>	Salmonberry	Woody	- - A1	-	NP	Moist areas, especially edges of woodland, streambanks (Mar–Jun)	
<input type="radio"/> <i>Rubus ursinus</i>	California Blackberry	Woody	- - -	-	NP	Open, disturbed areas (Mar–Jul)	
Rubiaceae							Rubiaceae
<input type="radio"/> <i>Cephalanthus occidentalis</i> (JM93: var. <i>californicus</i>)	California Button Willow	Woody	- - B	-	NP	Lake, stream edges (May–Sep)	
<input type="radio"/> <i>Galium andrewsii</i> subsp. <i>gatense</i>	Phlox-leaf Serpentine Bedstraw	Herbaceous	- 4.2 *A2	-	NP	Dry, rocky places in serpentine soil, chaparral or open oak/pine woodland (Apr–Jun)	
<input type="radio"/> <i>Galium aparine</i>	Goose Grass	Herbaceous	- - -	-	NA	Grassy, ± shady places (Apr–Jun)	
<input type="radio"/> <i>Galium californicum</i> subsp. <i>californicum</i>	California Bedstraw	Herbaceous	- - -	-	NP	Shady to open places, conifer or mixed forest, chaparral, sea cliffs, hillsides (Mar–Jul)	
<input type="radio"/> <i>Galium murale</i>	Tiny Bedstraw	Herbaceous	- - -	-	ZA	Damp, mossy places, grassy hillsides, dry disturbed areas (Apr–May)	
<input type="radio"/> <i>Galium parisiense</i>	Wall Bedstraw	Herbaceous	- - -	-	ZA	Warm, dry, generally rocky soil to moist areas, coastal-sage scrub, chaparral, grassy hillsides with oaks, roadsides (Apr–Aug)	
<input type="radio"/> <i>Galium porrigens</i> var. <i>porrigens</i>	Climbing Bedstraw	Herbaceous	- - -	-	NP	Among shrubs in chaparral, forest (May–Aug)	
<input type="radio"/> <i>Galium</i> sp.	Bedstraw	Herbaceous	- - -	-	--		
<input type="radio"/> <i>Galium trifidum</i> subsp. <i>columbianum</i> (JM93: var. <i>pacificum</i>)	Trifid Bedstraw	Herbaceous	- - A2	-	NP	Wet places, yellow-pine forest (Jul–Aug)	
<input type="radio"/> <i>Galium triflorum</i>	Sweet-scent Bedstraw	Herbaceous	- - B	-	NP	Damp, shady forest (May–Jul)	
<input type="radio"/> <i>Sherardia arvensis</i>	Field Madder	Herbaceous	- - -	-	ZA	Pastures, disturbed areas, grassland, dry meadows, oak woodland (Mar–Jul)	

Ruppiaceae	Sorted Alphabetically by Family, Scientific Name					Ruppiaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv OL	Habitat (Bloom)	
Ruppiaceae						Ruppiaceae
<input type="radio"/> <i>Ruppia maritima</i>	Straight Ditch-grass	Herbaceous	- - -	- NP	Brackish or saline waters, often coastal, marshes, ponds, sloughs (Mar–Aug)	
Ruscaceae						Ruscaceae
<input type="radio"/> <i>Maianthemum racemosum</i> (JM93: <i>Smilacina racemosa</i>)	Fat False Solomon's Seal	Herbaceous	- - -	- NP	Moist, open woodland, streambanks (Mar–Jul)	
<input type="radio"/> <i>Maianthemum stellatum</i> (JM93: <i>Smilacina stellata</i>)	Starry False Solomon's Seal	Herbaceous	- - -	- NP	Moist woodland, streambanks, open slopes (Apr–Jun)	
Rutaceae						Rutaceae
<input type="radio"/> <i>Ptelea crenulata</i>	Western Hop Tree	Woody	- - B	- NP	Scrub, woodland (Apr–May)	
Salicaceae						Salicaceae
<input type="radio"/> <i>Populus fremontii</i> subsp. <i>fremontii</i>	Fremont Cottonwood	Woody	- - -	- NP	Scattered. Alluvial bottomland, streambanks (Mar–Apr)	
<input type="radio"/> <i>Populus trichocarpa</i> (JM93: <i>P. balsamifera</i> subsp. <i>trichocarpa</i>)	Black Cottonwood	Woody	- - C	- NP	Alluvial bottomland, streambanks (Feb–Apr)	
<input type="radio"/> <i>Salix exigua</i> var. <i>hindsiana</i> (JM93: no var.)	Hinds' Willow	Woody	- - -	- NP	Common. Floodplains, sandy gravel (Apr–May)	
<input type="radio"/> <i>Salix gooddingii</i>	Goodding's Black Willow	Woody	- - B	- NP	Common. Streambanks, marshes, seepage areas, washes, meadows (Mar–Apr)	
<input type="radio"/> <i>Salix laevigata</i>	Red Willow	Woody	- - -	- NP	Common. Riverbanks, seepage areas, lakeshores, canyons (Dec–Jun)	
<input type="radio"/> <i>Salix lasiandra</i> var. <i>lasiandra</i> (JM93: <i>S. lucida</i> subsp. <i>Lasiandra</i>)	Pacific Willow	Woody	- - -	- NP	Wet meadows, lakeshores, riverbanks (May–Jun)	
<input type="radio"/> <i>Salix lasiolepis</i>	Arroyo Willow	Woody	- - -	- NP	Common. Shores, marshes, meadows, etc (Jan–Jun)	
<input type="radio"/> <i>Salix scouleriana</i>	Scouler Willow	Woody	- - A2	- NP	Common. Dry to moist forest, meadows, springs, swamps (Feb–Jun)	
<input type="radio"/> <i>Salix sitchensis</i>	Sitka Willow	Woody	- - -	- NP	Common. Tidal swamps, marshes, springs, streambeds (Mar)	
<input type="radio"/> <i>Salix</i> sp.	Willow	Woody	- - -	- ?P	- -	
Sapindaceae						Sapindaceae
<input type="radio"/> <i>Acer macrophyllum</i>	Big-leaf Maple	Woody	- - -	- NP	Common. Streambanks, canyons (Mar–Jun)	
<input type="radio"/> <i>Acer negundo</i> (JM93: var. <i>californicum</i>)	Box Elder	Woody	- - -	- NP	Streambanks, bottomland. (Mar–Apr)	
<input type="radio"/> <i>Aesculus californica</i>	California Buckeye	Woody	- - -	- NP	Dry slopes, canyons, borders of streams (May–Jun)	

Saururaceae		Sorted Alphabetically by Family, Scientific Name				Saururaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
Saururaceae						Saururaceae	
<input type="radio"/> <i>Anemopsis californica</i>	Yerba Mansa	Herbaceous	- - B	-	NP	Common. Saline or alkaline soil, wet or moist areas, seeps, springs (Mar-Sep)	
Saxifragaceae						Saxifragaceae	
<input type="radio"/> <i>Heuchera micrantha</i>	Small-flower Alumroot	Herbaceous	- - C	-	NP	Moist, rocky banks and cliffs (Apr-Jul)	
<input type="radio"/> <i>Lithophragma affine</i>	Woodland Star	Herbaceous	- - -	-	NP	Open, grassy slopes (Mar-Apr)	
<input type="radio"/> <i>Lithophragma heterophyllum</i>	Hill Starflower	Herbaceous	- - -	-	NP	Shaded slopes (Feb-Jun)	
<input type="radio"/> <i>Lithophragma parviflorum var. parviflorum</i>	Prairie Starflower	Herbaceous	- - B	-	NP	Common. Open areas (Mar-Jul)	
<input type="radio"/> <i>Lithophragma sp.</i>	Woodland Star	Herbaceous	- - -	-	NP	- -	
<input type="radio"/> <i>Micranthes californica (JM93: Saxifraga)</i>	California Saxifrage	Herbaceous	- - -	-	NP	Moist, shady places (Feb-May(Jun))	
<input type="radio"/> <i>Tellima grandiflora</i>	Fringe Cups	Herbaceous	- - B	-	NP	Moist slopes (Apr-Jul)	
<input type="radio"/> <i>Tolmiea diplomenziesii (JM93: T. menziesii)</i>	Pig-a-back Plant	Herbaceous	- - A1	-	NP	Moist streambanks (May-Aug)	
Scrophulariaceae						Scrophulariaceae	
<input type="radio"/> <i>Buddleja davidii</i>	Butterfly Bush	Woody	- - -	-	ZP	Disturbed, generally ± damp soil (May-Sep)	
<input type="radio"/> <i>Limosella acaulis</i>	Southern Mudwort	Herbaceous	- - A2	-	NA	Wet, muddy places, generally fresh water (May-Oct)	
<input type="radio"/> <i>Limosella australis (JM93: L. subulata)</i>	Delta Mudwort	Herbaceous	- 2.1 -	-	ZA	Muddy or sandy intertidal flats, brackish water (Apr)	
<input type="radio"/> <i>Myoporum laetum</i>	New Zealand Myoporum	Woody	- - -	M	ZP	Open areas in grassland, scrub, riparian habitats, generally coastal (Early spring)	
<input type="radio"/> <i>Scrophularia californica (JM93: subsp. californica, floribunda)</i>	California Figwort	Herbaceous	- - -	-	NP	Common; damp places, chaparral, roadsides (Mar-Jul)	
<input type="radio"/> <i>Verbascum thapsus</i>	Woolly Mullein	Herbaceous	- - -	L	ZB	Roadsides, streambanks, disturbed areas (May-Sep)	
Selaginellaceae						Selaginellaceae	
<input type="radio"/> <i>Selaginella bigelovii</i>	Spike-moss	Fern-like	- - A2	-	NP	Open sites, mineral soil on rock outcrops, amid shrubs -	
Simaroubaceae						Simaroubaceae	
<input type="radio"/> <i>Ailanthus altissima</i>	Tree Of Heaven	Woody	- - -	M	ZP	Disturbed areas, grassland, oak woodland, riparian areas (Jun)	
Solanaceae						Solanaceae	
<input type="radio"/> <i>Datura wrightii</i>	Wright Jimson Weed	Herbaceous	- - -	-	NAP	Sandy or gravelly open areas (Apr-Oct)	
<input type="radio"/> <i>Nicotiana attenuata</i>	Coyote Tobacco	Herbaceous	- - A1	-	NA	Open, well-drained slopes (May-Oct)	

Solanaceae		Sorted Alphabetically by Family, Scientific Name				Solanaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
<input type="radio"/> <i>Nicotiana glauca</i>	Tree Tobacco	Woody	- - -	M	ZP	Open, disturbed flats or slopes (Apr–Aug)	
<input type="radio"/> <i>Nicotiana quadrivalvis</i>	Indian Tobacco	Herbaceous	- - B	-	NA	Open, well-drained washes, slopes (May–Oct)	
<input type="radio"/> <i>Petunia parviflora</i>	Wild Petunia	Herbaceous	- - A1	-	NA	Open washes, dry streambeds (Apr–Aug)	
<input type="radio"/> <i>Solanum americanum</i>	Small-flowered Nightshade	Herbaceous	- - -	-	NAP	Open, often disturbed places (Apr–Nov)	
<input type="radio"/> <i>Solanum nigrum</i>	Black Nightshade	Herbaceous	- - -	-	ZA	Disturbed places (Mar–Oct)	
<input type="radio"/> <i>Solanum sp.</i>	Nightshade	Herbaceous	- - -	-	-	-	
<input type="radio"/> <i>Solanum umbelliferum</i>	Blue Witch	Woody	- - -	-	NP	Shrubland, mixed-evergreen forest, woodland (All year)	
<input type="radio"/> <i>Solanum xanti</i>	Purple Nightshade	Herbaceous	- - A1?	-	NP	Shrubland, oak/pine woodland, conifer forest (Feb–Jun)	
Tamaricaceae						Tamaricaceae	
<input type="radio"/> <i>Tamarix aphylla</i>	Athel	Woody	- - -	L	ZP	Uncommon. Washes, roadsides (May–Nov)	
<input type="radio"/> <i>Tamarix parviflora</i>	Smallflower Tamarisk	Woody	- - -	H	ZP	Common. Washes, streambanks, slopes, roadsides (Mar–Apr)	
<input type="radio"/> <i>Tamarix ramosissima</i>	Saltcedar	Woody	- - -	H	ZP	Common. Washes, streambanks (Apr–Aug)	
Themidaceae						Themidaceae	
<input type="radio"/> <i>Brodiaea elegans subsp. elegans</i>	Harvest Brodiaea	Herbaceous	- - -	-	NP	Grassland, meadows, open woodland, chaparral, occasionally serpentine (Apr–Aug)	
<input type="radio"/> <i>Brodiaea terrestris subsp. terrestris</i>	Dwarf Brodiaea	Herbaceous	- - A2	-	NP	Coastal prairie, foothill woodland (Apr–Jul)	
<input type="radio"/> <i>Dichelostemma capitatum subsp. capitatum</i>	Blue Dicks	Herbaceous	- - -	-	NP	Open woodland, scrub, desert, grassland (Mar–Jun)	
<input type="radio"/> <i>Dichelostemma congestum</i>	Fork-toothed Ookow	Herbaceous	- - -	-	NP	Open woodland, grassland (Apr–Jun)	
<input type="radio"/> <i>Dichelostemma sp.</i>	Blue Dicks/ookow	Herbaceous	- - -	-	NP	-	
<input type="radio"/> <i>Muilla maritima</i>	Common Muilla	Herbaceous	- - B	-	NP	Coastal scrub, grassland, woodland, valleys (Mar–Jun)	
<input type="radio"/> <i>Triteleia hyacinthina</i>	Fool's Onion	Herbaceous	- - -	-	NP	Grassland, vernal wet meadows, occ drier slopes (Mar–Jul)	
<input type="radio"/> <i>Triteleia laxa</i>	Ithuriel's Spear	Herbaceous	- - -	-	NP	Common. Open forest, conifer or foothill woodland, grassland on clay soil (Apr–Jun)	
Theophrastaceae						Theophrastaceae	
<input type="radio"/> <i>Samolus parviflorus</i>	Seaside Brookweed	Herbaceous	- - A1	-	NP	Moist sites (Spring–summer)	
Thymelaeaceae						Thymelaeaceae	
<input type="radio"/> <i>Dirca occidentalis</i>	Western Leatherwood	Woody	- 1B.2 *A2	-	NP	Gen n facing slopes, mixed-evergreen forest to chaparral, gen fog belt (Nov–Mar)	

Tropaeolaceae		Sorted Alphabetically by Family, Scientific Name				Tropaeolaceae	
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)	
Tropaeolaceae						Tropaeolaceae	
<input type="radio"/> <i>Tropaeolum majus</i>	Garden Nasturtium	Herbaceous	- - -	-	ZAP	Disturbed areas, moist or shaded areas in coastal scrub, wooded flats or slopes (Mar–Jun)	
Typhaceae						Typhaceae	
<input type="radio"/> <i>Sparganium eurycarpum</i> var. <i>eurycarpum</i> (JM93: subsp. <i>eurycarpum</i>)	Broad-fruited Bur-reed	Grass-like	- - A2	-	NP	Marshes, lakes, ponds, along streams (Jun–Aug)	
<input type="radio"/> <i>Sparganium eurycarpum</i> var. <i>greenii</i> (JM93: <i>S. erectum</i> subsp. <i>stoloniferum</i>)	Erect Bur-reed	Grass-like	- - A1	-	NP	Generally coastal (Jun–Aug)	
<input type="radio"/> <i>Typha angustifolia</i>	Narrow-leaved Cattail	Grass-like	- - -	-	NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (May–Aug)	
<input type="radio"/> <i>Typha domingensis</i>	Southern Cattail	Grass-like	- - -	-	NP	Nutrient-rich freshwater to brackish marshes, wet disturbed places (Jun–Jul)	
<input type="radio"/> <i>Typha latifolia</i>	Broad-leaved Cattail	Grass-like	- - -	-	NP	Unpolluted to nutrient-rich freshwater (brackish) marshes (Jun–Jul)	
<input type="radio"/> <i>Typha</i> sp.	Cattail	Grass-like	- - -	-	NP	--	
Ulmaceae						Ulmaceae	
<input type="radio"/> <i>Ulmus parvifolia</i>	Chinese Elm	Woody	- - -	-	ZP	Streams, springs, wetlands, roadsides, disturbed areas (Aug–Oct)	
Urticaceae						Urticaceae	
<input type="radio"/> <i>Hesperocnide tenella</i>	Western Creeping Nettle	Herbaceous	- - -	-	NA	Moist, shaded areas, often at base of rocks or shrubs, in chaparral, coastal scrub, riparian woodland, mesic oak woodland (Feb–Jun)	
<input type="radio"/> <i>Soleirolia soleirolii</i>	Baby's Tears	Herbaceous	- - -	-	ZA	Damp, shaded places (Mar–Jun)	
<input type="radio"/> <i>Urtica dioica</i> subsp. <i>holosericea</i>	Hoary Nettle	Herbaceous	- - -	-	NP	Meadows, seeps, springs, margins of marshes, streams, lakes, moist areas in chaparral, coastal scrub (Jun–Sep)	
<input type="radio"/> <i>Urtica urens</i>	Dwarf Nettle	Herbaceous	- - -	-	ZA	Disturbed areas, stream banks, shaded areas in grassland, oak woodland, chaparral, coastal-sage scrub, riparian woodland (Jan–Jun)	
Valerianaceae						Valerianaceae	
<input type="radio"/> <i>Centranthus ruber</i>	Red Valerian	Herbaceous	- - -	-	ZP	Disturbed places, rock or wall crevices, roadsides (Apr–Jul)	
<input type="radio"/> <i>Plectritis ciliosa</i> (JM93: subsp. <i>ciliosa</i> , <i>insignis</i>)	Longspur Plectritis	Herbaceous	- - B	-	NA	Common. Open, partly shaded slopes (Mar–Jun)	
<input type="radio"/> <i>Plectritis congesta</i> subsp. <i>brachystemon</i> (JM93: <i>P. brachystemon</i>)	Pale Plectritis	Herbaceous	- - B	-	NA	Common. Coastal bluffs, open, partly shaded slopes (Mar–Jun)	

Valerianaceae		Sorted Alphabetically by Family, Scientific Name				Valerianaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
<input type="radio"/> <i>Plectritis congesta</i> subsp. <i>congesta</i>	Sea Blush	Herbaceous	- - A2	-	NA	Coastal bluffs, open, partly shaded slopes (Mar–Jun)
<input type="radio"/> <i>Plectritis macrocera</i>	Longhorn Plectritis	Herbaceous	- - -	-	NA	Common. Open, partly shaded slopes (Mar–Jun)
Verbenaceae						Verbenaceae
<input type="radio"/> <i>Phyla lanceolata</i>	Fog-fruit	Herbaceous	- - A1	-	NP	Wet places, marshes (May–Nov)
<input type="radio"/> <i>Phyla nodiflora</i> (JM93: var. <i>nodiflora</i>)	Lemon Verbena	Herbaceous	- - -	-	NP	Wet places, pond margins (May–Nov)
<input type="radio"/> <i>Phyla</i> sp.	Verbena / Fog-fruit	Herbaceous	- - -	-	NP	--
<input type="radio"/> <i>Verbena bonariensis</i>	Tall Vervain	Herbaceous	- - -	-	ZAB	Disturbed, often wet places, fields (Jun–Oct)
<input type="radio"/> <i>Verbena lasiostachys</i> var. ?	Western Vervain	Herbaceous	- - -	-	NP	--
<input type="radio"/> <i>Verbena lasiostachys</i> var. <i>lasiostachys</i>	Western Vervain	Herbaceous	- - -	-	NP	Open, dry to wet places (May–Sep)
<input type="radio"/> <i>Verbena lasiostachys</i> var. <i>scabrida</i>	Robust Vervain	Herbaceous	- - -	-	NP	Open, dry to wet places (May–Sep)
Violaceae						Violaceae
<input type="radio"/> <i>Viola adunca</i> var. <i>adunca</i> (JM93: no var.)	Western Dog Violet	Herbaceous	- - A1	-	NP	Vernally moist meadows, damp streambanks, meadow edges in conifer forest, generally shade (Apr–Aug)
<input type="radio"/> <i>Viola douglasii</i>	Douglas' Violet	Herbaceous	- - A1	-	NP	Vernally moist flats, grassy slopes, often on serpentine (Feb–Jul)
<input type="radio"/> <i>Viola glabella</i>	Stream / Smooth Yellow Violet	Herbaceous	- - A2	-	NP	Moist to wet generally shady places in forest, streambanks, etc (Mar–Aug)
<input type="radio"/> <i>Viola pedunculata</i>	Johnny-jump-up	Herbaceous	- - C	-	NP	Open, grassy slopes, hillsides, chaparral, oak woodland, gen full sun (Feb–Apr)
<input type="radio"/> <i>Viola purpurea</i> subsp. <i>purpurea</i>	Mountain Violet	Herbaceous	- - A2	-	NP	In openings or beneath shrubs, generally in yellow-pine forest or higher (Mar–Jul)
<input type="radio"/> <i>Viola purpurea</i> subsp. <i>quercetorum</i>	Oak Violet	Herbaceous	- - A2	-	NP	Dry, grassy or brushy slopes, chaparral, generally below yellow-pine forest (Feb–Jul)
<input type="radio"/> <i>Viola sempervirens</i>	Redwood / Evergreen Violet	Herbaceous	- - A1	-	NP	Shady areas in coastal forest (Jan–Jul)
<input type="radio"/> <i>Viola sheltonii</i>	Shelton's Violet	Herbaceous	- - A2	-	NP	Fir, pine, or oak woodland, rich or gravelly soil (Mar–Jul)
<input type="radio"/> <i>Viola</i> sp.	Violet	Herbaceous	- - -	-	NP	--
Viscaceae						Viscaceae
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>macrophyllum</i> (JM93: <i>P. macrophyllum</i>)	Big-leaf Mistletoe	Herbaceous	- - C	-	NP	On trees other than <i>Quercus</i> (especially <i>Alnus</i> , <i>Fraxinus</i> , <i>Juglans</i> , <i>Platanus</i> , <i>Populus</i> , <i>Robinia</i> , <i>Salix</i>) (Dec–Mar)
<input type="radio"/> <i>Phoradendron serotinum</i> subsp. <i>tomentosum</i> (JM93: <i>P. villosum</i>)	Oak Mistletoe	Herbaceous	- - -	-	NP	Generally on <i>Quercus</i> , rarely on <i>Adenostoma</i> , <i>Arctostaphylos</i> , <i>Rhus</i> , <i>Umbellularia</i> (Jul–Sep)

Vitaceae		Sorted Alphabetically by Family, Scientific Name				Vitaceae
✓ Scientific Name (JM93 if different)	Common Name	Form	Status	Inv	OL	Habitat (Bloom)
Vitaceae						Vitaceae
<input type="radio"/> <i>Parthenocissus vitacea</i>	Woodbine	Woody	- - -	-	NP	Uncommon. Hillsides, thickets, ravines, open woodland, roadsides (Apr–Jun)
<input type="radio"/> <i>Vitis californica</i>	California Wild Grape	Woody	- - -	-	NP	Streamsides, springs, canyons (May–Jun)
<input type="radio"/> <i>Vitis vinifera</i>	Wine Grape	Woody	- - -	-	ZP	Abandoned fields, roadsides (May–Jun)
Woodsiaceae						Woodsiaceae
<input type="radio"/> <i>Athyrium filix-femina</i> var. <i>cyclosorum</i>	Western Lady Fern	Fern-like	- - -	-	NP	Woodland, along streams, seepage area -
<input type="radio"/> <i>Cystopteris fragilis</i>	Brittle Fern	Fern-like	- - C	-	NP	Shady, moist rock crevices, meadows, streamsides -
Zannichelliaceae						Zannichelliaceae
<input type="radio"/> <i>Zannichellia palustris</i>	Horned-pondweed	Herbaceous	- - -	-	NP	Streams, ponds, lakes (Mar–Nov)
Zosteraceae						Zosteraceae
<input type="radio"/> <i>Zostera marina</i>	Eel-grass	Grass-like	- - -	-	NP	Common. Shallow water, bays, estuaries (Mar–Jun)
Zygophyllaceae						Zygophyllaceae
<input type="radio"/> <i>Tribulus terrestris</i>	Puncture Vine	Herbaceous	- - -	-	ZA	Dry, disturbed areas incl roadsides, railways, vacant lots (Apr–Oct)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Acer negundo var. *californicum* (no var.)

Adenostoma fasciculatum (var. *fasciculatum*)

Agoseris grandiflora (var. *grandiflora*)

Agoseris heterophylla (var. *cryptopleura*)

Agrostis viridis (*Polypogon*)

Alisma plantago-aquatica (*A. triviale*)

Amsinckia menziesii var. *intermedia* (*A. intermedia*)

Amsinckia menziesii var. *menziesii* (no var.)

Antirrhinum vexillo-calyculatum ssp. *vexillo-calyculatum* (*A. vexillocalyculatum* subsp. *vexillocalyculatum*)

Arabis breweri var. *breweri* (*Boechera breweri* subsp. *breweri*)

Arabis glabra var. *glabra* (*Turritis glabra*)

Arctostaphylos tomentosa ssp. *crustacea* (*A. crustacea* subsp. *crustacea*)

Aster chilensis (*Symphyotrichum chilense*)

Aster lentus (*Symphyotrichum lentum*)

Aster radulinus (*Eurybia radulina*)

Aster subulatus var. *ligulatus* (*Symphyotrichum subulatum* var. *parviflorum*)

Atriplex joaquiniana (*A. joaquinana*)

Atriplex lentiformis ssp. *lentiformis* (no subsp.)

Atriplex triangularis (*A. prostrata*)

Azolla mexicana (*A. microphylla*)

Baccharis douglasii (*B. glutinosa*)

Baccharis pilularis (subsp. *consanguinea*)

Baccharis salicifolia (subsp. *salicifolia*)

Balsamorhiza macrolepis var. *macrolepis* (no var.)

Blepharizonia plumosa ssp. *plumosa* (no subsp.)

Blepharizonia plumosa ssp. *viscida* (*Blepharizonia laxa*)

Boschniakia strobilacea (*Kopsiopsis*)

Bromus catharticus (var. *catharticus*)

Camissonia boothii ssp. *decorticans* (*Eremothera*)

Camissonia graciliflora (*Tetrapteron graciliflorum*)

Camissonia intermedia (*Camissoniopsis*)

Camissonia micrantha (*Camissoniopsis*)

Camissonia ovata (*Taraxia*)

Cardaria draba (*Lepidium*)

Carduus pycnocephalus (subsp. *pycnocephalus*)

Carex dewayana ssp. *leptopoda* (*C. leptopoda*)

Carex lanuginosa (*C. pellita*)

Carex ovalis (*C. leporina*)

Ceanothus thyrsoiflorus (var. *thyrsoiflorus*)

Centaurium davyi (*Zeltnera*)

Centaurium muehlenbergii (*Zeltnera*)

Cephalanthus occidentalis var. *californicus* (no var.)

Chamomilla suaveolens (*Matricaria discoidea*)

Chenopodium ambrosioides (*Dysphania*)

Chenopodium macrospermum var. *halophilum* (no var.)

Chenopodium multifidum (*Dysphania multifida*)

Chenopodium rubrum (var. *rubrum*)

Chrysothamnus nauseosus ssp. *mohavensis* (*Ericameria nauseosa* var. *mohavensis*)

Cirsium cymosum (var. *cymosum*)

Claytonia parviflora ssp. *parviflora* (subsp. *utahensis*)

Collinsia bartsifolia var. *davidsonii* (var. *stricta*)

Collinsia heterophylla (var. *heterophylla*)

Conyza bilbaoana (*Erigeron sumatrensis*)

Conyza bonariensis (*Erigeron*)

Conyza canadensis (*Erigeron*)

Conyza coulteri (*Laennecia*)

Corallorhiza maculata (var. *maculata*)

Cordylanthus mollis ssp. *mollis* (*Chloropyron molle* subsp. *molle*)

Coronopus didymus (*Lepidium didymum*)

Corylus cornuta var. *californica* (subsp. *californica*)

Cotoneaster pannosa (*C. pannosus*)

Cryptantha decipiens (*C. rattanii*)

Cryptantha intermedia (var. *intermedia*)

Cryptantha muricata (var. *muricata*)

Cryptantha torreyana (var. *pumila*)

Cupressus macrocarpa (*Hesperocyparis*)

Cuscuta salina var. *major* (*C. pacifica* var. *pacifica*)

Cynara cardunculus (subsp. *flavescens*)

Danthonia californica var. *californica* (no var.)

Dicentra chrysantha (*Ehrendorferia*)

Disporum hookeri (*Prosartes*)

Disporum smithii (*Prosartes*)

Draba cuneifolia var. *integrifolia* (no var.)

Duchesnea indica (no var.)

Epilobium pygmaeum (*E. campestre*)

Eragrostis curvula var. *curvula* (no var.)

Erechtites glomerata (*Senecio glomeratus*)

Erechtites minima (*Senecio minimus*)

Eremocarpus setigerus (*Croton*)

Erigeron philadelphicus (var. *philadelphicus*)

Eriogonum umbellatum var. *bahiiforme* (var. *smallianum*)

Eriophyllum lanatum var. *achillaeoides* (var. *achilleoides*)

Eryngium aristulatum var. *aristulatum* (*E. jepsonii*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Erysimum capitatum var. *angustatum* (var. *capitatum*)

Filago californica (*Logfia filaginoides*)

Filago gallica (*Logfia*)

Fritillaria affinis var. *affinis* (no var.)

Galium trifidum var. *pacificum* (subsp. *columbianum*)

Gastridium ventricosum (*G. phleoides*)

Gnaphalium bicolor (*Pseudognaphalium biolettii*)

Gnaphalium californicum (*Pseudognaphalium*)

Gnaphalium canescens ssp. *beneolens*
(*Pseudognaphalium beneolens*)

Gnaphalium canescens ssp. *microcephalum*
(*Pseudognaphalium microcephalum*)

Gnaphalium luteo-album (*Pseudognaphalium luteoalbum*)

Gnaphalium purpureum (*Gamochaeta ustulata*)

Gnaphalium ramosissimum (*Pseudognaphalium*)

Gnaphalium stramineum (*Pseudognaphalium*)

Grindelia camporum var. *camporum* (no var.)

Grindelia hirsutula var. *davyi* (*G. camporum*)

Grindelia hirsutula var. *hirsutula* (no var.)

Guillenia flavescens (*Caulanthus*)

Guillenia lasiophylla (*Caulanthus lasiophyllus*)

Hedera helix ssp. *helix* (no subsp.)

Heliotropium curassavicum (var. *oculatum*)

Hemizonia congesta ssp. *congesta* (subsp. *lutescens*)

Hemizonia corymbosa ssp. *corymbosa* (*Deinandra corymbosa*)

Hemizonia fitchii (*Centromadia*)

Hemizonia lobbii (*Deinandra*)

Hemizonia parryi ssp. *congdonii* (*Centromadia*)

Hemizonia parryi ssp. *parryi* (*Centromadia*)

Hemizonia pungens ssp. *maritima* (*Centromadia pungens* subsp. *pungens*)

Hemizonia pungens ssp. *pungens* (*Centromadia*)

Heracleum lanatum (*H. maximum*)

Herniaria hirsuta ssp. *cinerea* (var. *cinerea*)

Holodiscus discolor (var. *discolor*)

Hordeum jubatum (subsp. *jubatum*)

Horkelia californica ssp. *californica* (var. *californica*)

Horkelia californica ssp. *dissita* (var. *elata*)

Horkelia californica ssp. *frondosa* (var. *frondosa*)

Isopyrum occidentale (*Enemion*)

Iva axillaris ssp. *robustior* (no subsp.)

Juglans californica var. *hindsii* (*J. hindsii*)

Juncus balticus (subsp. *ater*)

Juncus effusus var. *pacificus* (subsp. *pacificus*)

Juncus lesueurii (*J. lescurii*)

Koeleria phleoides (*K. gerardii*)

Lagophylla ramosissima ssp. *ramosissima* (no subsp.)

Lasthenia californica (subsp. *californica*)

Lavatera cretica (*Malva pseudolavatera*)

Lepidium dictyotum var. *dictyotum* (no var.)

Lepidium latipes var. *latipes* (no var.)

Lepidium nitidum var. *nitidum* (no var.)

Lepidium nitidum var. *oreganum* (no var.)

Leptochloa fascicularis (*L. fusca* subsp. *fascicularis*)

Lessingia filaginifolia var. *californica* (*Corethrogyne filaginifolia*)

Lessingia filaginifolia var. *filaginifolia*
(*Corethrogyne filaginifolia*)

Lewisia rediviva (var. *rediviva*)

Leymus condensatus (*Elymus*)

Leymus triticoides (*Elymus*)

Lilaea scilloides (*Triglochin*)

Limosella subulata (*Limosella australis*)

Linanthus acicularis (*Leptosiphon*)

Linanthus ambiguus (*Leptosiphon*)

Linanthus androsaceus (*Leptosiphon*)

Linanthus bicolor (*Leptosiphon*)

Linanthus ciliatus (*Leptosiphon*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus dichotomus (*Linanthus dichotomus* subsp. *dichotomus* or *meridianus*)

Linanthus parviflorus (*Leptosiphon*)

Linaria canadensis (*Nuttallanthus texanus*)

Lolium multiflorum (*Festuca perennis*)

Lolium perenne (*Festuca perennis*)

Lolium temulentum (*Festuca temulenta*)

Lonicera hispidula var. *vacillans* (no var.)

Lotus humistratus (*Acmispon brachycarpus*)

Lotus micranthus (*Acmispon parviflorus*)

Lotus purshianus var. *purshianus* (*Acmispon americanus* var. *americanus*)

Lotus scoparius var. *scoparius* (*Acmispon glaber* var. *glaber*)

Lotus strigosus (*Acmispon*)

Lotus wrangelianus (*Acmispon*)

Luzula comosa (var. *comosa*)

Madia madioides (*Anisocarpus*)

Malacothamnus fasciculatus (var. *nuttallii*)

Marah fabaceus (*M. fabacea*)

Marah oreganus (*M. oregana*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Melilotus alba (*M. albus*)

Melilotus indica (*M. indicus*)

Mentha spicata var. *spicata* (no var.)

Mimulus aurantiacus (var. *aurantiacus*)

Mirabilis californica (*M. laevis* var. *crassifolia*)

Monardella antonina ssp. *antonina* (*M. villosa* subsp. *villosa*)

Monardella douglasii ssp. *douglasii* (no subsp.)

Monardella villosa ssp. *globosa* (subsp. *villosa*)

Montia fontana ssp. *chondrosperma* (no subsp.)

Myrica californica (*Morella*)

Nassella cernua (*Stipa*)

Nassella lepida (*Stipa*)

Nassella pulchra (*Stipa*)

Osmorhiza chilensis (*O. berteroi*)

Oxalis albicans ssp. *pilosa* (*O. pilosa*)

Parvisedum pentandrum (*Sedella pentandra*)

Phacelia ramosissima var. *latifolia* (no var.)

Phacelia ramosissima var. *ramosissima* (no var.)

Phlox gracilis (*Microsteris*)

Phoradendron macrophyllum (*P. serotinum* subsp. *macrophyllum*)

Phoradendron villosum (*P. serotinum* subsp. *tomentosum*)

Phyla nodiflora var. *incisa* (no var.)

Phyla nodiflora var. *nodiflora* (no var.)

Picris echioides (*Helminthotheca*)

Piperia elegans (subsp. *elegans*)

Piptatherum miliaceum (*Stipa miliacea* var. *miliacea*)

Plagiobothrys canescens (var. *canescens*)

Plagiobothrys fulvus (var. *campestris*)

Plectritis brachystemon (*P. congesta* subsp. *brachystemon*)

Plectritis ciliosa ssp. *ciliosa* (no subsp.)

Plectritis ciliosa ssp. *insignis* (no subsp.)

Plectritis congesta (subsp. *congesta*)

Pleuropogon californicus (var. *californicus*)

Pluchea odorata (var. *odorata*)

Poa bulbosa (subsp. *vivipara*)

Polygonum amphibium var. *emersum* (*Persicaria amphibium*)

Polygonum amphibium var. *stipulaceum* (*Persicaria amphibium*)

Polygonum arenastrum (*P. aviculare* subsp. *depressum*)

Polygonum hydropiperoides (*Persicaria*)

Polygonum lapathifolium (*Persicaria lapathifolia*)

Polygonum persicaria (*Persicaria maculosa*)

Polygonum punctatum (*Persicaria punctata*)

Populus balsamifera ssp. *trichocarpa* (*P. trichocarpa*)

Potamogeton pectinatus (*Stuckenia pectinata*)

Potentilla glandulosa ssp. *glandulosa* (*Drymocallis glandulosa* var. *glandulosa*)

Psilocarphus tenellus var. *tenellus* (no var.)

Ranunculus aquatilis var. *capillaceus* (var. *diffusus*)

Ranunculus californicus (var. *californicus*)

Ranunculus canus (var. *canus*)

Ranunculus occidentalis (var. *occidentalis*)

Ranunculus sceleratus (var. *sceleratus*)

Rhamnus californica ssp. *californica* (*Frangula*)

Rhamnus tomentella ssp. *tomentella* (*Frangula californica* subsp. *tomentella*)

Rhus trilobata (*R. aromatica*)

Ribes menziesii (var. *menziesii*)

Rorippa nasturtium-aquaticum (*Nasturtium officinale*)

Rorippa palustris var. *occidentalis* (subsp. *palustris*)

Rosa gymnocarpa (var. *gymnocarpa*)

Rubus discolor (*R. armeniacus*)

Rumex maritimus (*R. fueginus*)

Rumex salicifolius var. *crassus* (*R. crassus*)

Rumex salicifolius var. *salicifolius* (no var.)

Rumex salicifolius var. *transitorius* (*R. transitorius*)

Salicornia europaea (*S. depressa*)

Salicornia subterminalis (*Arthrocnemum subterminale*)

Salicornia virginica (*S. pacifica*)

Salix exigua (var. *hindsiana*)

Salix lucida ssp. *lasiandra* (*Salix lasiandra* var. *lasiandra*)

Sambucus mexicana (*S. nigra* subsp. *caerulea*)

Satureja douglasii (*Clinopodium*)

Saxifraga californica (*Micranthes*)

Scirpus acutus var. *occidentalis* (*Schoenoplectus*)

Scirpus americanus (*Schoenoplectus*)

Scirpus californicus (*Schoenoplectus*)

Scirpus cernuus (*Isolepis*)

Scirpus maritimus (*Bolboschoenus maritimus* subsp. *paludosus*)

Scirpus robustus (*Bolboschoenus*)

Scrophularia californica ssp. *californica* (no subsp.)

Scrophularia californica ssp. *floribunda* (no subsp.)

Senecio breweri (*Packera*)

Senecio mikanioides (*Delairea odorata*)

Silene californica (*S. laciniata* subsp. *californica*)

East Bay Regional Park District

Wild Plant Name Changes

JM93 Scientific Name (JM2 name changes)

Smilacina racemosa (*Maianthemum racemosum*)

Smilacina stellata (*Maianthemum stellatum*)

Solidago californica (*S. velutina* subsp. *californica*)

Solidago canadensis ssp. *elongata* (*S. elongata*)

Sparganium erectum ssp. *stoloniferum* (*S. eurycarpum* var. *greenii*)

Sparganium eurycarpum ssp. *eurycarpum* (var. *eurycarpum*)

Spergula arvensis ssp. *arvensis* (no subsp.)

Spergularia media (var. *media*)

Stachys ajugoides var. *ajugoides* (no var.)

Stachys ajugoides var. *rigida* (*Stachys rigida* var. *quercetorum*)

Streptanthus albidus ssp. *peramoenus* (*S. glandulosus* subsp. *glandulosus*)

Stylomecon heterophylla (*Papaver heterophyllum*)

Taeniatherum caput-medusae (*Elymus*)

Thysanocarpus laciniatus (var. *laciniatus*)

Tolmiea menziesii (*T. diplomenziesii*)

Trifolium albopurpureum var. *albopurpureum* (no var.)

Trifolium albopurpureum var. *dichotomum* (*T. dichotomum*)

Trifolium albopurpureum var. *olivaceum* (*T. olivaceum*)

Trifolium barbigerum var. *andrewsii* (*T. grayi*)

Trifolium barbigerum var. *barbigerum* (no var.)

Trifolium gracilentum var. *gracilentum* (no var.)

Vicia americana var. *americana* (subsp. *americana*)

Vicia ludoviciana var. *ludoviciana* (subsp. *ludoviciana*)

Viola adunca (var. *adunca*)

Vulpia bromoides (*Festuca*)

Vulpia microstachys var. *ciliata* (*Festuca microstachys*)

Vulpia microstachys var. *confusa* (*Festuca microstachys*)

Vulpia microstachys var. *microstachys* (*Festuca microstachys*)

Vulpia microstachys var. *pauciflora* (*Festuca microstachys*)

Vulpia myuros var. *hirsuta* (*Festuca myuros*)

Vulpia myuros var. *myuros* (*Festuca myuros*)

Vulpia octoflora var. *hirtella* (*Festuca octoflora*)

Vulpia octoflora var. *octoflora* (*Festuca octoflora*)

Zigadenus fremontii (*Toxicoscordion*)

East Bay Regional Park District

Changes in Family Names by Genus

Genus	JM93 Family	JM2 Family
<i>Acer</i>	Aceraceae	Sapindaceae
<i>Aesculus</i>	Hippocastanaceae	Sapindaceae
<i>Allium</i>	Liliaceae	Alliaceae
<i>Amaryllis</i>	Liliaceae	Amaryllidaceae
<i>Anagallis</i>	Primulaceae	Myrsinaceae
<i>Antirrhinum</i>	Scrophulariaceae	Plantaginaceae
<i>Asclepias</i>	Asclepiadaceae	Apocynaceae
<i>Athyrium</i>	Dryopteridaceae	Woodsiaceae
<i>Bellardia</i>	Scrophulariaceae	Orobanchaceae
<i>Brodiaea</i>	Liliaceae	Themidaceae
<i>Buddleja</i>	Buddlejaceae	Scrophulariaceae
<i>Calandrinia</i>	Portulacaceae	Montiaceae
<i>Calyptridium</i>	Portulacaceae	Montiaceae
<i>Castilleja</i>	Scrophulariaceae	Orobanchaceae
<i>Chlorogalum</i>	Liliaceae	Agavaceae
<i>Chloropyron</i>	Scrophulariaceae	Orobanchaceae
<i>Claytonia</i>	Portulacaceae	Montiaceae
<i>Collinsia</i>	Scrophulariaceae	Plantaginaceae
<i>Cordylanthus</i>	Scrophulariaceae	Orobanchaceae
<i>Cuscuta</i>	Cuscutaceae	Convolvulaceae
<i>Cymbalaria</i>	Scrophulariaceae	Plantaginaceae

<i>Cystopteris</i>	Dryopteridaceae	Woodsiaceae
<i>Dichelostemma</i>	Liliaceae	Themidaceae
<i>Digitalis</i>	Scrophulariaceae	Plantaginaceae
<i>Emmenanthe</i>	Hydrophyllaceae	Boraginaceae
<i>Eriodictyon</i>	Hydrophyllaceae	Boraginaceae
<i>Eucrypta</i>	Hydrophyllaceae	Boraginaceae
<i>Fremontodendron</i>	Sterculiaceae	Malvaceae
<i>Glaux</i>	Primulaceae	Myrsinaceae
<i>Hydrocotyle</i>	Apiaceae	Araliaceae
<i>Hydrophyllum</i>	Hydrophyllaceae	Boraginaceae
<i>Keckiella</i>	Scrophulariaceae	Plantaginaceae
<i>Kickxia</i>	Scrophulariaceae	Plantaginaceae
<i>Landoltia</i>	Lemnaceae	Araceae
<i>Lewisia</i>	Portulacaceae	Montiaceae
<i>Linaria</i>	Scrophulariaceae	Plantaginaceae
<i>Maianthemum</i>	Liliaceae	Ruscaceae
<i>Mimulus</i>	Scrophulariaceae	Phrymaceae
<i>Montia</i>	Portulacaceae	Montiaceae
<i>Muilla</i>	Liliaceae	Themidaceae
<i>Myoporum</i>	Myoporaceae	Scrophulariaceae
<i>Nemophila</i>	Hydrophyllaceae	Boraginaceae

<i>Nitrophila</i>	Chenopodiaceae	Amaranthaceae
<i>Nuttallanthus</i>	Scrophulariaceae	Plantaginaceae
<i>Pedicularis</i>	Scrophulariaceae	Orobanchaceae
<i>Penstemon</i>	Scrophulariaceae	Plantaginaceae
<i>Phacelia</i>	Hydrophyllaceae	Boraginaceae
<i>Pholistoma</i>	Hydrophyllaceae	Boraginaceae
<i>Romanzoffia</i>	Hydrophyllaceae	Boraginaceae
<i>Ruppia</i>	Potamogetonaceae	Ruppiaceae
<i>Sambucus</i>	Caprifoliaceae	Adoxaceae
<i>Samolus</i>	Primulaceae	Theophrastaceae
<i>Sequoia</i>	Taxodiaceae	Cupressaceae
<i>Spirodela</i>	Lemnaceae	Araceae
<i>Tonella</i>	Scrophulariaceae	Plantaginaceae
<i>Toxicoscordion</i>	Liliaceae	Melanthiaceae
<i>Trientalis</i>	Primulaceae	Myrsinaceae
<i>Trillium</i>	Liliaceae	Melanthiaceae
<i>Triphysaria</i>	Scrophulariaceae	Orobanchaceae
<i>Triphysaria</i>	Scrophulariaceae	Orobanchaceae
<i>Triteleia</i>	Liliaceae	Themidaceae
<i>Veronica</i>	Scrophulariaceae	Plantaginaceae
<i>Viburnum</i>	Caprifoliaceae	Adoxaceae